



Contents lists available at ScienceDirect

Diabetes & Metabolic Syndrome: Clinical Research & Reviews

journal homepage: www.elsevier.com/locate/dsx

Original Article

Documented hypoglycemia is associated with poor short and long term prognosis among patients admitted to general internal medicine departments

Eyal Leibovitz^{a,*}, Israel Khanimov^a, Julio Wainstein^{b,c}, Mona Boaz^d^a Department of Internal Medicine "A" Yoseftal Hospital, Eilat, Israel^b Diabetes Clinic at the Wolfson Medical Center, Holon, Israel^c Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel^d The Department of Nutrition Sciences, Ariel University, Israel

ARTICLE INFO

Article history:

Received 1 July 2018

Accepted 15 July 2018

Keywords:

Diabetes mellitus

Hypoglycemia

1-Year mortality

Length of stay

ABSTRACT

Aim: To study the association of documented hypoglycemia with length of stay, 30-day mortality, and 1-year mortality, among patients with and without diabetes admitted to internal medicine units.

Methods: The electronic medical records of all patients hospitalized in internal medicine departments at E. Wolfson Medical Center, Holon, Israel, between 1/1/2010 and 31/12/2013, were reviewed. Data extracted included all glucose measurements (performed using an institutional blood glucose monitoring system). Patients were considered hypoglycemic if at least one hypoglycemic event was recorded. Regression analysis was used to assess the association between documented hypoglycemia and length of stay, 30-day and one-year mortality. Age, sex, reason for admission, and the Charlson comorbidity index were entered as covariates, and the most conservative model was developed.

Results: The study population included 45,272 patients (mean age 68.9 ± 17.8 years, 49.4% males, 21.0% had diabetes mellitus). The rate of hypoglycemia in the total study population was 7.5% (16.8% among DM patients, 6.0% among patients without diabetes, $p < 0.001$). Patients with documented hypoglycemia had a longer length of hospital stay (9.3 ± 18.7 vs. 3.1 ± 6.4 days, $p < 0.001$), as well as higher risk for both 30-day (23.7% vs. 7.0%, $p < 0.001$) and 1-year mortality (41.6% vs. 15.3%, $p < 0.001$). Cox regression analysis showed that hypoglycemia significantly increased risk death at one year (HR 2.436, 95% CI 2.298–2.582, $p < 0.001$) independent of age, sex, the Charlson comorbidity index, DM status and reason for admission.

Conclusion: Documented hypoglycemia is associated with prolonged length of hospital stay and increased risk for both 30-day and 1-year mortality, regardless of diabetes mellitus status.

© 2018 Published by Elsevier Ltd on behalf of Diabetes India.

1. Introduction

In the community settings, strict glucose control among diabetes mellitus patients is associated with increased risk for hypoglycemia [1]. This in turn leads to increased morbidity and mortality [2,3] and increased treatment costs [4]. In addition, it has been shown that hypoglycemia occurring in the hospital setting is also associated with increased risk of in-hospital mortality in several hospitalized patient populations. Among diabetes patients hospitalized in general medical departments, hypoglycemia was

associated with increased mortality [5], and prognosis was also correlated to hypoglycemia severity and number of hypoglycemic events [6].

Hypoglycemia has also been documented among non-diabetes patients. This is more frequent among intensive care unit (ICU) patients where hypoglycemia and hyperglycemia were shown to be associated with worst prognosis [7]. Similar results were found among non-critically ill patients, where glucose variability, rather than hyperglycemia or hypoglycemia per-se, was found to be associated with increased length of stay and higher mortality rates [8]. The etiology of this finding is unknown, but the occurrence of hypoglycemia among non-diabetes patients is considered to be associated with disease severity [9].

For diabetes patients, the occurrence of hyper and/or

* Corresponding author. Medicine "A", Yoseftal Hospital, Derech Yotam, Eilat, Israel.

E-mail address: eyallei1@post.tau.ac.il (E. Leibovitz).

hypoglycemia is considered to be mostly attributed to under/over treatment. However, this is not the case for non-diabetes patients. The hypothesis of this study is that hyper and hypoglycemia during hospitalization is detrimental to health, and carries similar short and long term risk, regardless of etiology and/or diabetes mellitus status. The aim of this study is to analyze the association of hyperglycemia and hypoglycemia incidence and short and long term prognosis among patients admitted to internal medicine departments.

2. Methods

2.1. Study design: historical prospective analysis of medical records

Study population: For this study, electronic medical records of all patients discharged from internal medicine units of Edith Wolfson Medical Center, a 700-bed tertiary government medical center, between January 1st 2010 and December 31st 2013, were reviewed. For patients with several admissions, only the first (index) admission was included. Extracted data included patient demographics, co-morbidities, laboratory information, and length of hospital stay and date of death when applicable. Diabetes mellitus (DM) status was defined as the presence of DM as a diagnosis in the discharge letter (ICD9 code 250 and sub codes) and/or glucose lowering medications recorded upon hospital admission. Laboratory information included all glucose measurements performed during the hospitalization period, using an institutional blood glucose monitoring system (IGMS), which consisted of a point-of-care, automated glucometer, and an interactive database [10].

Glucose control and hypoglycemia definitions: A hypoglycemic event was defined as any glucose measurement equal or below 70 mg/dL (3.89 mmol/L), regardless of symptoms or treatment. Patients were considered hypoglycemic if they had at least one documented hypoglycemic event. The number of hypoglycemic events was also recorded for each patient, and an average glucose during the stay was calculated. Patients were considered “controlled” if the average glucose during stay was lower than 180 mg/dL, regardless of the number of glucose measurements.

Statistical analysis: Data were analyzed using SPSS ver. 21.0 (IBM Inc., USA). Continuous variables were compared between patient groups using the *t*-test for independent samples, or ANOVA with Bonferonni post-hoc analysis as appropriate. Associations between nominal variables were assessed using the chi-square test. Logistic regression analysis was used to examine the effect of the number of hypoglycemic events on 30-day and, separately, 1-year mortality. General linear modeling was used to predict length of hospital stay. Cox regression analysis was used to model 30-day and 1-year survival. All tests are two-sided and considered significant at $p < 0.05$.

3. Results

3.1. Hypoglycemic events among diabetes and non-diabetes patients

Table 1 shows patient demographics, co-morbidities and baseline laboratory data across diabetes mellitus and hypoglycemia groups. During the acquisition period, 45,272 patients (mean age 68.9 ± 17.8 years, 49.4% males) were discharged from the hospital. Of the patients discharged, 9507 (21.0%) had DM.

The average glucose levels during stay for diabetes patient were 182 ± 63 mg/dL, and 57.7% were considered controlled with an average below 180 mg/dL. For non-diabetes patients, the average glucose levels during stay were 133 ± 46 mg/dL, and 88.8% were controlled ($p < 0.001$ between diabetes and non-diabetes results).

Rate of hypoglycemia was 7.5% of all patients. There were more individuals with documented hypoglycemic events among patients with vs. without diabetes: 16.8% vs. 6.0%, $p < 0.001$. Among patients with any hypoglycemic event, the mean number of events was 2.2 ± 2.6 (median 1 event). There was no clear association between the average glucose and the rate of hypoglycemia. Rate of hypoglycemia was the highest among the first and fifth quintiles of glucose control, regardless of diabetes mellitus status.

In general, hypoglycemic patients were older (73.8 ± 15.5 vs. 68.4 ± 17.9 years, $p < 0.001$), had higher rates of hypoalbuminemia upon admission (58.8% vs. 28.2%, $p < 0.001$) lower cholesterol (150 ± 50 vs. 169 ± 48 mg/dL, $p < 0.001$), lower hemoglobin (11.7 ± 2.2 vs. 12.6 ± 2.0 g/dL, $p < 0.001$) and higher white blood cell counts (12.2 ± 2.0 vs. $10.2 \pm 6.3 \times 10^3$ cells per mm^3 , $p < 0.001$). Acute infection was more prevalent as a reason for admission among patients with hypoglycemia (43.7% vs. 35.1%, $p < 0.001$). In addition, hypoglycemic patients were sicker, as indicated by the Charlson comorbidity index (1.9 ± 2.0 vs. 1.0 ± 1.6 arbitrary units, $p < 0.001$). They had higher rates of congestive heart failure (12.6% vs. 8.4%, $p < 0.001$), chronic renal failure (18.9% vs. 7.4%, $p < 0.001$) and dementia (7.6% vs. 4.0%, $p < 0.001$).

3.2. Short- and long-term prognosis of hypoglycemia

In the total study cohort, the mean length of hospital stay was 3.5 ± 8.2 days (median 2 days, Q1–Q3 0–4 days). Rates of 30-day and 1-year mortality were 8.3% and 17.3% respectively. Table 2 shows the length of hospital stay, 30-day and 1-year mortality rates according to hypoglycemia and diabetes mellitus status.

Regardless of diabetes mellitus status, patients with documented hypoglycemia during the hospitalization period had a significantly longer length of hospital stay compared to patients without hypoglycemia (9.3 ± 18.7 vs. 3.1 ± 6.4 , $p < 0.001$). Compared to patients without hypoglycemic events, patients with at least one hypoglycemic event had significantly greater 30-day mortality (7.0% vs. 23.7%, $p < 0.001$) and 1-year mortality (15.3% vs. 41.6%, $p < 0.001$).

A logistic regression model was used to examine the effects of the number of hypoglycemic events, and the glucose control status on 30-day, and 1-year mortality. This model showed that an increased number of hypoglycemic events significantly increased odds of both 30-day (OR 1.205, 95% CI 1.173–1.238, $p < 0.001$) and 1-year mortality (OR 1.370, 95% CI 1.327–1.414, $p < 0.001$).

In addition, age (OR 1.071, 95% CI 1.068–1.075, $p < 0.001$) also showed significantly increased odds of 30-day mortality. However, positive diabetes mellitus status (OR 0.733, 95% CI 0.662–0.810, $p < 0.001$) and positive glucose control status (OR 0.443, 95% CI 0.406–0.484, $p < 0.001$) all significantly reduced odds of 30-day mortality. Regarding 1-year mortality: Age (OR 1.076, 95% CI 1.073–1.079, $p < 0.001$), male sex (OR 1.154, 95% CI 1.090–1.221, $p < 0.001$) and the Charlson comorbidity index (OR 1.346, 95% CI 1.325–1.367, $p < 0.001$) all significantly increased odds of 1-year mortality. However, positive diabetes mellitus status (OR 0.797, 95% CI 0.737–0.862, $p < 0.001$) and positive glucose control during stay (OR 0.546, 95% CI 0.508–0.586, $p < 0.001$) were both significantly associated with reduced 1-year mortality.

In the total study cohort, general linear modeling showed that age, glucose control, and hypoglycemia occurrence significantly increased length of hospital stay, while sex did not. Findings were robust in analyses performed separately among patients with and without diabetes. In these models, both age and hypoglycemia occurrence significantly increased the length of hospital stay. However, when plotting the estimated marginal means for length of stay according to hypoglycemia, glucose control and diabetes mellitus status, it appears that hypoglycemia was a key parameter

Download English Version:

<https://daneshyari.com/en/article/10148551>

Download Persian Version:

<https://daneshyari.com/article/10148551>

[Daneshyari.com](https://daneshyari.com)