Accepted Manuscript

A High-Order interpolation for the Finite Volume Method: the Coupled Least Squares Reconstruction

Florian Haider, Bernard Courbet, Jean-Pierre Croisille


 PII:
 S0045-7930(18)30622-4

 DOI:
 https://doi.org/10.1016/j.compfluid.2018.09.009

 Reference:
 CAF 4004

To appear in: Computers and Fluids

Received date:24 March 2016Revised date:6 September 2018Accepted date:9 September 2018



Please cite this article as: Florian Haider, Bernard Courbet, Jean-Pierre Croisille, A High-Order interpolation for the Finite Volume Method: the Coupled Least Squares Reconstruction, *Computers and Fluids* (2018), doi: https://doi.org/10.1016/j.compfluid.2018.09.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Highlights

- A new cubic reconstruction for the finite volume method is proposed.
- The new method is presented in the general d-dimensional case.
- This reconstruction uses the Least Squares method in a coupled way.
- An algorithm efficiently calculating the successive derivatives is presented.
- Numerical illustrations with several hyperbolic problems in gas dynamics are given in dimensions 1, 2 and 3.

A CERTIFICATION AND SCALL

Download English Version:

https://daneshyari.com/en/article/10226401

Download Persian Version:

https://daneshyari.com/article/10226401

Daneshyari.com