




Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Synthesis and biological evaluation of aryloxyacetamide derivatives as neuroprotective agents

Yan Zhong^{a,†}, Yi Xu^{b,†}, Ai-Xia Zhang^b, Xiao-Feng Li^b, Zhao-Ying Xu^b, Ping Li^c, Bin Wu^{b,*}

^aSchool of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, China

^bSchool of Pharmacy, Nanjing Medical University, Nanjing 211166, China

^cSchool of Basic Medical Sciences, Nanjing Medical University, Nanjing 210029, China

ARTICLE INFO

Article history:

Received 22 December 2015

Revised 16 March 2016

Accepted 25 March 2016

Available online xxx

Keywords:

Aryloxyacetamide

Neuroprotective activity

Cerebral ischemic stroke

Caspase-3

ABSTRACT

A series of new aryloxyacetamide derivatives **10a–s** and **14a–m** are designed and synthesized. Their protective activities against the glutamate-induced cell death were investigated in differentiated rat pheochromocytoma cells (PC12 cells). Most compounds exhibited neuroprotective effects, especially for **10m**, **10r**, **14b** and **14c**, which showed potential protection of PC12 cells at three doses (0.1, 1.0, 10 μ M). MTT assay, Hoechst 33342/PI double staining, and high content screening (HCS) revealed that pretreatment of the cells with **10m**, **10r**, **14b** and **14c** has significantly decreased the extent of cell apoptosis in a dose-dependent manner. The results of western blot analysis demonstrated these compounds suppressed apoptosis of glutamate-induced PC12 cells via caspase-3 pathway. These compounds can be lead compounds for further discovery of neuroprotective agents for treating cerebral ischemic stroke. Basic structure–activity relationships are also presented.

© 2016 Published by Elsevier Ltd.

Stroke is the second commonest cause of death and leading cause of adult disability worldwide. Over two-thirds of stroke deaths worldwide are in developing countries.^{1,2} In China, stroke is the second cause for mortality in all diseases. Particularly, ischemic strokes account for 60–80% of these strokes.³ The pathological mechanisms of ischemic strokes are complex and not fully understood, but it is generally accepted that a pathological release of glutamate from neurons plays a central role in mediating subsequent neuronal cell injury and death.⁴ Excessive excitatory transmission can be transformed into an implement of neuronal destruction resulting in CNS disorder, such as cerebral ischemia, hypoxia, autoimmune, Alzheimer's, Parkinson's diseases, and so on.⁵

So far, treatment options for stroke-related brain damage are very limited. Most pharmacological agents have focused on mechanisms that occur in acute stage of stroke, such as restoration of blood flow with antithrombosis and thrombolytic therapy, or reducing the effects of ischemia by neuroprotective therapy. Unfortunately, thrombolytic therapy can only be given to highly selected patients.⁶ Therefore, there is an urgent need for effective neuroprotective agents to treat stroke-related brain damage.

Natural products are the single most productive source of lead molecules for development as clinically useful drugs for human disorders.⁷ It has been reported that the cinnamon extract has vasodilative, antithrombotic, anti-ulcerous, and anti-allergic action.⁸ It is believed that the extracts (aqueous and/or organic solvent extraction) would provide different compositions of phytochemicals (such as cinnamic acid, cinnamaldehyde, proanthocyanidins), and these are responsible for the above effect.⁹

Encouraged by these observations and in continuation of our ongoing research program, we designed and synthesized several cinnamide derivatives to explore their neuroprotective properties.¹⁰ We found that cinnamide scaffold often affords neuroprotective compounds (Fig. 1). Especially the compound **NY-308** (**1**), which has the (*E*)-*p*-methoxycinnamoyl moiety, exhibited good neuroprotection in vitro PC12 cells and in vivo rat focal cerebral ischemic animal model.^{10,11} Structurally, the (*E*)-*p*-methoxycinnamoyl moiety in **NY-308** was believed to play a very important role in its activity. To further elucidate the structure–activity relationship, we designed and synthesized a novel series of phenoxyacetamide derivatives using –OCH₂CO– connecting bridge as the surrogate replacing –CH=CHCO– moiety of cinnamide. Furthermore, we replaced diphenylmethylpiperazine of **NY-308** with benzylpiperazine and changed substituent groups of benzene rings to find new chemical entities with better neuroprotective activity (Fig. 2).

* Corresponding author. Tel./fax: +86 025 8686 8467.

E-mail address: wubin@njmu.edu.cn (B. Wu).

† These authors contributed equally to this work.




Figure 1. Structures of new cinnamide derivatives.




Figure 2. Structure exploration strategy based on compound NY-308 (1).

Therefore, a total of 32 target compounds were designed and synthesized. Their *in vitro* neuroprotective activities in the cell injury induced by glutamate, apoptosis assays, and the inhibition of caspase-3 were evaluated. Herein, the synthesis and preliminary biological evaluation of these compounds were reported.

Synthetic route for substituted diphenylmethylpiperazine analogs **10a–s** was depicted in Scheme 1. Commercially available substituted aromatic phenols **2a–g** were transformed into the desired intermediates **3a–g**, via reacting with ethyl chloroacetate in acetonitrile. Subsequent hydrolysis of **3a–f** with 10% sodium hydroxide gave carboxylic acids **4a–f**, which were chloridized with oxalyl chloride to give acyl chlorides **5a–f**. The synthesis of other key intermediates **9a–d** commenced with the reduction reactions of benzophenones **6a–d** under sodium borohydride. Subsequent chlorination of benzhydrols **7a–d** with SOCl_2 , followed by alkylation with anhydrous piperazine, afforded diphenylmethylpiperazines **9a–d**. The target compounds **10a–s** were finally obtained by acylation of various substituted diphenylmethylpiperazines with the corresponding acyl chloride under mild conditions (at RT in acetone, and with triethylamine as base).

The second series of benzylpiperazine compounds **14a–m** were conveniently synthesized as outlined in Scheme 2. Substituted aryloxy acetyl piperazine intermediates **11a–g**, which were obtained by aminolysis of substituted ethyl aryloxyacetates **3a–g** with anhydrous piperazine, were used to react with benzyl bromides **13a–d** to directly provide the target compounds **14a–m**. The key intermediates benzyl bromides **13a–d**, were prepared by free-radical bromination of commercially available substituted toluenes **12a–d** with little excess of *N*-bromosuccinimide (NBS).

In order to study the potential neuroprotective activities of the title compounds, a preliminary screening was performed

investigating neuroprotection on impairment induced by glutamine (Glu) in differentiated PC12 cells,^{12,13} as evaluated by MTT assay.^{14–16} The results are showed in Table 1. The vast majority of compounds tested exhibited protection of PC12 cells against glutamate-induced cell death, indicating that its bioactivity remained after introduction of the $-\text{O}-\text{CH}_2-\text{CO}-$ connecting bridge into structures. Potency and toxicity were highly sensitive to structural variations. Remarkably, compounds **10j**, **10k**, **10m**, **10r**, **14b**, **14c**, **14f**, **14i**, and **14j** showed good neuroprotective activity for all three test concentrations (0.1, 1.0, 10 μM) (protection >20%). From Table 1, it was found that the cumulative addition of the compounds **10m**, **10r**, **14b**, and **14c** (0.1–10 μM) caused concentration-dependent neuroprotective effects with the maximal effect observed at 10 μM . Compounds **10j**, **10k**, **14d**, and **14j** showed a pattern of increased protection with increasing concentrations (0.1–1 μM) in terms of cell protection. Compounds **10n**, **14f**, **14h** and **14k** were observed to have the highest protection at the lowest concentrations of 0.1 μM (cell protection: 45.45%, 35.28%, 53.34% and 39.86%, respectively). Compounds **10a**, **10b**, **10f**, **10g**, **10i**, **10l**, and **14e** were observed to have the highest protection at the highest concentrations of 10 μM (cell protection: 25.64%, 30.24%, 21.17%, 56.98%, 80.50%, 50.34% and 25.10%, respectively). Though many of derivatives showed interesting neuroprotective effects, unfortunately, some of them possess unexpected cytotoxicity toward PC12 cells. Moreover, some compounds have limited solubility in cell culture medium, which complicates interpretation of these negative results.

Derivatives of I and II were substituted with different functional groups to study of the substituent variability influence on the biological activity and find new chemical entities with better neuroprotective activity. Based on lipophilic and electronic

Download English Version:

<https://daneshyari.com/en/article/10590228>

Download Persian Version:

<https://daneshyari.com/article/10590228>

[Daneshyari.com](https://daneshyari.com)