Accepted Manuscript


Biphasic changes in angiopoietin-like 8 level after laparoscopic sleeve gastrectomy and type 2 diabetes remission during a 1-year follow-up

Eman Al-Shawaf Ph.D., Ebaa Al-Ozairi M.D., Fahad Al-Asfar M.D., Shaima Al-Beloushi M.Sc., Surbhi Kumari M.Sc., Jaakko Tuomilehto MD, MA, PhD., Hossein Arefanian Ph.D.

 PII:
 S1550-7289(18)30301-0

 DOI:
 10.1016/j.soard.2018.05.026

 Reference:
 SOARD 3392



Received date:23 January 2018Revised date:22 May 2018Accepted date:23 May 2018

Please cite this article as: Eman Al-Shawaf Ph.D., Ebaa Al-Ozairi M.D., Fahad Al-Asfar M.D., Shaima Al-Beloushi M.Sc., Surbhi Kumari M.Sc., Jaakko Tuomilehto MD, MA, PhD., Hossein Arefanian Ph.D., Biphasic changes in angiopoietin-like 8 level after laparoscopic sleeve gastrectomy and type 2 diabetes remission during a 1-year follow-up, *Surgery for Obesity and Related Diseases* (2018), doi: 10.1016/j.soard.2018.05.026

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Highlights

- 1. ANGPTL8 elevation is associated with T2D, with a minimal BMI effect
- 2. Increase in ANGPTL8 post-LSG was exclusive to participants with T2D
- 3. Increase in ANGPTL8 post-LSG is independent of the increase in %EWL and %EBMIL
- 4. Unique biphasic pattern of change in ANGPTL8 post-LSG in participants with T2D

ACERTIN

Download English Version:

https://daneshyari.com/en/article/11018527

Download Persian Version:

https://daneshyari.com/article/11018527

Daneshyari.com