

Accepted Manuscript

Solitary fibrous tumor of thyroid: a case report with review of literature

Mohammad Ghasemi Rad, Kevin Yuqi Wang, Shilpa Jain, Christie M. Lincoln



PII: S0899-7071(18)30251-1
DOI: doi:[10.1016/j.clinimag.2018.09.011](https://doi.org/10.1016/j.clinimag.2018.09.011)
Reference: JCT 8547

To appear in: *Clinical Imaging*

Received date: 6 August 2018
Revised date: 13 September 2018
Accepted date: 17 September 2018

Please cite this article as: Mohammad Ghasemi Rad, Kevin Yuqi Wang, Shilpa Jain, Christie M. Lincoln , Solitary fibrous tumor of thyroid: a case report with review of literature. Jct (2018), doi:[10.1016/j.clinimag.2018.09.011](https://doi.org/10.1016/j.clinimag.2018.09.011)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Solitary fibrous tumor of thyroid: a case report with review of literature

Mohammad Ghasemi Rad¹, Kevin Yuqi Wang¹, Shilpa Jain², Christie M. Lincoln³

1. Resident Physician, PGY-3, Department of Radiology, Baylor College of Medicine, Houston, TX, USA

2. Assistant Professor, Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, USA

3. Assistant Professor, Department of Radiology, Baylor College of Medicine, Houston, TX, USA

Corresponding Author:

Christie M. Lincoln

Assistant Professor, Department of Radiology, Baylor College of Medicine, Houston, TX, USA

Email: Christie.Lincoln@bcm.edu

ABSTRACT

Solitary fibrous tumors are rare spindle cell tumors of mesenchymal origin most commonly found in the pleura and peritoneum. In contrast, extra-serosal locations are rare. We present a 41-year-old female with a progressively enlarging left-sided neck mass of one-year duration with subsequent rapid-onset growth over two months, and was pathologically confirmed to be a benign solitary fibrous tumor of the thyroid gland.

Download English Version:

<https://daneshyari.com/en/article/11021866>

Download Persian Version:

<https://daneshyari.com/article/11021866>

[Daneshyari.com](https://daneshyari.com)