

# Accepted Manuscript

## EXPERIMENTAL REFERRED PAIN EXTENDS TOWARDS PREVIOUSLY INJURED LOCATION – AN EXPLORATIVE STUDY

Thorvaldur Skuli Palsson , Shellie A. Boudreau ,  
Hans Jørgen Krebs , Thomas Graven-Nielsen

PII: S1526-5900(18)30184-6  
DOI: [10.1016/j.jpain.2018.04.018](https://doi.org/10.1016/j.jpain.2018.04.018)  
Reference: YJPAI 3586



To appear in: *Journal of Pain*

Received date: 29 January 2018  
Revised date: 30 April 2018  
Accepted date: 30 April 2018

Please cite this article as: Thorvaldur Skuli Palsson , Shellie A. Boudreau , Hans Jørgen Krebs , Thomas Graven-Nielsen , EXPERIMENTAL REFERRED PAIN EXTENDS TOWARDS PREVIOUSLY INJURED LOCATION – AN EXPLORATIVE STUDY, *Journal of Pain* (2018), doi: [10.1016/j.jpain.2018.04.018](https://doi.org/10.1016/j.jpain.2018.04.018)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- It is unclear if a recovered pain condition leaves traces of higher sensitivity of pain mechanisms
- Pain is more frequently referred to the previous area of nociceptive activity
- The pain area in the previously painful area is enlarged compared with controls
- The ability to dampen pain via endogenous inhibition seems improved after recovery from pain
- The findings shed a light on the mechanisms involved in recovery from musculoskeletal pain

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/11022857>

Download Persian Version:

<https://daneshyari.com/article/11022857>

[Daneshyari.com](https://daneshyari.com)