

Accepted Manuscript

Gray matter volume differences between affective and non-affective first episode psychosis: A review of Magnetic Resonance Imaging studies

A. Calvo , G. Delvecchio , A.C. Altamura , J.C. Soares ,
P. Brambilla

PII: S0165-0327(17)32317-0
DOI: [10.1016/j.jad.2018.03.008](https://doi.org/10.1016/j.jad.2018.03.008)
Reference: JAD 9649

To appear in: *Journal of Affective Disorders*

Received date: 8 November 2017
Revised date: 22 January 2018
Accepted date: 14 March 2018

Please cite this article as: A. Calvo , G. Delvecchio , A.C. Altamura , J.C. Soares , P. Brambilla , Gray matter volume differences between affective and non-affective first episode psychosis: A review of Magnetic Resonance Imaging studies, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.03.008](https://doi.org/10.1016/j.jad.2018.03.008)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- *This review summarizes MRI studies exploring brain deficits in both NA-FEP and A-FEP patients.*
- *The evidence suggested that the two diagnostic groups showed common and distinct pattern of GM deficits.*
- *This review highlights the importance of identifying neural biomarkers of FEP, which may ultimately be useful for more targeted treatments.*

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/11025350>

Download Persian Version:

<https://daneshyari.com/article/11025350>

[Daneshyari.com](https://daneshyari.com)