



Frequency of bloodstream infection in febrile neutropenic patients, experience from a developing country



Basit Siddiqui^{a,*}, Rabeea Azmat^b, Shiyam Sunder Tikmani^g, Shumaila Rafi^c, Beenish Syed^d, Muhammad Tahir Khan^e, Hareem Rehman^f, Saleemullah Paracha^a

^a Department of Medicine, Abbasi Shaheed Hospital, Karachi, 74700, Pakistan

^b Department of Medicine, Aga Khan University Hospital, Karachi, Pakistan

^c Department of Medicine, Al-Tibri Medical College and Hospital, Karachi, Pakistan

^d Department of Medicine, Liaquat National Hospital, Karachi, Pakistan

^e School of Public Health, Dow University of Health Sciences, Karachi, Pakistan

^f Medical College, The Aga Khan University, Karachi, Pakistan

^g Department of Community Health Sciences, Aga Khan University, Karachi, Pakistan

ARTICLE INFO

Keywords:

Fever
Neutropenia
Staphylococcus
E. coli
Pseudomonas
Klebsiella

ABSTRACT

Introduction: About 20% of febrile neutropenic patients are presented with bloodstream infection which is a leading cause of mortality among these patients. Awareness of the locally prevalent pathogens and their susceptibility pattern is important for proper treatment of infection which improves survival in these patients. The objective of this study was to determine the frequency of bloodstream infection in patients with febrile neutropenia admitted in Abbasi Shaheed Hospital.

Methods: This cross-sectional study was conducted in the Medical wards in Abbasi Shaheed Hospital, Karachi from 21-2-2016 to 20-8-2017. Patients of over 15 years of age of either gender with severe neutropenia were included in this study. Patients with noninfectious cause of fever, or fever prior to neutropenia were excluded. Patients meeting inclusion criteria were enrolled after taking informed consent. About 2–5 ml blood was collected under aseptic measures in Bactac culture bottles. Blood samples were sent to lab within 30 min of collection. Antimicrobial sensitivity testing of all isolates was performed on diagnostic Sensitivity test plates by Kerby – Bauer Method. Patients were referred back to treating physician if blood culture is positive. The data was analyzed using SPSS version 20.

Results: Of 200 patients, the mean age was 25.8 ± 5.7 years, 120 (60%) were male with male to female ratio of 1.5:1, 89 (44.5%) had low grade fever (Temp. $\leq 102^\circ\text{C}$) and 111 (55.5%) had high grade fever (Temp. $> 102^\circ\text{C}$). The frequency of *staphylococcus aureus* was 16%, *E. coli* was 14.5%, *Pseudomonas* 8.5% and *Klebsiella* 15.5%

Conclusions: It is concluded that the frequency of *staphylococcus aureus* was 16%, *E. coli* was 14.5%, *Pseudomonas* 8.5% and *Klebsiella* 15.5%

1. Introduction

Patients with febrile neutropenia are more susceptible to life threatening bacterial infections due to lack of inflammatory response [1]. Hence, febrile neutropenia should be dealt as an emergency and administration of empirical antibiotics may improve survival in these patients [2]. According to American college of physicians and society of critical care medicine which include axillary temperature $> 38\text{C}$, heart rate > 90 /mainland respiratory rate > 20 /min. Absolute neutrophil count is $500/\text{mm}^3$ The reported risk of bloodstream infection in patients with febrile neutropenia is between 11 and 38% [3].

Staphylococcus aureus was the most common isolate among the Gram-positive organisms (53%), while *Pseudomonas aeruginosa*, *Escherichia coli*, and *Klebsiella pneumoniae* were the 3 leading Gram-negative isolates (42%) [4]. Mortality rate due to gram-negative pathogens was 18% and gram-positive pathogen was 5% [5,6].

Compared to previous reports, the pattern of bacterial isolates and their resistance to antibiotics has changed over the past years. Gram-positive organisms predominated in the 50s and 60s, gram negative later, the emergence of multi-drug resistant aerobic rods in 90s and a shift back to gram positive in last two decades in some regions has created problems of effective therapy [7,8].

* Corresponding author. Department of Medicine, Abbasi Shaheed Hospital, Karachi, 74700, Pakistan.

E-mail address: dr.basit.siddiqui@gmail.com (B. Siddiqui).

<https://doi.org/10.1016/j.amsu.2018.09.004>

Received 2 April 2018; Received in revised form 20 August 2018; Accepted 9 September 2018

2049-0801/© 2018 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Management of patients with febrile neutropenic is complex and involves careful consideration of multiple factors like primary underlying etiology, duration, and severity of the neutropenia, Improved broad-spectrum antibiotic agents, combined with improved supportive care, have improved the prognosis for most patients with severe neutropenia. A study showed more than 30 years ago that there was an inverse relationship between the absolute neutrophil count and life threatening infections [9,10]. A delay of more than 48 h in the administration of appropriate antibiotics may result in a mortality rate of approximately 50% [10].

Identification of the locally prevalent pathogens and their susceptibility pattern is important, before putting neutropenic patients on empiric antimicrobial regimens, Bacterial infections are the major cause of morbidity and mortality among neutropenic patients. Thus, the objective of the study was to determine the frequency of bloodstream infection responsible for blood stream infection (BSI) in febrile neutropenic patients admitted in xxxxxx Hospital so that appropriate antibiotic therapy should be started to reduce mortality.

2. Material and methods

This cross sectional study was conducted in Department of Medicine, xxxxxxxx Hospital from 21-February-2016 to 20-August-2017. Department of Medicine is a 120 bedded medical unit admitting approximately X patients per day.

Patients of age over 15 years with moderate to severe neutropenia due to infection (viral, bacterial, protozoal and fungal) or immune neutropenias or hypersplenism or hematological and non-hematological malignancies or vitamin B12, folate deficiency or radiation or chemotherapy or hemodialysis were included in this study. Patients of age < 15 years developing fever due to noninfectious causes as after 6–24 h of transfusion of blood products or drug infusion or had fever prior to neutropenia or those who did not give written consent were excluded from the study. Patients from medical unit were selected after the approval of the ethical board review committee of hospital and College of Physicians and Surgeons. Patients who fulfilled inclusion criteria were enrolled in the study after taking informed consent. The data were collected from all neutropenic patients (defined as, an absolute neutrophil count (ANC) of less than 1500/ml and running fever). Fever was defined as a single oral temperature $\geq 38.3^\circ\text{C}$ (101°F) for at least 1 h or a temperature of $\geq 38^\circ\text{C}$ or 100.4°F on two or more occasions within 12-24-hour period. Four to five ml of venous blood was taken under aseptic technique and is collected in a Bactec blood cultures bottle. Blood culture was sent to laboratory within 20 min of collection. Antimicrobial sensitivity testing of all isolates was performed on diagnostic sensitivity test plates by Kerby – Bauer Method; *Staphylococcus aureus* was identified as round grape like colonies under microscopy; *Pseudomonas aeruginosa*, as rod shaped bacteria with unipolar motility; *E. coli*, as rod shaped, 2 mm long and 0.5 μm in diameter and *Klebsiella pneumoniae* was identified as small rod shaped structures inoculated in MacConkey culture. Data were recorded in a structured questionnaire. Patients were referred back to treating physician if blood culture is positive for further assessment and treatment.

The sample size of 200 was calculated assuming prevalence of neutropenia of 7.2%, bond on error of estimation of 4% and confidence level of 95%. Data analysis was performed through SPSS Version-20. Mean and SD was calculated for age, and temperature. Frequency and percentages was calculated for various bacterial agents which were *staphylococcus aureus*, *E.coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. Stratification was done based on age and gender to see effect on outcome variable through Chi-square test and p-value of < 0.05 was considered significant.

3. Results

A total of 200 patients were enrolled in this study. Mean age was

Table 1
Characteristics of patients.

Age	
Mean \pm SD	25.8 \pm 5.7 years
Sex	
Male	80 (40%)
Female	120 (60%)
Temperature	
Low grade (102 F)	89 (44.5%)
High grade (> 102 F)	111 (55.5%)
Pathogens	
<i>Staphylococcus aureus</i>	32 (16%)
<i>E. coli</i>	29 (14.5%)
<i>Pseudomonas aeruginosa</i>	17 (8.5%)
<i>Klebsiella pneumoniae</i>	31 (15.5%)

25.8 \pm 5.7 years, 120 (60%) were male with male to female ratio of 1.5:1, 89 (44.5%) had low grade fever (Temp. 102 C) and 111 (55.5%) had high grade fever. The frequency of *Staphylococcus aureus* was 16%, *E. coli* was 14.5%, *Pseudomonas aeruginosa* 8.5% and *Klebsiella pneumoniae* was 15.5% (Table 1) (see Table 2 and Table 3).

Stratified analysis showed: Mean age of *Staphylococcus aureus* positive cases was 25.2 \pm 4.7 years compared to 25.9 \pm 5.8 years in *Staphylococcus aureus* cases (p-0.49) and among male patients 19.2% had *Staphylococcus aureus* cases compared to 11.3% in female cases (p-0.16); mean age of *E. coli* positive cases was 28.2 \pm 6.1 years compared to 25.5 \pm 5.5 years in *E. coli* negative cases (p-0.02) and among male patients 14.2% had *E. coli* positive cases compared to 15% in female cases (p-0.51); Mean age of *Pseudomonas aeruginosa* positive cases was 25.6 \pm 3.6 years compared to 25.8 \pm 5.8 years in *Pseudomonas aeruginosa* negative cases (p-0.82) and among male patients 7.5% had *Pseudomonas aeruginosa* positive cases compared to 10% in female cases (p-0.35) and mean age of *Klebsiella pneumoniae* positive cases was 25.9 \pm 4.6 years compared to 25.8 \pm 5.8 years in *Klebsiella pneumoniae* negative cases (p-0.93) and among male patients 14.2% had *Klebsiella pneumoniae* positive cases compared to 17.5% in female cases (p-0.32).

4. Discussion

Febrile neutropenia is the development of fever, often with other signs of infection, in a patient with neutropenia, an abnormally low number of neutrophil granulocytes (a type of white blood cell) in the blood. The term neutropenic sepsis is also applied, although it tends to

Table 2
Stratified analysis of pathogens by age.

	STAPHYLOCOCCUS AUREUS		P-value
	Positive	Negative	
Mean age	25.2 \pm 4.7	25.9 \pm 5.8	0.49
	PSEUDOMONAS AERUGINOSA		P-value
	Positive	Negative	
Mean age	25.6 \pm 3.6	25.8 \pm 5.8	0.82
	E. COLI		P-value
	Positive	Negative	
Mean age	28.2 \pm 6.1	25.5 \pm 5.5	0.02
	KLEBSIELLA PNEUMONAE		P-value
	Positive	Negative	
Mean age	25.9 \pm 4.6	25.8 \pm 5.8	0.93

Download English Version:

<https://daneshyari.com/en/article/11030521>

Download Persian Version:

<https://daneshyari.com/article/11030521>

[Daneshyari.com](https://daneshyari.com)