

X. ISTERH CONFERENCE
Review

Update on the possible nutritional importance of silicon



Forrest H. Nielsen

U.S. Department of Agriculture, Agricultural Research Service,¹ Grand Forks Human Nutrition Research Center, Grand Forks, ND, USA

ARTICLE INFO

Keywords:

Silicon
Trace elements
Bone
Connective tissue
Collagen

ABSTRACT

Convincing evidence that silicon is a bioactive beneficial trace element continues to accumulate. The evidence, which has come from human, animal, and in vitro studies performed by several laboratories, indicate that silicon in nutritional and supra nutritional amounts promotes bone and connective tissue health, may have a modulating effect on the immune or inflammatory response, and has been associated with mental health. A plausible mechanism of action for the beneficial effects of silicon is the binding of hydroxyl groups of polyols such that it influences the formation and/or utilization of glycosaminoglycans, mucopolysaccharides, and collagen in connective tissue and bone. In addition, silicon may affect the absorption, retention or action of other mineral elements (e.g., aluminum, copper, magnesium). Based on findings from both animal and human experiments, an intake of silicon of near 25 mg/d would be a reasonable suggestion for an adequate intake that would assure its nutritional benefits. Increased intakes of silicon through consuming unrefined grains, certain vegetables, and beverages and cereals made from grains should be recognized as a reasonable dietary recommendation.

Published by Elsevier GmbH.

Contents

Introduction	379
Silicon, bone and connective tissue	380
Silicon and immune and inflammatory response	380
Silicon and mental health	381
Plausible silicon mechanism of action	381
Silicon metabolism	381
Silicon status indicators	381
Beneficial intakes of silicon	381
Conclusion	381
Conflict of Interest	382
References	382

Introduction

Silicon is nutritionally essential for some lower forms of life [1]. Silicon has a structural role in diatoms, radiolarians, and some sponges. Diatoms, which are unicellular microscopic plants, have an absolute requirement for silicon as monomeric silicic acid for normal cell growth. Silicon also may be essential for

some higher plants (e.g., rice). Because silicon deprivation has not been shown to interrupt the life cycle in mammals, or to have a defined biochemical function, silicon is not generally accepted as an essential nutrient for higher animals and humans. However, for over 40 years, reports about silicon having beneficial, especially on connective tissue and bone formation, in higher animals and humans have appeared. Initial experiments performed in the 1970s used supra nutritional supplemental amounts of silicon (100 and 500 mg/kg diet) to prevent abnormalities in animal models fed low-silicon diets of questionable nutritional quality based on growth data. The silicon supplementation alleviated abnormal bone structure and strength in chicks and rats; abnormal bone cartilage characterized by decreased hexosamine in chicks; and decreased

E-mail address: forrest.nielsen@ars.usda.gov

¹ The U.S. Department of Agriculture, Agricultural Research Service, Northern Plains Area, is an equal opportunity/affirmative action employer and all agency services are available without discrimination.

collagen and prolyhydroxylase activity in skull bone from cultured chick embryos [1,2]. Concern about dietary quality resulted in the question of whether the supra nutritional supplements were alleviating abnormalities caused by a silicon deprivation or induced by some other sub-optimal dietary factor.

Silicon, bone and connective tissue

Experiments performed since 2000 have indicated that only nutritional amounts of silicon are needed to prevent bone, hexosamine, and collagen metabolism abnormalities, similar to, but of less magnitude than those reported in the 1970s, in animal models fed apparently nutritionally adequate diets low in silicon. In three studies, silicon supplementation of 10 or 35 mg/kg diet containing <2.0 mg/kg silicon, was used to show that the silicon deprivation in rats decreased collagen formation [3] and femur hydroxyproline concentration [4] and increased urinary helical peptide excretion [5]. Another study provided drinking water containing 53.2 µg/g silicon to rats fed diets containing 3.2 µg/g silicon for 26 weeks [6]. The silicon deprivation reduced bone growth plate thickness and increased chondrocyte density. In mice, a silicon supplement of 50 µg/g to a diet containing a soluble silicon content of 0.2 µg/g increased femur dry and ash weights, calcium and hydroxyproline contents, and alkaline phosphatase activity, and decreased femur tartrate-resistant acid phosphatase and urinary excretion of hydroxyproline [7]. Silicon supplementation also increased femur strength and stiffness, and bone marrow mRNA expression related to osteoblastogenesis.

Recent *in vitro* studies also indicate that silicon promotes bone formation. Orthosilicic acid at physiological concentrations was found to stimulate collagen type 1 synthesis in human osteoblast-like cells and enhance osteoblastic differentiation in culture [8]. Silicon in silica-based bioactive glass and ceramics has been implicated in the *in vivo* efficiency of bone engineering implants through involvement in osteoblast proliferation and differentiation, type 1 collagen synthesis, and apatite formation [9,10].

Supra nutritional amounts of silicon also have beneficial effects on bone in ovariectomized rats. A silicon supplement of 500 mg/kg diet for 30 d increased longitudinal growth and mineral content of the femur [11]. In another study, ovariectomized rats were fed a calcium-deficient AIN-93M diet and compared to rats fed the same diet supplemented with 500 mg/kg silicon as sodium metasilicate for 10 weeks [12]. The silicon supplementation significantly increased bone mineral density of the femur and tibia and C-telopeptide type 1 collagen levels in serum. Choline-stabilized orthosilicic acid orally supplemented at a dose of 1 mg/kg body weight daily for 30 weeks was found to partially prevent femoral bone loss in aged ovariectomized rats [13]. Another study with aged (17 weeks) ovariectomized rats found that daily oral administration of 20 mg silicon/kg body weight significantly increased femur and tibia bone mineral density [14].

Recent epidemiological studies have indicated that nutritional intakes of silicon are beneficial for bone health in humans. Dietary silicon was positively associated with bone mineral density in four hip sites of men and premenopausal women in the Framingham Offspring cohort of 1251 men and 1596 women [15]. Large differences of up to 10% were found between the highest (>40 mg/d) and lowest (<14 mg/d) quintiles of silicon intake. Also in the Framingham Offspring Cohort, the beneficial effect of a moderate consumption of beer on hip and spine bone mineral density was associated with silicon in the beer [16]. In addition, silicon intake was positively associated with bone mineral density at the femur neck in late premenopausal women and postmenopausal women on hormone replacement therapy in the Aberdeen Prospective Osteoporosis Screening Study [17]. The lowest quartile of silicon intake was 16 mg/d and highest quartile was 31.5 mg/d. Quartile of

energy-adjusted silicon intakes were negatively associated with urinary markers of bone resorption (pyridinoline and deoxypyridinoline crosslinks) and positively associated with the serum N-terminal propeptide of type 1 collagen, a marker of bone formation.

Although animal, *in vitro*, and epidemiological findings indicate increased dietary silicon is associated with improved bone mass or density and bone turnover markers, human silicon supplementation or intervention studies have found bone mass improvements only in a small number of subjects, only a limited number of positive bone turnover marker changes, and just subjective changes in skin and hair. In a review, it was reported that a 28 mg/d silicon supplement for 12 weeks increased spine bone mineral density by 2.5% in six women with low bone mass [18]. In support of this report was a double-blind, placebo-controlled study completed by 136 women supplemented daily with 1 g calcium and 20 µg vitamin D₃, daily supplements of 3, 6, or 12 mg silicon as choline-stabilized orthosilicic acid, or a placebo for 12 months [19]. The 6 and 12 mg doses of silicon significantly increased the bone formation marker of type 1 collagen at 12 months. Another study that used choline-stabilized orthosilicic acid as a supplement involved 50 women with photo-damaged skin [20]. A supplement of 10 mg/d silicon for 20 weeks improved the photo-damaged skin surface and mechanical properties, and decreased hair and nail brittleness.

In contrast to the limited positive reports, is a study involving 17 postmenopausal women with low bone mass that were randomized to drinking either purified water or artesian water containing 86 mg/L silicon for 12 weeks [21]. The silicon-rich artesian water significantly increased urinary silicon excretion but the purified water did not. Bone turnover markers procollagen type 1, N-terminal propeptide, bone specific alkaline phosphatase, and osteocalcin in serum were not significantly affected within or between groups.

The limited supplementation findings assert the need for randomized controlled trials of silicon supplementation to establish whether achieving a specific dietary intake of this element results in benefits for bone and connective tissue health. These trials should involve a significant number of participants that could be stratified by serum silicon and with supplementation long enough to have measurable effects on changes in bone or connective tissue health.

Silicon and immune and inflammatory response

In 1988, it was suggested that silicon had a regulatory role in the cell cycle of lymphocytes because monomethylsilanetriol at an optimal concentration of 10 mg/L silicon in culture media stimulated peripheral lymphocyte proliferation and decreased lymphoblast proliferation [22]. This possible effect of silicon in nutritional amounts received little attention until 2002 when it was reported that mitogen-induced DNA synthesis of splenic T-lymphocytes from silicon deprived (2.3 mg/kg diet) compared to silicon supplemented (35 mg/kg diet) rats was increased when the rats were fed normal adequate dietary arginine but decreased when fed a supplemental 5 g/kg diet [23]. Since then, findings from two other studies with rats have suggested that silicon supplementation (35 mg/kg diet) vs. silicon-deprivation (2.8 and 1.9 mg/kg diet) might affect the immune or inflammatory response. When injected with type II collagen to induce a long-term (four weeks) inflammatory response, circulating lymphocyte counts were higher and neutrophil counts were lower in silicon-deprived than-supplemented rats [24]. On the other hand, silicon deprivation did not affect acute-phase (2 h) inflammatory markers changes induced by the injection of the endotoxin lipopolysaccharide [25]. However, the endotoxin increased the liver and femur concentrations of silicon in silicon-deprived rats but not in silicon-supplemented rats, which suggests a relation between silicon and the inflammatory or immune response. Further research is required

Download English Version:

<https://daneshyari.com/en/article/1226450>

Download Persian Version:

<https://daneshyari.com/article/1226450>

[Daneshyari.com](https://daneshyari.com)