



International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015)

Does processing have a considerable effect on the nutritional and functional properties of Mung bean (*Vigna radiata*)?

S.D. Chandrasiri^a, R. Liyanage^b, J.K. Vidanarachchi^a, P.Weththasinghe^a, B.C. Jayawardana^{a*}

^aDepartment of Animal Science, Faculty of Agriculture, University of Peradeniya, Peradeniya, 20400, Sri Lanka

^bInstitute of Fundamental Studies, Hanthana road, Kandy, 20000, Sri Lanka

Abstract

Mung bean is an important grain legume which is rich in nutrients and other bioactive compounds with many beneficial physiological effects. However, the effect of processing on properties of mung bean has not been widely studied in Sri Lanka. In the current study, effect of processing (boiling and sprouting) on some nutritional and functional properties of mung bean was investigated in comparison with raw mung bean. Mung bean seeds (*Vigna radiata*), variety MI6 were obtained from the Government Seed Center, Dambulla, Sri Lanka. Proximate composition, gross energy, insoluble dietary fibre (IDF) and soluble dietary fibre (SDF) contents, total phenolic content (TPC), total flavonoid content (TFC), α -amylase inhibition capacity, DPPH and ABTS scavenging activities were determined. Crude Protein (CP) content of boiled mung bean was significantly ($p < 0.05$) lower than that of raw mung bean while it was significantly ($p < 0.05$) higher in sprouted mung bean than raw mung bean. Crude fat content (CF) was significantly ($p < 0.05$) lower in sprouted mung bean when compared to raw mung bean. IDF contents in raw mung bean was significantly ($p < 0.05$) higher than that in sprouted and boiled mung bean whereas, SDF content in boiled and sprouted mung bean were significantly ($p < 0.05$) higher than that in raw mung bean. processing significantly ($p < 0.05$) reduced antioxidant activity and total phenol content compared to raw mung bean. However, flavonoids could only be detected in sprouted mung bean. Further, α -amylase inhibition in sprouted mung bean was significantly ($p < 0.05$) higher than that in boiled mung bean. In conclusion, processing (boiling and sprouting) influences the CP content, CF content, dietary fibre content, antioxidant capacity, α -amylase inhibition and fermentation ability in mung bean.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015).

* Corresponding author. Tel.: +9 471 8585388.

E-mail address: baranaj2000@yahoo.com

Keywords: Mung bean; Processing method; Functional properties; Dietary fibre; *In vitro* study

Main text

1. Introduction

Mung bean [*Vigna radiata* (L.)Wilczek] is one of the important legume crops widely cultivated in Asia¹. From the 5.5 million ha of world mung bean production, about 90% is in Asia and among that, India is the biggest producer where about 2.99 million ha are cultivated⁵. Although mung bean is widely known for its fibre, mineral and protein, at present it is considered not only as a rich source of nutrients but also a source of other bioactive compounds with many beneficial physiological effects such as antioxidant, antidiabetic, anticholesteromic and anticancer effect in controlling and preventing various metabolic diseases². Previous studies have shown that processing alters the nutritional and functional properties of food². Some processing methods can increase and some can decrease the nutritional and functional properties of food. Therefore a great attention should be paid not only on what is eaten but also on the way of preparing it. Mung bean is processed into various forms such as sprouted, cooked and boiled before consumption. However, the effect of processing on nutritional and functional properties of mung bean has not been widely studied in Sri Lanka. Thus, the present study was carried out to investigate the nutritional and functional properties such as antioxidant effect, α - amylase inhibitory activity, dietary fibre and fermentation ability of boiled mung bean (BMB) and sprouted mung bean (SMB) in comparison with raw mung bean (RMB) using *in vitro* techniques.

2. Materials and methods

Mung bean (*Vigna radiata*) seeds, variety MI6 were obtained from the Government Seed Center, Dambulla, Sri Lanka, which was cultivated in Yala season, 2014. The seeds were then cleaned and soaked overnight at room temperature (28°C) before preparing boiled (BMB) and sprouted (SMB) mung beans. Beans were boiled for 30 minutes at 60°C for the preparation of BMB and to prepare SMB, beans were germinated for 48 h in folded papers. Raw, boiled and sprouted seeds were air dried and oven dried at 60°C until a constant weight was obtained and ground to get a fine consistency. Powders were stored in sealed air-tight plastic containers in a refrigerator at 5°C until analysis.

Proximate composition of mung bean was analyzed according to the AOAC procedure and gross energy content was determined using the bomb calorimeter. Insoluble dietary fibre (IDF) and soluble dietary fibre (SDF) contents were also determined on dry matter basis³. Water extract of dried mung bean samples were prepared and antioxidant capacity was determined using DPPH and ABTS assays and total phenolic content (TPC) and total flavonoid content were determined by Folin-ciocalteu and Aluminium chloride method, respectively. Antidiabetic effect was determined by α -amylase inhibitory assay and microbial fermentation characteristics were measured using unadapted microorganisms obtained from swine caecum. In the present study, hydrogen and carbon dioxide gases were measured which are by products of microbial fermentation to determine fermentation characteristics of mung bean.

Data were analyzed by one-way analysis of variance (ANOVA) using the General Linear Model (GLM) procedure of SAS. Significant differences among means were separated by the Duncan's Multiple Rang Test (DMRT). Differences at $p < 0.05$ were considered as significant.

3. Results, discussion, conclusion and recommendation

The gross energy in RMB, BMB and SMB samples were not significantly different ($p > 0.05$). Crude Protein (CP) content in SMB was significantly higher ($p < 0.05$) than that in RMB whereas it was significantly lower ($p < 0.05$) in BMB. Further, the crude fat content (CF) was significantly lower ($p < 0.05$) in SMB compared to RMB (Table 1). Insoluble dietary fibre content in RMB powder was significantly ($p < 0.05$) higher than that of BMB and SMB powder whereas, soluble dietary fibre content in BMB and SMB powder was significantly ($p < 0.05$) higher than that in RMB powder (Table 1). Total phenolic content (TPC) in RMB was significantly higher ($p < 0.05$) than

Download English Version:

<https://daneshyari.com/en/article/1266237>

Download Persian Version:

<https://daneshyari.com/article/1266237>

[Daneshyari.com](https://daneshyari.com)