



Original Research Article

Renal carnitine excretion following abstinence after chronic drinking



Alina Kępką^{a,*}, Sylwia Chojnowska^b, Rafał Śnitko^a, Krzysztof Zwierz^c,
Napoleon Waszkiewicz^d

^a Department of Biochemistry, Radioimmunology and Experimental Medicine, The Children's Memorial Health Institute, Warsaw, Poland

^b Medical Institute, College of Computer Science and Business Administration, Łomża, Poland

^c Medical College of the Universal Education Society, Łomża, Poland

^d Department of Psychiatry, Medical University of Białystok, Białystok, Poland

ARTICLE INFO

Article history:

Received 8 June 2015

Accepted 17 November 2015

Available online 8 December 2015

Keywords:

Abstinence

Alcohol

Carnitine

Fatty acids

Urinary excretion

ABSTRACT

Purpose: Carnitine participates in the metabolism of lipids and cognitive activity. Excessive consumption of alcohol disturbs renal tubular canalicules, that increases urinary excretion of carnitine and its esters.

The study evaluates restoration of the urinary free- and total carnitine as well as acylcarnitine excretion after chronic drinking and during the 49-days of controlled abstinence.

Materials/methods: In 32 patients (6♀; 26♂), 26–60 years old, 2–30 years of alcohol dependence: 75–700 g of pure alcohol (166 ± 94 g) of alcohol daily consumption, 2–360 (35 ± 67) days of intoxication and 1.25 ± 0.8 days of abstinence at admission, we determined urinary free (FC) and total carnitine (TC) as well as acylcarnitine (AC) and acylcarnitine/free carnitine ratio (AC/FC) at admission (T0), after 30 (T30) and 49 (T49) days of the controlled abstinence.

Results: At T0 excretion of FC, TC and AC as well as AC/FC ratio were significantly higher as compared to the control group. After 30- and 49-days of abstinence, excretion of FC and TC decreased to the level of control group with an exception of the AC and AC/FC ratio at T30 that remained significantly increased.

Conclusion: 30 days for the FC and TC and 49 days of abstinence for the AC and AC/FC ratio was sufficient to normalize urinary excretion of the carnitines.

© 2015 Medical University of Białystok. Published by Elsevier Sp. z o.o. All rights reserved.

1. Introduction

World Health Organization estimates that 140 million people worldwide suffer from the alcoholism. Alcoholic disturbance of the body metabolism [1–3] is increased due to malnutrition that is a common finding in chronic alcoholics [4,5]. Normally, free fatty acids are re-esterified with glycerol to form triacylglycerols or they enter mitochondria for β -oxidation [6]. One of the major consequences of alcohol ingestion is an excessive production of NADH, which reduces glucose synthesis and fatty acids oxidation [7,8]. Alcoholism causes structural and functional damage of many body organs including kidney [9]. Renal injury may be associated with ethanol-induced changes in the membrane composition of renal tubules and lipid peroxidation in epithelial cell membranes [10–12].

Chronic alcohol dependence is associated with an excessive excretion of the L-carnitine and its fatty acid esters into the urine

[11,13]. L-carnitine (2-hydroxy-4-trimethylammonium butyrate) is a small hydrophilic molecule synthesized in the liver, brain and kidneys from protein-bound lysine and methionine. L-carnitine participates in the transport of long chain fatty acids from cytoplasm into the mitochondrial matrix for their oxidation. It takes part in the intracellular decomposition and excretion of branched-chain ketoacids, improves cognitive abilities in neurodegenerative diseases and provides acetyl groups for the acetylcholine synthesis [10,14,15]. Demand for the carnitine is covered by endogenous synthesis (25%) and by diet (75%) [14,16]. Carnitine is not degraded in the human body but it is excreted with urine. Kidney plays a major role in the homeostasis of carnitine [13,17,18]. Proximal renal tubules are the intra-renal site of carnitine acylation and regulators of the blood and/or urinary carnitine acylation [19]. In health, tubular reabsorption of free carnitine, a process facilitated by the active transport of carnitine and its short-chain esters by the OCTN2 transporter on the renal brush border membranes, reaches 96–99% [17,20] and is higher than that for carnitine esters [20,21].

An important question is if and when the renal carnitine excretion recovery is possible after heavy drinking period. Therefore, the aim of this study was to evaluate urinary excretion

* Corresponding author at: Department of Biochemistry, Radioimmunology and Experimental Medicine, The Children's Memorial Health Institute, Al. Dzieci Polskich 20, 04-730 Warsaw, Poland. Tel.: +48 22 815 77 52; fax: +48 22 815 13 13.
E-mail address: a.kepka@czd.pl (A. Kępką).

of the free-(FC), total-(TC) and acyl-(AC) carnitine in persons chronically addicted to alcohol before and during controlled abstinence.

2. Material and methods

2.1. Patients

The study group consisted of 32 patients from the Department of Detoxification and Therapy for Alcohol Dependence in Choroszcz Psychiatric Hospital, Poland, 6 women and 26 men, aged from 26 to 60 years (44.6 ± 8.9 years). Alcohol dependence ranged from 2 to 30 years (14.5 ± 7.9 years). Amount of alcohol consumed ranged from 75 to 700 g/day of pure alcohol (166.7 ± 94.3 g/day). Intoxication lasted 3–360 (35.6 ± 67.1) days. At admission, abstinence from alcohol lasted 1.25 ± 0.8 days. Patients with alcoholic liver disease (steatosis, inflammation, fibrosis and cirrhosis) were excluded from the study. During abstinence the patients were hydrated, balanced with electrolytes and glucose for 2.4 ± 0.6 days and treated with the following drugs: diazepam for 8.6 ± 4.6 days, hydroxyzinum for 3.6 ± 2.7 days, haloperidolum for 4.8 ± 2.5 days, carbamazepine for 14.7 ± 5.4 days, promazine for 11.4 ± 10.8 days, tiazolidyncarboxylic acid for 5–36 days and NSAIDs (mostly ketoprofen) for 2–8 days.

The control group consisted of 18 healthy adult volunteers, aged from 22 to 60 years (8 women and 10 men) remaining on the overall diet with an occasional drinking. Prior to collection of the urine samples all persons from the control group did not consume alcohol for at least one week.

2.2. Carnitine determination

In the group of alcoholics urinary FC and TC were determined at admission to hospital (T0), after 30 (T30) and 49 days (T49) of the abstinence. Carnitine excretion was expressed as $\mu\text{mol/g}$ creatinine [18].

Samples of the urine collected during 12-h were centrifuged for 10 min at $2000 \times g$ using 5702R Eppendorf AG centrifuge, Hamburg, Germany, and kept at -86°C until measurements. Urinary free carnitine was determined by enzymatic method of Cederblad et al. [22], which is based on the reaction of free carnitine with acetyl-CoA catalyzed by CAT (carnitine acetyltransferase). Free carnitine reacts with acetyl moiety of acetyl-CoA releasing CoA-SH, which is determined by reaction with 5,5'-dithiobis-2-nitrobenzoic acid

(DTNB). Increase of the absorbance at 412 nm was measured using Hitachi UV/VIS Spectrophotometer – Model U-2900, Tokio, Japan. For the determination of total carnitine, 100 μL of centrifuged urine was added to 10 μL 1 mol/L KOH solution, mixed and incubated at 56°C for 1 h (for hydrolysis of the carnitine esters) and finally neutralized to $\text{pH} \sim 7.0$ with 2 μL 5 mol/L HCl, and assayed for the free carnitine as described above. Acylcarnitine (AC) level was calculated by subtracting the concentration of free carnitine from the total carnitine. Acylcarnitine/free carnitine ratio (AC/FC) was calculated as total carnitine minus free carnitine/free carnitine, according to Schmidt-Sommerfeld et al. [23] and Seccombe et al. [24].

Creatinine in the urine was measured by Jaffe's method in modification of Larsen [25]. The absorbance of chromogen was measured at a 450 nm on Creatinine Analyzer 2 (Beckman, Munich, Germany).

2.3. Statistical analysis

The data are expressed as a mean \pm SD and were analyzed by the Statistica version 10.0 (Statsoft, Cracow, Poland). Independent groups were compared using Student's *t*-test or Mann–Whitney *U*-test depending on their distribution, that was assessed with Shapiro–Wilk test for normality. Dependent groups were evaluated using Wilcoxon signed-rank test. A *p* value less than 0.05 was considered to be significant.

3. Results

Urinary excretion of the FC, TC and AC during abstinence after a period of heavy drinking is shown in Fig. 1.

Urinary excretion of the FC in healthy subjects was $88.6 \pm 54.1 \mu\text{mol/g}$ creatinine, TC excretion was $138.1 \pm 94.5 \mu\text{mol/g}$ creatinine, and the excretion of AC was $49.4 \pm 48.4 \mu\text{mol/g}$ creatinine. Fig. 1 shows significantly increased excretion of the FC, TC ($p < 0.05$) and AC ($p < 0.001$) at the first day (T0) of hospitalization as compared to the control group. Abstinence for 30 days (T30) significantly decreased excretion of the FC ($p < 0.001$), TC ($p < 0.001$) and AC ($p < 0.05$) as compared to the carnitine excretion at admission (T0). After 30 days of abstinence, excretion of FC and TC did not differ significantly from excretion in the control group, whereas excretion of AC remained significantly increased ($p < 0.001$). After 49 days of hospitalization (T49), urinary FC, TC and AC levels were not significantly different from those in the T30 and in control group.




Fig. 1. Urinary excretion of free (FC), total (TC) and acylcarnitine (AC) during abstinence after chronic drinking of the persons dependent on alcohol. Abbreviations: T0 admission day, T30 and T49 days of abstinence; significantly different: * ($p < 0.05$); *** ($p < 0.001$); ns, not significant.

Download English Version:

<https://daneshyari.com/en/article/2031986>

Download Persian Version:

<https://daneshyari.com/article/2031986>

[Daneshyari.com](https://daneshyari.com)