



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Biomedicine

journal homepage: [www.elsevier.com/locate/apjtb](http://www.elsevier.com/locate/apjtb)Original article <http://dx.doi.org/10.1016/j.apjtb.2015.12.018>

## *Agave sisalana* extract induces cell death in *Aedes aegypti* hemocytes increasing nitric oxide production

Louise Helena Guimarães de Oliveira<sup>1</sup>, Patricia Alexandria Paiva Silva de Sousa<sup>1</sup>, Fabrine Felipe Hilario<sup>1</sup>, Gabriel Joventino Nascimento<sup>1</sup>, João Paulo Saraiva Moraes<sup>2</sup>, Everaldo Paulo de Medeiros<sup>2</sup>, Manoel Francisco de Sousa<sup>2</sup>, Fabiola da Cruz Nunes<sup>1\*</sup>

<sup>1</sup>Biotechnology Center, Federal University of Paraíba, Caixa Postal 58.051-900, João Pessoa, PB, Brazil

<sup>2</sup>Embrapa Algodão, Brazilian Agricultural Research Corporation, Campina Grande, PB, Brazil



## ARTICLE INFO

## Article history:

Received 24 Sep 2015

Received in revised form 22 Oct, 2nd revised form 18 Nov 2015

Accepted 2 Dec 2015

Available online 21 Mar 2016

## Keywords:

Sisal

Cell culture

Mosquito

Insecticide

## ABSTRACT

**Objective:** To investigate the effects of *Agave sisalana* (*A. sisalana*) extract on *Aedes aegypti* (*Ae. aegypti*) primary cell culture.

**Methods:** Cells of *Ae. aegypti* were exposed to different concentrations of *A. sisalana* crude extract (0.18–6.00 mg/mL) for 24 h. Then, the cells were labeled with propidium iodide and subjected to fluorescence microscopy to verify cell viability. In addition, nitric oxide production was measured.

**Results:** Results showed that cells exposed to 6 mg/mL of the crude extract presented a greater percentage of death when compared to control (73.8% ± 9.6% vs. 34.6% ± 9.6%). Furthermore, there was an increase in the nitric oxide production in cells exposed to 6 mg/mL of *A. sisalana* crude extract [(0.81 ± 0.08) μmol/L] compared to control group [(0.41 ± 0.18) μmol/L].

**Conclusions:** The results show that *A. sisalana* is cytotoxic to *Ae. aegypti* and may be used as raw material for new eco-friendly and inexpensive insecticides, since sisal industry discards the liquid waste for the extraction of plant fiber.

## 1. Introduction

*Aedes aegypti* (*Ae. aegypti*) is a mosquito that is responsible for transmitting many diseases such as dengue fever, Chikungunya fever, yellow fever and Zika fever [1–3]. These diseases cause significant morbidity and mortality in developing countries [3]. Recently, dengue transmission has increased in urban and semi-urban areas of tropical countries around the world, becoming a major global public health problem. The World Health Organization estimates that there may be 50–100 million dengue infections worldwide every year [4,5].

Considering that there is no specific treatment or vaccine for dengue fever, its control is performed by combating with the mosquito vector [6–9].

Usually, mosquito control is done with chemical insecticides [9]. Mosquito larvae are usually targeted using organophosphates, insect growth regulators, and microbial control agents. Indoor residual spraying and insecticide-treated bed nets are also employed to reduce transmission of mosquito-borne disease in tropical countries [10,11]. However, studies have shown the development of resistance to the most common insecticides in many countries [9,12]. Therefore, the seek for new insecticides is needed, and natural products can be a reliable and eco-friendly source of raw materials for that purpose [13].

*Agave sisalana* (*A. sisalana*) is a monocotyledonous plant from the Agavaceae family, originally from Mexico. The plant, commonly known as sisal, is cultivated mainly in tropical and semi-arid areas around the world [14–16]. Several studies have been conducted to explore the possible biological activities of *A. sisalana*. Researchers have described anti-inflammatory, analgesic, anthelmintic and bactericidal activities of *A. sisalana* [17–19]. In addition, a recent study from our laboratory reported that *A. sisalana* possessed larvicidal activity against *Ae. aegypti* [20].

Therefore, the aim of this study was to investigate the effects of *A. sisalana* crude extract on *Ae. aegypti* hemocytes death,

\*Corresponding author: Fabiola da Cruz Nunes, PhD, Biotechnology Center, Federal University of Paraíba, Zip code 58.051-900, João Pessoa, PB, Brazil.

Tel/Fax: +55 083 3216 7511

E-mail: [fabiola@cbiotec.ufpb.br](mailto:fabiola@cbiotec.ufpb.br)

Foundation Project: Supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) [Grant No. 467147/2014-0 (VSL)].

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

exploring the possible mechanisms involved, especially as regards the nitric oxide production, which is an important substance involved in the insects immune system.

## 2. Materials and methods

### 2.1. Plant material

*A. sisalana* was cultivated in the city of Pocinhos (7°24'54" S, 39°24'36" W, 624 m above sea level) in Paraíba State, Brazil. The crude extract was obtained by grinding the leaves in a manual grinder until complete extraction of the crude extract. Then, the crude extract was strained and placed in a plastic container, protected from light, in a freezer at  $-10^{\circ}\text{C}$  until use. The crude extract was characterized regarding its physical and chemical composition, for determination of soluble solids ( $^{\circ}\text{Brix}$ ), total solids, saponins, reducing sugars and pH. Soluble solids ( $^{\circ}\text{Brix}$ ) were determined by manual refractometer. Total solids were determined according to the method described by Carle and Reinhardt [21], and saponins were determined by gas chromatography. Reducing sugars were determined by the dinitrosalicylic acid method [22], and pH was measured using a digital pH meter.

The Brazilian Agricultural Research Corporation (Embrapa), located in Campina Grande, Paraíba, was responsible for harvesting the plant and extracting the crude extract.

### 2.2. Primary cell culture

*Ae. aegypti* third instar larvae were sanitized with 70% alcohol and rinsed in phosphate buffer solution, and then they had their hemolymph collected and placed in tube containing phosphate buffer solution. The pool of hemolymph was centrifuged at 1500 r/min for 7 min in a refrigerated centrifuge at  $4^{\circ}\text{C}$  and the supernatant was discarded. The hemocytes were placed in cell culture vials containing Leibovitz medium (Himedia), supplemented with 10% of fetal bovine serum, penicillin-streptomycin (300 IU/mL), gentamicin (100  $\mu\text{g/mL}$ ), and fungizone (2.5 mg/mL). The culture was incubated at  $28^{\circ}\text{C}$  for 5 days.

### 2.3. Evaluation of cell viability by fluorescence microscopy

For the assay,  $1 \times 10^5$  cells from the primary cell culture were plated in 12-well plates and exposed to different concentrations of *A. sisalana* crude extract (0.18–6.00 mg/mL) for 24 h. Cells were centrifuged at 1500 r/min for 7 min in a refrigerated centrifuge at  $4^{\circ}\text{C}$  and the supernatant was discarded. The final volume was supplemented with 2 mL of Leibovitz medium (Himedia) and transferred to 15 mL tubes containing 15  $\mu\text{L}$  of propidium iodide to differentiate intact hemocyte from those necrosis. After 15 min of incubation in the dark, the cells were counted using a fluorescence microscope (Motic, Xiamen). In the control groups cells were exposed only to PBS. Assay was performed in triplicate.

### 2.4. Measurement of NO

For this assay,  $1 \times 10^5$  cells were plated in 12-well plates and exposed to *A. sisalana* crude extract (0.18–6.00 mg/mL)

for 24 h. The NO production was evaluated in the cultures exposed to different concentrations of the *A. sisalana* crude extract (0.18–6.00 mg/mL), according to method of Green *et al.* [23]. For this purpose, 50  $\mu\text{L}$  of the cell supernatant was collected and placed in 96-well plates containing 50  $\mu\text{L}$  of Griess reagent and incubated for 15 min. In the control groups cells were exposed only to PBS. The assay was performed in triplicate. The absorbance was measured using a microplate reader under 562 nm and the NO was quantified using a standard curve of  $\text{NaNO}_2$  as a reference.

### 2.5. Statistical analysis


Statistical analysis was performed using GraphPad Prism version 5.0 for Windows (GraphPad Software, San Diego, CA). Significant differences between groups were analyzed by ANOVA followed by Tukey's HSD test ( $P < 0.05$ ).

## 3. Results

The composition of the crude extract included: soluble solids ( $7.70 \pm 0.09$ )  $^{\circ}\text{Brix}$ , pH 4.82, phenolics ( $1.93 \pm 0.02$ ) mg/L, nitrogen 980 mg/L, protein 6.13 mg/L, glucose 12.4 g/L, fructose 1.2 g/L and saponins 2.4 mg/mL.

In order to evaluate the cytotoxicity of the *A. sisalana* crude extract, hemocytes were exposed to different concentrations of the plant extract (0.18–6.00 mg/mL) for 24 h. Fluorescence microscopy showed that the cells exposed to concentrations below 6 mg/mL had no significant difference in cell death compared to the control group. The cells exposed to 6 mg/mL of the crude extract of *A. sisalana* showed ( $73.8 \pm 9.6$ )% of necrotic cells vs. ( $34.6 \pm 9.6$ )% in the control group, after 24 h of exposure (Figure 1). It is worth mentioning that it presented more 39.2% necrotic cells than that in the control group.

The NO is involved in immunological reactions from the insect's metabolism, and for this reason it was measured in this study. The results showed that the cells exposed to concentrations below 6 mg/mL had no significant difference in the NO production compared to the control group. After 24 h of exposure to the 6 mg/mL of *A. sisalana* crude extract, the NO concentration in the cell supernatant was ( $0.81 \pm 0.08$ )  $\mu\text{mol/L}$ , which was twice as much as that in the control group [ $(0.41 \pm 0.18)$   $\mu\text{mol/L}$ ] (Figure 2).



**Figure 1.** Percentage of cell necrosis of *Ae. aegypti* hemocytes exposed to different concentrations of crude extract of *A. sisalana*. Bars with the same letter are not significantly different by Tukey test, 5%.

Download English Version:

<https://daneshyari.com/en/article/2032529>

Download Persian Version:

<https://daneshyari.com/article/2032529>

[Daneshyari.com](https://daneshyari.com)