



## Microbial turnover of above and belowground litter components in shrublands



Venkata S.S.R. Marella, Paul W. Hill, Davey L. Jones, Paula Roberts\*

School of Environment, Natural Resources and Geography, Bangor University, Gwynedd, LL57 2UW, UK

### ARTICLE INFO

#### Article history:

Received 28 April 2016

Received in revised form 4 July 2016

Accepted 10 July 2016

#### Keywords:

Belowground carbon storage

Mineralisation

Nutrient cycling

Litter decomposition

Root turnover

### ABSTRACT

Shrublands cover a large proportion of the world's land surface, yet they remain poorly studied in comparison to other ecosystems. Within shrublands, soil organic matter (SOM) is replenished from inputs of both above- and below-ground plant litter, however, their relative importance depends on their respective turnover rates. To critically address this, we measured the biodegradation rates of the soluble and insoluble components of  $^{14}\text{C}$ -labelled above- and below-ground plant litter in soil. During the 150 day incubation, the amount of plant-derived soluble-C lost as  $^{14}\text{CO}_2$  was similar for the different plant parts being  $64.7 \pm 2.3\%$  for roots,  $72.1 \pm 7.4\%$  for stems, and  $72.4 \pm 1.8\%$  for leaves. In comparison, the turnover of the insoluble fraction was much slower. However, again little difference in mineralisation was seen for the different plant parts with the total losses being  $21.1 \pm 0.9\%$  for roots,  $19.5 \pm 1.6\%$  for stems, and  $19.6 \pm 1\%$  for leaves. A double exponential first order kinetic model fitted well to the experimental data. It also allowed the partitioning of C between microbial anabolic and catabolic processes for the soluble C component. Using this model, we deduced that the soluble fraction turns over *ca.* 40 times annually, whereas it takes *ca.* 2.5 years to turnover the insoluble fraction. For the soluble plant component, the overall microbial carbon use efficiency (CUE) was estimated to be greater for root-derived C in comparison to that derived from aboveground (no difference was observed for the insoluble component). From this, we tentatively suggest that C sourced from belowground plant components may persist longer in soil than C derived from aboveground plant components.

© 2016 Elsevier GmbH. All rights reserved.

Soil organic matter (SOM) represents a major store of terrestrial carbon (C) (Schlesinger, 1997) and its turnover and replenishment represent a critical component of the global C cycle. SOM is primarily derived from the continual input of above- and below-ground plant components, however, their relative importance, particularly in shrubland ecosystems, remains poorly understood (Vogt et al., 1986). Earlier studies have suggested that plant roots contribute a larger proportion of C to soil organic carbon (SOC) than plant shoots, due to their greater chemical recalcitrance in relation to microbial enzymatic breakdown (Broadbent and Nakashima, 1974; Jane et al., 2007). In contrast, within some agroecosystems, significant contributions by crop shoots have also been observed (Barber, 1979).


The input of organic matter to the soil can be broadly classified into two pools (van Hees et al., 2005). The first pool is described as

the dissolved organic C component that includes low molecular weight, highly bioavailable compounds such as organic acids, peptides, amino acids, mono- and oligo-saccharides, amino sugars, phenolics and siderophores (Mc Keague et al., 1986). The second pool consists of plant polymers such as cellulose, hemicellulose, lignin and some proteins, which are relatively resistant to microbial attack (Kalbitz et al., 2000). These two pools can have vastly different C:N:P ratios which may subsequently influence their rate of processing and also microbial carbon use efficiency (CUE; Schmidt et al., 2011).

Numerous studies have described the mineralisation of individual low molecular weight compounds (Glanville et al., 2012), plant material (Simfukwe et al., 2011) and have measured the subsequent rates of  $^{14}\text{CO}_2$  evolution and/or microbial incorporation. These studies have enhanced our understanding of the  $^{14}\text{C}$  mineralisation process of single or occasionally combinations of simple C compounds by the microbial community. However, plant material consists of vast range of compounds (Buckingham, 1993) and the mineralisation capacity of

\* Corresponding author.

E-mail address: [p.roberts@bangor.ac.uk](mailto:p.roberts@bangor.ac.uk) (P. Roberts).



**Fig. 1.** Cumulative mineralisation of soluble and insoluble fractions of either  $^{14}\text{C}$ -labelled shrub roots, stems or leaves after addition to soil. Data points represent means  $\pm$  standard error of the mean ( $n=3$ ).

microorganisms to act upon more complex suite of substrates provides a more representative estimate of the potential for C storage in soil. Therefore, the aim of this study was to assess the microbial turnover of the soluble and insoluble fractions of above- and below-ground plant components (root, stem, leaf) from a common shrubland plant to assess their persistence in the soil under laboratory conditions.

Soil was obtained from the Henfaes experimental station located in Abergwyngregyn, Gwynedd, North Wales ( $53^{\circ}14'N$ ,  $4^{\circ}01'W$ ) UK. The sandy clay loam textured soil is classified as a Eutric Cambisol (FAO) or Dystric Eutrudepts (US Soil Taxonomy) (see SM<sub>1</sub> and Table S<sub>1</sub>). *Cistus monspeliensis* L. plants were grown in a hydroponic system consisting of 50% strength Long Ashton nutrient solution under laboratory conditions. Plants were labelled with  $^{14}\text{C}$  twice, 3 days apart for 5 h each time to get sufficient translocation of  $^{14}\text{C}$  to all plant components (see SM<sub>2</sub>). Immediately after the second labelling, the plant components were separated into leaves, stem, and roots and air-dried. The dried plant parts were finely ground using a ball mill and stored in 50 ml polypropylene tubes at 20 °C for further analysis. The distribution of  $^{14}\text{C}$  label among soluble and structural fractions of plant material was determined by performing a sequential chemical extraction. These results were tested in parallel with unlabelled plants, using an automated fibre analyser (see SM<sub>3</sub>). The soluble and insoluble fraction from each of the three plant components were separated using a hot water extract (see SM<sub>4</sub>) and amended to field-moist soil contained in 50 cm<sup>3</sup> polypropylene tubes. The mineralisation of the  $^{14}\text{C}$ -labelled components was studied for 150 days and values were expressed as a percentage of the initial amount of  $^{14}\text{C}$  applied to the soil (see SM<sub>5</sub>). Similar extraction process was conducted with unlabelled plant components and the soluble fraction from each component was analysed for distribution of low molecular weight ( $\leq 300$  Da) compounds using MALDI-TOF mass spectrometry (Bruker Reflex IV) with TiO<sub>2</sub> as a matrix. At the end of the incubation period, the amount of soluble  $^{14}\text{C}$  remaining in the soil either as unaltered plant material or fixed in the microbial biomass was determined by extracting the soil in 0.5 M K<sub>2</sub>SO<sub>4</sub> (see SM<sub>6</sub>). A

double exponential first order decay model was then fitted to the experimental data (Glanville et al., 2016). Substrate-C pool distribution within the microbial community, decay constants, CUE and half-lives (Newton-Raphson iteration method) (Oburger and Jones, 2009) were calculated (see SM<sub>7</sub>). The data was analysed by one-way ANOVA with Post-Hoc least significant difference test using SPSSv20.0 (SPSS Inc., Chicago, IL) using  $P < 0.05$  as an indication of statistical significance.

Following the labelling process, the distribution of  $^{14}\text{C}$  into soluble and structural fractions of the different plant components was broadly similar to the total amount of unlabelled  $^{12}\text{C}$  in each chemical fraction, although the data for stems is not available (Table S<sub>2</sub>). This indicates a fairly uniform dilution of the  $^{14}\text{C}$  isotope within the plant. The addition of  $^{14}\text{C}$ -labelled soluble and insoluble fractions to soil caused an initial rapid phase of  $^{14}\text{CO}_2$  evolution followed by a secondary slower phase, irrespective of plant tissue type (Fig. 1). The overall amount of  $^{14}\text{C}$  mineralisation in soils amended with soluble fractions was substantially higher compared to the values obtained for the insoluble fractions ( $P < 0.001$ ). This was presumably due to the presence of more labile low molecular weight compounds in the soluble fractions. Conversely, insoluble fractions broadly consist of structural polymers which require enzymatic depolymerisation to promote solubilisation prior to uptake and assimilation by the microbial community (van Hees et al., 2005). Among the soluble fractions, root-derived  $^{14}\text{C}$  showed the fastest mineralisation rate followed by stem and leaf  $^{14}\text{C}$  during the first hour, presumably because of relatively higher quantities of low molecular weight compounds which exist in roots (Figs. S<sub>1</sub> and S<sub>2</sub>). After 24 h, the amount of  $^{14}\text{C}$  mineralisation of the root soluble fraction ( $19.7 \pm 0.4\%$ ) was substantially higher than for the stems ( $8.7 \pm 0.3\%$ ) and leaves ( $5.7 \pm 0.3\%$ ). Similarly, among the insoluble fractions, the root-derived  $^{14}\text{C}$  fraction had the highest initial mineralization rate ( $0.62 \pm 0.2\%$ ) within 24 h, followed by the stems ( $0.43 \pm 0.02\%$ ) and leaves ( $0.26 \pm 0.01\%$ ). However, at the end of 150 days, the pattern had changed with  $64.7 \pm 2.3$ ,  $72.1 \pm 7.4\%$ , and  $72.4 \pm 1.8\%$  of the soluble fraction lost for the root, stem and leaf-derived  $^{14}\text{C}$ , respectively. In contrast, for

Download English Version:

<https://daneshyari.com/en/article/2060983>

Download Persian Version:

<https://daneshyari.com/article/2060983>

[Daneshyari.com](https://daneshyari.com)