


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/pjnns>

Case report

Hemorrhagic stroke, cerebral amyloid angiopathy, Down syndrome and the Boston criteria[☆]

Karol Jastrzębski^{a,1}, Magdalena Justyna Kacperska^{a,1,*}, Agata Majos^b,
Magdalena Grodzka^b, Andrzej Głabiński^a

^a Department of Neurology and Stroke, Medical University of Lodz, Lodz, Poland

^b Department of Radiological and Isotopic Diagnostics and Therapy, Medical University of Lodz, Łódź, Poland

ARTICLE INFO

Article history:

Received 6 October 2014

Accepted 21 April 2015

Available online 7 May 2015

Keywords:

Amyloid-beta

Hemorrhagic stroke

Down syndrome

Cerebral amyloid angiopathy

The Boston criteria

ABSTRACT

A stroke, or a cerebrovascular accident (CVA) is a life-threatening condition which often results in permanent or significant disability in the adult population. Several classifications of CVAs exist, one of them being based on the mechanism of injury of brain tissue: ischemic (85–90%) and hemorrhagic (10–15%). In a hemorrhagic stroke an intercranial bleeding occurs, leading to the formation of a focal hematoma typically located in the basal ganglia of the brain (approx. 45% of cases). A common yet underestimated cause of intracerebral hemorrhage is cerebral small vessel disease with microhemorrhages, including the cerebral amyloid angiopathy (CAA). This condition is associated with the deposition of amyloid-beta in arterial walls (in soft meninges, subcortical areas and the cerebral cortex). Research has shown that causes of hemorrhagic changes in the brain include genetic disorders, such as Down syndrome. The association is caused by the so-called 'gene dosage effect', as the gene for the precursor protein for amyloid-beta is located in chromosome 21. We wish to present the case of a 60 year old patient with Down syndrome who suffered a hemorrhagic stroke without antecedent hypertension. Based on the history taken, diagnostic imaging and the source literature, a diagnosis of cerebral amyloid angiopathy as the source of the bleeding was made (however it must be noted that without a full post-mortem examination, the Boston criteria allow only for a 'probable cerebral amyloid angiopathy' diagnosis to be made). The authors hereby also report the need to modify the Boston criteria for cerebral amyloid angiopathy.

© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

1. Introduction

According to the World Health Organisation report, cerebrovascular diseases including strokes are the second leading

cause of death in high and middle-income countries, surpassed only by the ischemic heart disease [1]. Projections of global mortality and disease burden until the year 2030 predict an increase in the number of deaths caused by strokes, which is attributed to an aging population [2]. Poland is one of

[☆] By the decision of the Rector of the Medical University of Lodz, the resolution no. 277/2014 dated 17 Apr 2014 renamed the Department of Neurology, Epileptology and Stroke into the Department of Neurology and Stroke.

* Corresponding author at: Department of Neurology and Stroke, Medical University of Lodz, Zeromskiego 113, 90-549 Lodz, Poland.

Tel.: +48 042 639 35 91; fax: +48 042 639 35 91.

E-mail address: karol.jastrzebski@umed.lodz.pl (K. Jastrzębski).

¹ These authors contributed equally to this work.

<http://dx.doi.org/10.1016/j.pjnns.2015.04.006>

0028-3843/© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

the few Eastern European countries with rising mortality rates due to strokes [3]. The poorest outcomes were associated with hemorrhagic strokes: the intracerebral hemorrhage and the subarachnoid hemorrhage [4,5]. Classic risk factors for intracerebral hemorrhage are hypertension, aneurysms, vascular malformations, bleeding diatheses, the use of psychoactive substances, brain tumors and cerebral amyloid angiopathy, among others [6]. According to the source literature, the cerebral amyloid angiopathy was found in approximately one third of patients with lobar intracerebral hemorrhages [7]. Pathological changes associated with CAA may be found in patients with Alzheimer's disease as well as Down syndrome, although their importance in the latter case is not fully understood. There are reported cases from various countries, including Poland, of histologically confirmed intracerebral hemorrhages in patients with Down syndrome with coexisting cerebral amyloid angiopathy [8-10]. In Down syndrome there is an overexpression of genes located in the triplicated chromosome 21. In case of APP (amyloid precursor protein APP) the expression is increased four to five times in individuals with DS (Ann N Y Acad Sci. 1993 Sep 24;695:91-102. Regulation and expression of the Alzheimer's beta/A4 amyloid protein precursor in health, disease, and Down's syndrome. Beyreuther K, Pollwein P, Multhaup G, Mönning U, König G, Dyrks T, Schubert W, Masters CL). This overexpression of APP is thought to be responsible for Alzheimer's-like dementia in patients with DS (Neurology 2004; 62:1996-1998, M. Margallo-Lana, C.M. Morris, A.M. Gibson, et al.). The AD-like dementia start at the median age of 50 years, which strongly suggests that beta-amyloid exists not only in the brain but also in small vessels. However, only 22% of patients suffering from AD have cerebral microbleeds. (Cerebral microbleeds and Alzheimer's disease, Charlotte Cordonnier and Wiesje M. van der Flier, in Cerebral Microbleeds edited by David J. Werring, Cambridge University Press, London, 2011). The post-mortem examination of 117 brains of patients with AD revealed signs of CAA in 97 brains (83%), out of which 30 (25,6%) showed moderate to severe CAA. Brains in the latter group revealed more ischemic and hemorrhagic changes compared to the group with little to no amyloid angiopathy (Neurology. 1996 Jun;46(6):1592-6.

Cerebral amyloid angiopathy in the brains of patients with Alzheimer's disease: the CERAD experience, Part XV. Ellis RJ, Olichney JM, Thal LJ, Mirra SS, Morris JC, Beekly D, Heyman A). The available body of research indicates that there is a connection between the APP gene expression, CAA and hemorrhagic strokes.

This study presents the case of a Down syndrome patient; after taking into consideration the patient's history, pathogenesis, and additional diagnostic studies, it was possible to establish an antemortem diagnosis of probable cerebral amyloid angiopathy based on the Boston criteria. However, the authors maintain that by applying the Occam's razor principle, the degree of probability is much higher than the criteria would lead one to believe.

2. Case report

A 60-year-old patient with Down syndrome (diagnosis based on patient's phenotype as well as available documentation) was admitted to the clinic with an acute onset of right-sided hemiparesis preceded by loss of consciousness. There were no signs of cranial trauma, nor any information indicative thereof from the paramedics. Neurological examination on admission revealed that the patient was conscious, somnolent, with incoherent speech, had eye deviation towards the left side, central facial palsy, paralysis of right upper limb and vestigial movement in right lower limb. Plantar reflex was present in the right lower limb. Deep tendon reflexes were diminished on the right side. The CBC, PT, PTT, LFTs, glucose and BUN values were within normal limits. A CT scan revealed a hemorrhagic focus in deep structures of the left hemisphere, which corresponded to the presented symptoms (Fig. 1) as well as bilateral lacunae of approx. 6 mm in diameter in basal ganglia.

In addition, calcifications in lenticular nuclei were noted. The patient was disqualified from surgical management. Due to the lack of history of hypertension, a contrast enhanced head MRI with a susceptibility weighted imaging (SWI) sequence was ordered to differentiate from other causes of bleeding, bearing in mind the association between Down




Fig. 1 – Hemorrhagic region in early subacute stage in deep structures of the left hemisphere with surrounding edema and modeling of the lateral ventricle. Also periventricular lacunar infarcts can be noticed on both sides. A – CT scan, B – T2-dependent axial image, C – T1-dependent axial image.

Download English Version:

<https://daneshyari.com/en/article/2152829>

Download Persian Version:

<https://daneshyari.com/article/2152829>

[Daneshyari.com](https://daneshyari.com)