



Hydroxycinnamic acid amides from *Scopolia tangutica* inhibit the activity of M1 muscarinic acetylcholine receptor in vitro



Yan Zhang^{a,*}, Zhen Long^{b,1}, Zhimou Guo^b, Zhiwei Wang^c, Xiuli Zhang^b, Richard D. Ye^a, Xinmiao Liang^b, Olivier Civelli^c

^a School of Pharmacy, Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China

^b Key Laboratory of Separation Science for Analytical Chemistry, Key Lab of Natural Medicine, Liaoning Province, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, People's Republic of China

^c Department of Pharmacology, University of California, Irvine, CA 92697, United States

ARTICLE INFO

Article history:

Received 15 September 2015
Received in revised form 12 November 2015
Accepted 13 November 2015
Available online 14 November 2015

Keywords:

Scopolia tangutica Maxim
Hydroxycinnamic acid amides
M1 muscarinic acetylcholine receptor

ABSTRACT

Scopolia tangutica Maxim (*S. tangutica*) extracts have been traditionally used as antispasmodic, sedative, and analgesic agents in Tibet and in the Qinghai province of China. Their active compositions are however poorly understood. We have recently isolated five new hydroxycinnamic acid (HCA) amides along with two known HCA amides, one cinnamic acid amide from these extracts. In this study, we evaluate their abilities to inhibit carbachol-induced activity of M1 muscarinic acetylcholine receptor along with the crude extracts. Chinese hamster ovary cells stably expressing the recombinant human M1 receptor (CHO-M1 cells) were employed to evaluate the anticholinergic potentials. Intracellular Ca^{2+} changes were monitored using the FLIPR system. Five HCA amides as well as the crude *S. tangutica* extract displayed dose-dependent inhibitory effects against M1 receptor. These findings demonstrate that HCA amides are part of the M1 receptor-inhibiting principles of *S. tangutica*. Since blockade of parasympathetic nerve impulse transmission through the inhibition of the M1 receptor lessens smooth muscle spasms, our findings provided a molecular explanation for the traditional use of *S. tangutica* against spasm.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Scopolia tangutica Maxim (*S. tangutica*) is one of the traditional Chinese medicines used in Tibet and in the Qinghai province of West China. Its extracts have been traditionally used as antispasmodic, sedative, and analgesic agents [1,2]. Its medicinal importance is associated with high biologically active tropane alkaloids, in particular anisodine and scopolamine [3–5]. These alkaloids primarily affect the parasympathetic nervous system and act as muscarinic acetylcholine receptor (mAChR) antagonists [6–8]. mAChR antagonists block the transmission of parasympathetic nerve impulses by acetylcholine and consequently lessen the spasms of smooth muscle [9,10]. Recently we have isolated five new hydroxycinnamic acid (HCA) amides along with two known HCA amides, one cinnamic acid amide from *S. tangutica* [11,12]. The conjugates of HCAs with polyamines have been found in a wide variety of plants and are concentrated in the reproductive organs of many higher plants [13–15]. The HCA-spermidine conjugates have been proven to be useful for chemotaxonomic purposes in several plant

families [16–18]. The HCA amides have been proposed to have a number of biological activities, including the effects of anti-fungi, anti-bacteria and anti-tumor [19–21], suggesting that HCA amides may have great potential for identifying new molecules for different diseases. It is interesting to note the similarity of the plant derived HCA amides to polyamine conjugates found in the invertebrate venom [22,23]. Some of these invertebrate toxins have been found as neurotransmitter receptor ligands [24,25]. In this study, we examined the anticholinergic potential of these HCA amides and compared the anticholinergic properties to the crude extract. The goal of the study is to foster a better understanding of the traditional use of *S. tangutica* and pursuit of the discovery of novel drugs.

mAChRs belong to G protein-coupled receptors (GPCR) family A. They are found in nervous system where they mediate the metabotropic actions of the neurotransmitter acetylcholine, in particular its parasympathetic action. They are major drug targets in human disease therapies. Five subtypes of mAChRs, classified as M1–M5, have been identified [26]. M1, M3 and M5 receptors couple to Gq proteins, which mediate intracellular calcium mobilization upon stimulation. M2 and M4 receptors couple to Gi proteins which inhibit adenylyl cyclase activity [27]. The sequence similarities found among these receptors infer that binding of a ligand to one of them should insure binding to all. In this study we chose to use the M1 receptor as the lead receptor.

* Corresponding author.

E-mail address: zhangyan_sjtu@sjtu.edu.cn (Y. Zhang).

¹ These authors contributed as co-first authors.

2. Materials and methods

2.1. Plant materials

The roots of *S. tangutica* were collected from Jiuzhi county (at 100°40'19" E; 33°16'56" N), Qinghai province, People's Republic of China. Voucher specimens (QTPMB) were deposited at Qinghai–Tibet Plateau Museum of Biology, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, and were identified by Senior Engineer Lijuan Mei.

2.2. Extract preparation and compound purification

The crude extract was prepared as described before [28]. In brief, 95 g *S. tangutica* was grounded into powder and extracted twice with 1 L ethanol–water (95:5, v/v) at 80 °C for 3 h. The crude extracts were obtained by rotary evaporation of the solvent. 1 mL of crude extract was dissolved in 5 mL of methanol. An aliquot of 0.1 mL of crude extract sample was applied to the conditioned SCX-based solid phase extraction column. The isolation and identification of pure compounds from *S. tangutica* are extensively described by our most recent report [12]. Briefly, *S. tangutica* were extracted by 95% ethanol. The isolation of pure compounds was achieved by multi-dimensional HPLC. The identity of the isolated compounds was confirmed by spectroscopic and spectrometric methods (1D- and 2D-NMR, LC-MS, optical rotation). The purity of all isolated compounds was $\geq 98\%$.

2.3. M1 receptor activity measurement

Chinese hamster ovary cells stably expressing the recombinant human M1 receptor (CHO-M1 cells) were kindly provided by Dr. Naoto Hoshi (University of California, Irvine). The M1 receptor inhibitory activity was determined by Fluorometric Imaging Plate Reader (FLIPR). The assay was performed as reported earlier with slight modification [29]. Briefly, the cells were seeded into black wall, clear-bottom 96-well plates at a density of 60,000 cells per well. Twenty-four hours later the medium was removed and replaced with 100 μ l of dye loading solution (2 μ M Fluo-4 AM dissolved in FLIPR buffer, which consists of pluronic acid in 1 \times Hank's buffer supplemented with 20 mM HEPES, pH 7.4) for 1 h at 37 °C. The cells were then washed 3 times with FLIPR buffer prior to FLIPR assay. The compounds were dissolved in dimethyl sulphoxide (DMSO) and diluted with FLIPR buffer. For antagonist tests, the compounds were first incubated with the cell for 10 min, before the addition of 50 nM carbacol to induce the activity of M1 receptor. Data were expressed as fluorescence (arbitrary units) versus time.

2.4. Data processing

IC₅₀ values and curve fitting were determined using Graphpad Prism (GraphPad Software, Inc., San Diego, CA). Data from each dose-response curve were normalized to the carbacol (50 nM) stimulation. At least three independent experiments were performed. Data are presented as means \pm standard error of the mean (SEM).




Fig. 1. Structures of compounds 1–11 isolated from *S. tangutica*. Seven HCA amides (1–7): scotanamine A–D (1–4), N¹, N¹⁰-di-dihydrocaffeoylspermidine (5), cafeoylputrescine (6), N¹-caffeoyl-N³-dihydrocaffeoylspermidine (7); one cinnamic acid amide: carboxycotangamine A (8) and three well-known anticholinergic compounds (9–11): scopolamine (9), atropine (10) and anisodine (11).

Download English Version:

<https://daneshyari.com/en/article/2538234>

Download Persian Version:

<https://daneshyari.com/article/2538234>

[Daneshyari.com](https://daneshyari.com)