



ELSEVIER

Contents lists available at ScienceDirect

## American Journal of Infection Control

journal homepage: [www.ajicjournal.org](http://www.ajicjournal.org)

## Commentary

Challenges in assessing transmission of *Mycobacterium tuberculosis* in long-term-care facilities

David A. Jackson MD<sup>a</sup>, Karen Mailer MD<sup>b</sup>, Kimberly A. Porter PhD<sup>c</sup>,  
R. Todd Niemeier MS, CIH<sup>d</sup>, Donna A. Fearey MSN, ANP<sup>e</sup>, Linda Pope MD<sup>b</sup>,  
Lauren A. Lambert MPH<sup>f</sup>, Kiren Mitruka MD, MPH<sup>f</sup>, Marie A. de Perio MD<sup>d,\*</sup>

<sup>a</sup> University of Cincinnati Medical Center and Cincinnati Children's Hospital Medical Center, Cincinnati, OH

<sup>b</sup> Providence Health & Services, Anchorage, AK

<sup>c</sup> Epidemic Intelligence Service, Alaska State Health Department, Centers for Disease Control and Prevention, Atlanta, GA

<sup>d</sup> National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Cincinnati, OH

<sup>e</sup> Alaska Department of Health and Social Services, Anchorage, AK

<sup>f</sup> Division of Tuberculosis Elimination, Centers for Disease Control and Prevention, Atlanta, GA

In 2012, the Centers for Disease Control and Prevention (CDC) reported 3.2 cases of tuberculosis (TB) per 100,000 persons in the United States.<sup>1</sup> Although TB incidence has declined during the past several decades, the 2010 goal of < 1 case per 1,000,000 persons—as established in the national strategic plan for TB elimination—has yet to be achieved.<sup>2</sup> Although persons aged ≥ 65 years accounted for only 14% of the population in 2012, this group represented 22% of reported cases of TB.<sup>3,4</sup> An analysis of 1993–2008 cases reported in the United States showed that the rate of TB among elderly adults was as much as 30% higher than among younger adults.<sup>5</sup> Even more striking are the disproportionate rates documented among those living in long-term-care facilities (LTCFs). Previous reports have estimated that adults aged ≥ 65 years residing in LTCFs may have between 4 and 50 times the risk of developing TB disease than elderly persons living in the community.<sup>5–7</sup>

As of April 2014, approximately 3.2 million workers were employed in LTCFs.<sup>8</sup> The size of this occupational group will grow significantly in the coming years if LTCF resident populations increase as expected. Past estimates suggest the TB case rates are 3 times higher among LTCF workers compared with those working in any other job.<sup>9</sup> Therefore, prevention and control of TB in LTCFs are essential to protect both the residents and employees in these settings. The goal of this article is to summarize findings of an LTCF TB outbreak investigation to highlight the unique

challenges posed by *Mycobacterium tuberculosis* transmission in these settings.

## CASE STUDY

During 2011–2012, the Alaska Department of Health and Social Services, the Anchorage Department of Health and Human Services, CDC's National Institute for Occupational Safety and Health (NIOSH), and CDC's Division of Tuberculosis Elimination investigated suspected *M tuberculosis* transmission at a 190-bed LTCF in Alaska. In April 2011, the investigation was initiated when the facility's annual employee TB screening program identified an aberration in the number of tuberculin skin test (TST) conversions. In contrast to no TST conversions in previous annual screenings, 8 of 230 evaluated employees had documented TST conversions (defined as ≥ 10 mm increase). In this LTCF, employees without a history of TB disease or latent TB infection (LTBI) were screened with a TST upon hire and then every April. Similarly, residents without a history of TB disease or LTBI were screened via TST within 72 hours of admission and then annually during the month of their admission anniversary. Residents and employees with a history of TB disease or LTBI undergo a baseline chest radiograph and annual TB symptom screening via questionnaire.

Subsequent screening of residents from April–November 2011 used an interferon-gamma release assay (IGRA) and identified newly positive IGRA test results, defined as ≥ 0.35 IU/mL, among 8 of 17 residents (47%) of a single secure unit for dementia special care.

Using a risk-stratified approach to preselect those at highest risk of infection, thorough screening of 216 of 350 (62%) current and former employees and 85 of 155 (55%) current and former residents was undertaken via varying combinations of symptom review; TST or IGRA; chest imaging; and sputum collection through

\* Address correspondence to Dr. Marie A. de Perio, MD, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, 1090 Tusculum Ave., R-10, Cincinnati, OH 45226.

E-mail address: [mdeperio@cdc.gov](mailto:mdeperio@cdc.gov) (M.A. de Perio).

The findings and conclusions presented here are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Conflicts of interest: None to report.

expectoration, induction, or bronchoscopy as needed. Deciding between TST or IGRA was based on which test had been used previously for each person evaluated.<sup>10</sup>

Ultimately, 12 residents and 11 employees were found to have new LTBI, and 1 resident was found to have pulmonary TB disease. This resident had a history of pulmonary TB disease in the remote past and resided in the secure dementia special care unit. The resident had a comprehensive negative prior workup for TB between April and November 2011, including chest radiographs, bronchoscopy with bronchoalveolar lavage for acid-fast bacilli (AFB) smear and culture, and stool polymerase chain reaction test for TB. From a pulmonary specimen collected in May 2012, the resident was found to have both a positive AFB sputum smear result and a positive AFB culture result for *M tuberculosis*, which were interpreted as probable reactivation of TB. All 11 employees and 8 of 12 residents with newly diagnosed LTBI were epidemiologically linked to this resident with TB disease, who reportedly did not move outside of this unit during the time at the LTCF. However, no clear epidemiologic link to the other 4 residents, who resided outside of the secure dementia special care unit, was found.

We cross-checked the limited paper visitor and volunteer logs available against the state TB disease database, without any matches. Evaluation of the ventilation system did not find an airflow pattern that could explain how the other 4 residents could have been infected from the resident with TB disease. One of these LTCF residents with significant weight loss, productive cough, and TST conversion died from respiratory failure during August 2012 while our investigation was still ongoing. Because no postmortem examination was performed, we were unable to determine whether this individual had TB disease and was the source of exposure to the other 3 residents. The 1 resident with documented TB disease received appropriate therapy promptly upon diagnosis, and all those with LTBI were offered treatment. Improvements to administrative controls such as training of employees and improved screening procedures were implemented at the LTCF. No further evidence of ongoing transmission has been noted as of July 2014.<sup>11</sup>

## SITUATIONAL ANALYSIS

Throughout the course of the investigation, many diagnostic, staffing, and treatment difficulties were encountered. LTCFs present a unique set of challenges for TB prevention and control, and diagnosing TB among elderly persons can be complicated. Therefore, understanding issues encountered in outbreak investigations aids in refining areas for continued clinical and public health research and informing updated guidelines and policies to protect residents and employees.

### Diagnostic challenges

#### Recognizing signs and symptoms

Administrative controls, starting with early detection of contagious TB, are the cornerstone of preventing transmission in health care settings.<sup>12</sup> However, TB has long been recognized as an imitator of multiple other diseases, making diagnosis challenging even under ideal conditions.<sup>13</sup> Further, aging reduces the ability to mount a cell-mediated immune response, making atypical presentations of TB due to dissemination of TB disease or localization to other organs more common.<sup>6,14</sup> In elderly persons, classic presenting features of pulmonary TB disease such as weight loss, cough, hemoptysis, and night sweats may be either absent or attributable to alternative diagnoses.<sup>6,14,15</sup> Conditions such as dementia and strokes can be associated with dysphagia, increasing the possibility of chronic cough from recurrent aspiration.<sup>16</sup> Elderly persons are also more prone to weight loss because of poor

nutrition and increased metabolic demand from chronic diseases.<sup>17</sup> Obtaining reliable information from symptom screening can be especially challenging given the prevalence of cognitive impairments.<sup>18</sup>

TB disease in elderly persons may have an insidious onset with persistently decreased ability to perform activities of daily living, new onset or worsening of preexisting cognitive impairment, and increasing fatigue and dyspnea.<sup>15</sup> Given the broad differential diagnosis necessary for chronic signs and symptoms such as these, TB diagnosis is delayed in some instances, increasing patient morbidity and mortality and allowing more time for transmission, especially in congregate settings.<sup>5,6</sup>

### Diagnostic testing

CDC recommendations for TB screening of residents and employees of LTCFs are based on initial and ongoing facility-specific risk assessments.<sup>12</sup> Medium-risk LTCFs, such as the 1 in this investigation before the outbreak, are defined as settings where patients with TB are expected to be encountered. In these settings, initial screening via a 2-step TST or a single IGRA should be used for baseline LTBI evaluation. Thereafter, CDC recommends annual screening with a single TST or IGRA unless there is documentation of a prior positive TST or IGRA result. In individuals with a baseline positive test result for LTBI, or completion of LTBI or TB disease therapy, a single chest radiograph should be performed to exclude TB disease with routine symptom screening thereafter.<sup>12</sup>

The limitations of TST have been well described. Exposure to nontuberculous mycobacteria can cause a false-positive TST result.<sup>6,12,19,20</sup> False-negative TST results are more common in aging populations due to impaired immunity leading to the possibility of unrecognized TB infection.<sup>5,21</sup> Additionally, TST findings are only valid if the tests are performed and interpreted properly. Therefore, TB should remain in the differential diagnosis in elderly patients residing in LTCFs with clinical signs and symptoms of TB, even for those with a negative TST result.<sup>6</sup>

IGRAs can be performed at a single patient encounter; however, IGRAs have their own limitations. IGRA is generally more expensive than TST, requires proper blood collection and transport, and must be processed promptly.<sup>10</sup> Recent reports have documented variable results with serial IGRA testing, making interpretation challenging.<sup>22</sup> False-positive IGRA conversions have been noted among health care workers in areas or settings with a low prevalence of TB.<sup>22</sup> Additionally, scant data exist on IGRA use in elderly populations.<sup>6,23,24</sup> Finally, use of TSTs and IGRAs in combination can lead to diagnostic dilemmas from discordant results.<sup>10</sup> We encountered both discordance between serial IGRA tests and discordance between IGRA and TST results during our investigation.<sup>11</sup>

A chest radiograph is recommended for persons with signs or symptoms of TB disease or with a positive test result for LTBI and should be performed before initiation of therapy for LTBI to rule out TB disease. Upper-lobe infiltrates, often with evidence of cavities, are the classic TB findings on chest radiograph.<sup>12</sup> The most easily interpretable chest radiograph includes 2 upright views: posterior-anterior and lateral. Proper technique dictates use of the correct exposure, appropriate body positioning, and patient-held maximally inspired breath.<sup>25</sup>

Obtaining adequate radiographs in the LTCF population is challenging. Curvature of the spine and cognitive impairment can make proper positioning and capturing of images at full inspiration difficult.<sup>7</sup> Additionally, most LTCFs are not equipped to perform 2-view, upright chest radiographs within their facilities, but rather are limited to lower quality, portable, single-view, anterior-posterior images. Ideally, LTCF residents with suspected TB disease should be transported to a facility equipped for full diagnostic imaging.

# دانلود مقاله



<http://daneshyari.com/article/2636729>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات