



Feature Article

Caring for people with dementia and challenging behaviors in nursing homes: A needs assessment geriatric nursing



Jeanette M. Daly, RN, PhD^{a,*}, Camden P. Bay, MS^b, Barcey T. Levy, PhD, MD^{a,b}, Ryan M. Carnahan, MS, PharmD^b

^aDepartment of Family Medicine, Carver College of Medicine, The University of Iowa, Iowa City, IA, USA

^bDepartment of Epidemiology, College of Public Health, The University of Iowa, Iowa City, IA, USA

ARTICLE INFO

Article history:

Received 5 November 2014

Received in revised form

29 December 2014

Accepted 5 January 2015

Available online 9 February 2015

Keywords:

Nursing home

Director of nursing

Dementia

Challenging behaviors

Antipsychotic medication

ABSTRACT

An estimated 50% of nursing home residents have a dementia diagnosis. The purpose of this research was to conduct a needs assessment of directors of nursing (DON) in Iowa nursing homes in relation to caring for patients with Behavioral and Psychological Symptoms of Dementia. DON responses were linked to Online Survey Certification and Reporting/Certification and Survey Provider Enhanced Reporting (OSCAR/CASPER) data to examine how facility characteristics may be associated with use of and confidence in non-drug management strategies. From 431 questionnaires mailed to DONs, 160 (37%) were returned. Regression analysis showed that those who were more confident in managing challenging behavior were more likely to have satisfaction with current training on managing challenging behaviors and had a psychiatrist available to visit the facility. Facilities with a larger proportion of patients with challenging behaviors being treated with non-drug approaches instead of antipsychotics had DONs who were more likely to be confident in non-drug management strategies and have knowledge about the FDA antipsychotic medications risks.

© 2015 Elsevier Inc. All rights reserved.

Currently, there are more than 15,000 Medicare and Medicaid-certified nursing homes in the United States¹ with almost 3.3 million patients receiving care in those nursing homes.² An estimated 50% of nursing home residents have a diagnosis of dementia^{3,4} with 13% having dementia or Alzheimer's disease as their primary diagnosis.⁵

Neuropsychiatric disturbances occur in 60–98% of people with dementia, frequently triggering nursing home placement and antipsychotic treatment.⁶ Antipsychotics have been a mainstay of pharmacologic management. In 2006, 25.9% of nursing home residents in the United States received antipsychotics,⁷ compared to 16.4% just 10 years earlier. Recent estimates place the prevalence of nursing home antipsychotic use around 22%.⁸ Evidence suggests that antipsychotic use is often suboptimal, based on lack of appropriate indications, inappropriate dosing,^{9,10} and high variability in usage rates across facilities.¹⁰ This is of particular concern since antipsychotics are associated with serious side effects in people with dementia, including an increased risk of mortality

which was the reason for a black box warning required by the Food and Drug Administration. In 2014 the Centers for Medicare and Medicaid Services (CMS) launched the National Partnership to Improve Dementia Care initiative to enhance dementia care quality and address antipsychotic overuse, with a goal of reducing antipsychotic use in nursing homes by 30% at the end of 2016.¹¹ Regulatory requirements have facilitated a marked improvement in decreasing the use of antipsychotic medications in nursing homes from 24% in the fourth quarter of 2011 to 20% in the first quarter of 2014.¹¹ The current focus of the initiative is to not only reduce antipsychotics but also to encourage person-centered care and non-drug management of behavioral symptoms of dementia in order to reduce unnecessary antipsychotic use.¹²

Challenging behaviors for persons with dementia are common and multi-faceted being attributable to the environment, chronic conditions, health, or medications. More than 80% of nursing home residents with dementia exhibit one or more forms of challenging behaviors.¹³ Examples of challenging behaviors include agitation, anger, depression, disrobing, eating abnormalities, hoarding, inappropriate sexual behavior, irritability, paranoia, physical and verbal aggression, repetition, swearing, and withdrawal.

There are situations in which antipsychotic use in dementia patients is considered appropriate by most clinicians as well as CMS, particularly when a patient's behaviors are dangerous to

Funding: ARHQ R18 HS19355-01.

* Corresponding author. 01290-F PFP, 200 Hawkins Drive, Iowa City, IA 52242, USA. Tel.: +1 319 384 8995.

E-mail address: jeanette-daly@uiowa.edu (J.M. Daly).

themselves or others.¹⁴ Initial therapy for patients with dementia who have challenging behaviors is environmental modifications and non-pharmacologic approaches. However, if that is unsuccessful and antipsychotic use is warranted, an appropriate indication for use of the medication needs to be documented. Once antipsychotic medication is started ongoing monitoring to evaluate its effectiveness and observation for adverse effects is necessary. If the medication is ongoing, the lowest effective dose should be established and maintained. In 2013, the percentage of nursing home residents taking antipsychotic medication ranged from 12.3% in Hawaii to 33.1% in Louisiana with a mean of 24.2% in the U.S.¹⁵

In Iowa the percentage of nursing home residents taking antipsychotic medication was 21.2% in 2012.¹⁵ The purpose of this research study was to conduct a needs assessment of directors of nursing (DON) in Iowa nursing homes in relation to caring for patients with dementia and those with challenging behaviors. That is a needs assessment to establish knowledge deficits in which to appropriately care for people with Behavioral and Psychological Symptoms of Dementia (BPSD) and then to best determine how they would like to receive additional training. This needs assessment would help to inform the development of educational materials and tools to help guide care for people with BPSD. In addition, the DON responses were linked to the Online Survey Certification and Reporting/Certification and Survey Provider Enhanced Reporting (“OSCAR/CASPER”) survey and inspection data to provider responses to examine how information on facility operations, patient census and regulatory compliance may be associated with use of and confidence in non-drug management strategies for behavioral symptoms of dementia. Comparison of both resident and facility data allows us to look at the variation and influence of these factors. In particular, we utilized regression modeling to aid in assessing the relationship between the proportion of dementia patients with challenging behavior treated using non-drug approaches and nurse’s knowledge, confidence, and education, family involvement, medical director education, availability of a psychiatrist or nurse practitioner with a dementia/psychiatric specialty, facility operations, and patient census (see [Table 1](#) for the entire list OSCAR/CASPER Data). Additionally, we wanted to look at nurse’s confidence in managing challenging patient behaviors associated with dementia and non-drug approaches.

Methods

After institutional review board approval, a mailed survey was sent to Iowa nursing homes. The Medicare nursing home compare website (medicare.gov/NursingHomeCompare) listed 443 nursing homes in Iowa. Of the 443, 12 were hospitals and eliminated from the list because the cover letter read director of nursing and would go to the hospital director and not the nursing home director, leaving 431 nursing homes. All nursing home DONs were invited to participate and questionnaires were mailed in January 2011. DONs were asked to complete the questionnaire while skipping any questions they did not want to answer and mail it back. If no response was given by a DON within three weeks, a second questionnaire was mailed. Following the two mailings, no further mailings or contacts were made.

Instrument

A 22-item questionnaire (see [Appendix 1](#)) entitled, “Needs Assessment: Caring for People with Dementia and Challenging Behaviors” developed by the research team of clinicians, researchers, and education specialists, was used to assess the needs of DONs in regard to caring for individuals with BPSD. To ensure

content validity, the questionnaire was critique by current or former nursing home employees that were on the research team. Questions included information pertaining to the treatment of dementia patients and the education of nursing home staff and family members about dementia. To better understand current dementia management approaches, DONs were asked about which antipsychotics were most often prescribed (a list of commonly used antipsychotics was provided), satisfaction with current training on managing dementia behaviors, if the nursing staff provides influence on the selection of the antipsychotic medication prescribed, their awareness of the FDA warning on antipsychotics in dementia, and their usage as well as confidence in non-drug treatment approaches for managing BPSD.

To gather information related to the education of nursing home staff and family members of dementia patients, DONs were asked how they obtain information about dementia treatment, what dementia topics interest them the most, preferred methods of delivery (such as, CD-ROM, e-mail, handbook, mail, online resources, or pocket guides, poster) for dementia treatment information and what method would be best for communicating about the treatment of dementia symptoms with family members. Information about the medical director and availability of a psychiatrist or dementia specialist nurse consultant to consult or make a visit to the nursing home was collected. DON demographic information was collected and included age, gender, race, ethnicity, total years in practice, and total years employed at the current nursing home.

OSCAR/CASPER data

OSCAR is an administrative database of the Centers for Medicare and Medicaid Services (CMS). In July 2012, OSCAR data was replaced by the CASPER system. CASPER is a part of a large relational database operating within CMS Automated Survey Processing Environment (ASPEN). Every nursing home in the United States that is certified to provide services under either Medicare or Medicaid is listed in the survey data. The number of variables in a data file depends on the survey process and extensive data is collected for nursing homes. Data are available for download, but new survey data overwrites the previous data. Archived longitudinal data for this study were purchased from Cowles Research Group so the data was current with the survey.¹⁶

OSCAR/CASPER data were linked to each nursing home’s DON responding to the questionnaire through the nursing home’s respective National Provider Identifier (NPI). Thirty-two CMS OSCAR/CASPER variable descriptions were retrieved from their respective forms which included: 1) Long Term Care Facility Application for Medicare and Medicaid, 2) Resident Census and Conditions of Residents, 3) Statement of Deficiencies and Plan of Correction (see [Table 1](#)). The research team reviewed the forms and determined which variables to include in the data analysis.

Data analysis

Using the returned questionnaires, basic descriptive statistics for questions of interest were calculated. For any statistics derived from the questionnaires and reported as proportions, missing responses have been removed from the denominator for calculation; any missing data exceeding 10% for a question has been noted in the results.

The analysis explored predictors for two of the questions asked on the questionnaire: 1) the proportion of dementia patients who had challenging behavior treated with non-drug approaches instead of medications (questionnaire takers were given five choices: “less than 1/4,” “less than 1/2 but more than 1/4,” “about 1/2,” “more than 1/2 but less than 3/4,” and “more than 3/4”), 2) the

دانلود مقاله



<http://daneshyari.com/article/2648351>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات