



King Saud University
The Saudi Dental Journal

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Experimentation and correlates of electronic nicotine delivery system (electronic cigarettes) among university students – A cross sectional study



K.H. Awan*

Department of Oral Medicine & Diagnostic Sciences, College of Dentistry, King Saud University, Riyadh, Saudi Arabia

Received 4 October 2015; revised 23 November 2015; accepted 19 December 2015

Available online 20 May 2016

KEYWORDS

E-cigarettes;
University students;
Smoker;
Questionnaire;
Prevention

Abstract Objective: E-cigarettes are becoming popular among youth as safe nicotine delivery systems. Many have expressed concern, however, that e-cigarettes may serve as a gateway to future smoking, given their low perceived risk, or that their use may prevent regular smokers from quitting by maintaining their nicotine addiction. The aim of this study was to assess experimentation with and correlates of e-cigarette use among university students.

Material and methods: A cross-sectional study was carried out among 480 university students from four faculties at a university in Riyadh in August–October 2014. A modified version of the World Health Organization's Global Adult Tobacco Survey was used, and multinomial logistic regression was carried out to assess correlations with e-cigarette variables in the whole study sample and among smokers.

Results: Almost all students, including the majority of ex-smokers (96.3%) and smokers (94.4%), reported having heard about e-cigarettes. In addition, about one-quarter of the sample (54.2% of smokers, 24.7% of ex-smokers, 6% of never smokers) had experimented with e-cigarettes at least once during their lifetime. Curiosity and peer influence were reported as the main reasons for the use of e-cigarettes. Factors found to be correlated significantly with e-cigarette use were male gender, being a traditional cigarette smoker, having friends who have tried e-cigarettes, and having a strong belief that e-cigarettes could aid smoking cessation.

Conclusion: E-cigarettes are popular among Saudi youth, especially among smokers and ex-smokers. Well-designed health education programs and regulatory interventions are required to address this issue.

© 2016 The Author. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Tel.: +966 (1) 467 7422; fax: +966 (1) 467 9018.

E-mail address: kamranhabibawan@gmail.com.

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

Electronic nicotine delivery systems, more commonly called electronic cigarettes or e-cigarettes, are battery-operated devices based on an inhalation-activated system that heats a

cartridge, producing vapor that the user inhales. The cartridges contain liquids that consist of various combinations of propylene glycol, glycerin, nicotine, tobacco extracts, flavorants, and/or adulterants. As e-cigarettes do not rely on combustion, users do not expose themselves or others to many of the harmful tobacco smoke constituents and particles produced by regular cigarettes (Benowitz, 2011).

In recent years, the marketing and use of e-cigarettes have increased markedly throughout the world due to manufacturers' claims that these nicotine delivery systems are safe. Safety data, however, are sparse and inconsistent (Bahl et al., 2012; Kuschner et al., 2011; Lee et al., 2011). In addition, many have expressed concerns that the popularity of e-cigarettes among youth because of their low perceived risk may represent a gateway to future smoking (Grana, 2013), or that their use may prevent regular smokers from quitting by maintaining their nicotine addiction (Cobb and Abrams, 2011). Well-designed clinical trials need to be conducted to examine the safety of e-cigarettes and their efficiency in smoking termination.

Due to the lack of safety data and global regulations, as well as the novelty of these products, the performance of clinical trials may be challenging. Until trials can be undertaken, the collection of data on usage patterns through surveys may provide information about the effects of these products on a range of outcomes. Although the e-cigarette literature is expanding rapidly, very few data on the awareness and use of e-cigarettes among youth in Saudi Arabia are available. Hence, the aim of this study was to assess awareness of, beliefs about, and experimentation with e-cigarettes among university students in Saudi Arabia.

2. Material and methods

2.1. Population and study design

This cross-sectional study was conducted in August–October 2014 among university students from four faculties (medical, dental, pharmacy, and engineering) of King Saud University, Riyadh, Saudi Arabia. Participants were given a self-administered questionnaire with detailed instructions for completing it individually. Questionnaires were distributed during breaks from lectures or work. The participants were informed that their participation in the study was voluntary, and they were required to complete the questionnaire on site and to return it immediately to the research team. Standard informed consent procedures, including the protection of participants' anonymity and confidentiality, were used. The institutional research and ethics committees approved the study [FR 0319].

2.2. Instrument and data collection

The questionnaire was developed from the e-cigarette part of the World Health Organization's Global Adult Tobacco Survey (<http://www.cdc.gov/tobacco/global/gats/questionnaire/index.htm> [updated 25 May 2009, accessed June 2009]) and was peer-reviewed, piloted, and determined to be comprehensive. Questionnaire items solicited data on sociodemographic characteristics (age, gender, year of study) and awareness of and beliefs about e-cigarettes. E-cigarette use was assessed by questions concerning use in one's lifetime, reasons for use, use in the last month, intention to use, and experimentation

with e-cigarettes by peers, parents, and/or siblings. The questionnaire also assessed respondents' smoking behavior (smoker [smoked traditional cigarettes in the last month], ex-smoker [smoked in the past but not in the last month], never-smoker [never smoked traditional cigarettes]).

2.3. Data analysis

Data were analyzed using SPSS Inc. (Chicago). Responses were coded as numeric to facilitate data entry. Multinomial logistic regression analysis was carried out to assess correlations with e-cigarette variables in the whole study sample and among smokers. The dependent variable was experimentation with e-cigarettes (never [reference] versus ever during one's lifetime). The independent variables were gender; smoking status (0 = never smoker, 1 = ex-smoker, 2 = smoker); attitudes toward e-cigarettes (e.g., can aid smoking cessation, are less dangerous, only smokers use e-cigarettes; totally disagree/partially disagree/do not know versus totally agree/partially agree); experimentation with e-cigarettes by friends, parents, or siblings (yes versus no); frequency of cigarette smoking (≤ 1 pack/day versus > 1 pack/day); and intention to quit cigarette smoking within the next year (yes versus no). The level of significance was set at $P < 0.05$.

3. Results

A total of 480 students completed the questionnaire (response rate, 89.6%). The majority (63.3%) of participants was male and 36.7% were female, with a mean age of 24.0 ± 1.3 years.

3.1. Awareness, attitudes, and practice

More than one-third (33.8%) of participants were smokers and 16.6% were ex-smokers. Never-smokers comprised about half (50.4%) of the study population. Regarding awareness of e-cigarettes, the majority (93.8%) of participants reported having heard about e-cigarettes (ex-smokers, 96.3%; smokers, 94.4%; never-smokers, 89.3%).

More than half of the students who were aware of e-cigarettes believed that e-cigarettes are less dangerous than traditional cigarettes, with the highest percentage (63.3%) recorded among smokers (Table 1). More than two-thirds of the students reported that e-cigarettes could aid smoking cessation, and almost half of the students reported that only smokers use e-cigarettes.

Regarding the use of e-cigarettes, one-quarter of the study population reported that they had tried e-cigarettes at least once during their lifetimes, with the highest percentage of users (54.2%) found among smokers. Interestingly, no ex-smoker or never smoker reported the use of e-cigarettes in the last month. Curiosity and peer influence were the main reported reasons for ever trying e-cigarettes; only one-third of smokers used these products for smoking cessation.

Almost two-thirds of the study population reported that they had friends who had experimented with e-cigarettes, whereas less than 10% reported that their parents or siblings had used e-cigarettes. The majority (87.5%) of participants reported that they had no intention to use e-cigarettes in the next year; 12.7% of smokers declared their intention to use e-cigarettes in the next year.

3.2. Correlates of experimenting with e-cigarettes

The results of multinomial logistic regression analyses are shown in Table 2. Male students, smokers, and those whose peers who had tried e-cigarettes were more likely to experiment with e-cigarettes. Students who strongly believed that e-cigarettes play an important role in smok-

Download English Version:

<https://daneshyari.com/en/article/2688661>

Download Persian Version:

<https://daneshyari.com/article/2688661>

[Daneshyari.com](https://daneshyari.com)