



## Brief communication

## Visual acuity in an opportunistic raptor, the chimango caracara (*Milvago chimango*)



Simon Potier<sup>a,\*</sup>, Francesco Bonadonna<sup>a</sup>, Almut Kelber<sup>b</sup>, Olivier Duriez<sup>a</sup>

<sup>a</sup> CEFE UMR 5175, CNRS, Université de Montpellier, Université Paul-Valéry Montpellier, EPHE, 1919 route de Mende, 34293 Montpellier cedex 5, France

<sup>b</sup> Department of Biology, Lund University, Sölvegatan 35, S-22362 Lund, Sweden

## HIGHLIGHTS

- We estimated the visual acuity of the chimango caracara using operant conditioning experiments
- Chimango caracara visual acuity ranges from 15.08 to 39.83 c/deg
- Chimango caracaras have the lowest visual acuity estimated in any raptor to date
- Their relatively lower visual acuity may reflect their foraging on the ground

## ARTICLE INFO

## Article history:

Received 14 September 2015

Received in revised form 18 January 2016

Accepted 23 January 2016

Available online 25 January 2016

## Keywords:

Chimango caracara

Opportunistic raptor

Visual acuity

Operant conditioning

## ABSTRACT

Raptors are always considered to have an extraordinary resolving power of their eyes (high visual acuity). Nevertheless, raptors differ in their diet and foraging tactics, which could lead to large differences in visual acuity. The visual acuity of an opportunist bird of prey, the Chimango caracara (*Milvago chimango*) was estimated by operant conditioning. Three birds were trained to discriminate two stimuli, a positive grey uniform pattern and a negative grating pattern stimulus. The visual acuity range from 15.08 to 39.83 cycles/degrees. When compared to other birds, they have a higher visual acuity than non-raptorial birds, but they have the lowest visual acuity found in bird of prey so far. We discuss this result in the context of the ecology of the bird, with special focus on its foraging tactic.

© 2016 Elsevier Inc. All rights reserved.

### 1. Introduction

Among vertebrates, birds are certainly the most dependent on vision [1] although some groups, like procellariiforms and vultures may use olfaction more than previously recognized [2, 3]. Visual acuity is one of the most frequent parameters that has been studied in bird vision and many studies assessed visual acuity using operant conditioning experiments in various species ranging from passerines [4, 5] to raptors [6–8]. The eyes of diurnal raptors are often considered to afford elevated spatial resolving power, yet only few species have been extensively studied: the wedge-tailed eagle (*Aquila audax*), the brown falcon (*Falco berigora*), the American kestrel (*Falco sparverius*) and some vulture species [6, 7, 9–11]. Although visual acuity in the wedge-tailed eagle is the highest estimated in vertebrate to date, we cannot generalize these findings to all raptors. Raptors differ highly in their ecology: some are exclusive predators, others are scavengers or opportunists [12]. Because diet and foraging represent important factors that influence spatial resolving

power in a given species (e.g. active predators have higher visual acuity than herbivores among mammals) [13], raptors with different diets may thus have different resolving power.

Here, we estimated the visual acuity of an opportunistic species, the chimango caracara *Milvago chimango* (order falconiformes). In contrast to eagles, falcons and vultures that search for food in flight, chimango caracaras are opportunists in their feeding behavior and mainly forage on the ground [12]. This would suggest that they might have high visual acuity to search for small moving prey, but probably lower than predatory birds such as eagles or falcons that search for prey from a perch or in flight.

### 2. Methods

#### 2.1. Animals

The three captive born chimango caracaras used in this study were living in individual aviaries (dimensions 4 × 4 × 3 m high) at the falconry park “le Rocher des Aigles”, France (44°48'4"N, 1°36'55"E). Birds were weighed every day and were maintained at approximately 90%

\* Corresponding author.

E-mail address: [simon.POTIER@cefe.cnrs.fr](mailto:simon.POTIER@cefe.cnrs.fr) (S. Potier).

of their free feeding weights during the experiment (see Table 1). Because these birds are usually used in a flight show (except during the experiment), all had previous experience of operant conditioning.

## 2.2. Experimental aviary and experimental device

The trial aviary (8 × 20 m in area and 6 m high) was located outdoors and covered with a net. A starting perch was placed at 10 m distance from two wooden boxes (6 m from each other), in which two monitors (Samsung S22C300H) were placed to present the visual stimuli (created on R.3.1.2 (R Development Core Team 2014) and presented on PowerPoint 2010). Monitor sizes were 510 × 398 mm that correspond to 2.92 × 2.28° of visual angle for the birds at 10 m. The two boxes were 70 cm wide, 80 cm high and 120 cm long and painted in black to create a “dark room”, in order to avoid any effect of direct sunlight on the screens. The luminance in the two boxes while screens were turned off was measured for each test phase and was not different between the two boxes (Left box: 115.63 ± 8.57 Lux; right box: 109.08 ± 10.59 Lux, *t*-test, *t* = 0.48, *df* = 44.07, *p* = 0.63). A feeding box with ten compartments and a perch was placed below each box with the screens. Each compartment contained a piece of chicken meat as a reward. To avoid any effect of possible olfactory cues, each compartment was perforated. When a bird made a correct choice, the observer (S.P.) opened a compartment from a distance using electric motor with a switch.

## 2.3. Assessment of visual acuity

Before each experiment, we ensured that the cage was oriented in such way that birds never had to fly against the sun. The birds were trained to fly towards one of the screens and to choose between a negative stimulus (composed of black and white stripes) and a positive stimulus (uniform grey). The grey stimulus was either slightly darker (10%) or slightly brighter (10%) than the pattern to make sure the birds could not use brightness as a cue. The birds indicated a choice by flying to one of the two perches below the monitors. The determination of visual acuity involved two phases.

Phase 1 (training): the birds had to learn to fly from the starting perch to the uniform stimulus while the other monitor presented a coarse grating (1.71 cycles per degree, abbreviated c/deg). During the first two days, the feeding boxes were presented to the birds to ensure that they could see into the compartments. This step was necessary to train the birds to fly to the perch. Later, the food was hidden. During the training phase, the monitors were switched off after 5 s if the bird made an incorrect choice, and after the bird had finished eating if the birds made a correct choice. Two sessions of 30 trials were conducted every day. For each trial, if the bird made a correct choice, one compartment was opened and the bird received a reward (3 g of chicken meat). For a session of 30 trials, gratings and uniform greys were presented 15 times, on each side, while the side of the stimulus was changed in quasi-random order. To prevent side preferences, the positive stimulus was presented on the same side for a maximum of three consecutive trials [6].

Phase 2 (Test): when the bird made more than 80% of correct choice during 3 consecutive training sessions, the test phase could begin. Two sessions were conducted per day. Before each test session, we presented 5 coarse gratings to ensure that the birds were still conditioned to the grey pattern. We conducted 5 sessions of 30 trials and 3 sessions of 28

trials, with 9 different gratings (1.71, 9.59, 17.98, 20.55, 23.97, 28.77, 41.10, 47.95 and 143.84 cycles per degree) that were presented randomly across trials and sessions. Each grating was presented 26 times. When the birds were about to leave the starting perch, the monitors were switched off to ensure that they could not alter their decision on the way. The observer (S.P.) was hiding in a cabin to avoid any bias to the choice of the bird but he was not blind of the experiment in order to open a box compartment with a reward if the birds made a correct choice.

## 2.4. Eye size

Corneal diameters were measured with ImageJ 1.49 from close-up photographs of the three individuals, according to [14]. The mean corneal diameter (CD, in mm) for each species was translated to axial length (AL), using the formula [15] for diurnal animals:

$$AL = CD/10^{-0.22} \quad (1)$$

We then calculated the visual acuity (VA) using the allometric function [16]

$$VA = 10^{(1.42 \times \log_{10}(AL) - 0.11)} \quad (2)$$

We compared the visual acuity obtained by the allometric function with the visual acuity obtained by operant conditioning experiment.

## 2.5. Data analysis

Analyses of behavioral data were performed with R.3.1.2 (R Development Core Team 2014) using psyphy [17] and ggplot2 [18] packages. To determine the threshold of visual acuity (72.5% correct choices, binomial test, *n* = 26, *p* < 0.05), we fitted a psychometric function for each individual using a generalized linear model (GLM).

## 3. Results

The visual acuity of three chimango caracaras assessed by operant conditioning ranged from 15.08 to 39.83 c/deg (Fig. 1, Table 1). Individuals differed in their ability to learn the task. Especially, the individual (Flow) showing the lowest value of visual acuity needed twice as much time to be conditioned to the grey pattern than as the two others birds (Table 1).

Mean corneal diameter was 8.37 ± 0.17 mm. Visual acuity obtained by the allometric function was 32.55 ± 0.92 c/deg (Table 1).

## 4. Discussion

Using operant conditioning tests, we estimated the visual acuity in chimango caracaras ranging from 15.08 to 39.83 c/deg. The three individuals tested differed in their visual acuity, particularly one individual showed an extremely low visual acuity compared to the two other birds. This individual took more time to be conditioned and made more errors than the others in the test phase (see Fig. 1 and Table 1). We may have underestimated the visual acuity of this individual because of motivation problems, or this bird may also have a vision defect. Nevertheless, this bird did not behave differently in other situations

**Table 1**  
Information on birds used in the experiment.

Bird	Age (year)	Corneal diameter (mm)	Free feeding weight (g)	Weight in experiment (g)	Weight lost (%)	Number of training session before conditioned	Visual acuity operant conditioning (c/deg)	Visual acuity allometric function (c/deg)
Richard	4	8.2	320	275	14.06	10	39.83	31.62
Kougounia	2	8.2	290	280	3.45	12	30.65	31.62
Flow	2	8.7	285	270	5.26	22	15.07	34.40

Download English Version:

<https://daneshyari.com/en/article/2844035>

Download Persian Version:

<https://daneshyari.com/article/2844035>

[Daneshyari.com](https://daneshyari.com)