



Acute ischemic stroke associated with nephrotic syndrome: Incidence and significance – Retrospective cohort study



Hiroataka Iwaki^{a,c}, Masaru Kuriyama^{a,*}, Shuichiro Neshige^a, Shinichi Takeshima^a, Takahiro Himeno^{a,1}, Kazuhiro Takamatsu^a, Yutaka Shimoe^a, Hiromitsu Kobayashi^b, Masahiro Nomoto^c, Akio Tanaka^b

^a Brain Attack Center Ota Memorial Hospital, Department of Neurology, 3-6-28 Okinogami, Fukuyama, Hiroshima 720-0825, Japan

^b Department Radiology, 3-6-28 Okinogami, Fukuyama, Hiroshima 720-0825, Japan

^c Department of Neurology and Clinical Pharmacology, Ehime University Graduate School of Medicine, Japan

ARTICLE INFO

Article history:

Received 18 August 2015

Received in revised form 9 October 2015

Accepted 9 October 2015

Available online 22 October 2015

Keywords:

Arterial ischemic stroke

Nephrotic syndrome

Atherosclerosis

Diabetic nephropathy

Hypercoagulability

ABSTRACT

We report 10 cases with arterial ischemic stroke (AIS) with nephrotic syndrome (NS), and clarified its incidence and clinical characteristics. The patients having albumin less than 3.0 g/dl and serum cholesterol greater than 250 mg/dl at the same time were retrospectively screened from 11,161 cases of stroke. Furthermore, the patients of AIS showing heavy proteinuria were selected. The 10 cases were diagnosed as AIS with NS. Its incidence was 0.09% of all kinds of stroke and 0.12% of AIS. Their subtypes were 6 large-artery atherosclerosis, 3 small-vessel occlusion, and 1 cardioembolism. We carried out a retrospective cohort study to assess the association between NS and atherosclerosis progression in AIS patients. Seven AIS patients with NS due to diabetic nephropathy (cases; NS group) were compared with patients with AIS and diabetes mellitus (DM) without NS (control group). Control group subjects were matched in a 2:1 ratio to cases by age, sex, use of medications for DM, and hemoglobin A1c (HbA1c) level. The NS group had high cerebral artery atherosclerosis scores, especially in the anterior circulation. The NS group demonstrated atherosclerosis of the internal carotid and lower extremity arteries, although there were no statistical differences between the two groups. Study subjects had high serum fibrinogen and D-dimer levels, suggesting that AIS patients with NS have a greater degree of hypercoagulability than AIS patients without NS.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Nephrotic syndrome (NS) is defined by the presence of heavy proteinuria (protein excretion greater than 3.5 g/24 h), hypoalbuminemia (less than 3.0 g/dl), hyperlipidemia and peripheral edema [1]. Thrombotic diseases are also frequently observed. Arterial and venous thromboses are potential complications of NS. Arterial thromboses are less frequent than venous thromboses and the most common locations are femoral arteries, although other arteries may be involved [1–3]. Stroke associated with NS has rarely been reported in several case reports [4–10]. We report 10 cases of acute ischemic stroke (AIS) associated with NS and the results of a retrospective cohort study comparing AIS patients with and without NS. The study's purpose is to define the incidence of AIS with NS and assess whether NS is associated with atherosclerosis progression.

2. Subjects and methods

2.1. Subjects

This retrospective study analyzed hospitalized patients enrolled in our stroke registry. First, the patients having albumin less than 3.0 g/dl and serum cholesterol higher than 250 mg/dl at the same time were screened from 11,161 cases of stroke for 9 years from April 2004 to August 2013. From these patients, furthermore, the cases associated with AIS showing heavy proteinuria were selected. The proteinuria was estimated quantitatively by measuring the protein in a urine/24 h or qualitatively by using paper kit (Eiken Chemical Co. Japan). Stroke subtypes were determined according to the classification of the Trial of Org 10172 in Acute Stroke Treatment (TOAST) [11]. In the retrospective cohort study, AIS patients with NS due to diabetic nephropathy (DN) were compared with AIS patients without NS who were matched to cases in a 2:1 ratio by age, sex, use of medication for diabetes mellitus (DM), and hemoglobin A1c (HbA1c) level. Cases were compared with controls in terms of body mass index (BMI), morbidity of hypertension, and laboratory findings, including serum albumin, total cholesterol, low density lipoprotein (LDL) and high density lipoprotein (HDL) cholesterol,

* Corresponding author.

E-mail address: kuriyama@shouwa.or.jp (M. Kuriyama).

¹ Present address: Department of Neurology, Oita Red Cross Hospital, Japan.

triglycerides, creatinine, estimated glomerular filtration rate (eGFR) [12], hematocrit, platelet counts, prothrombin time (PT), activated partial thromboplastin time (aPTT), fibrinogen, and D-dimer.

2.2. Evaluation of cerebral arterial stenosis

In this retrospective study, the exact percentage of arterial stenosis could not be determined, because the enhanced MRA could not be available. Extent of atherosclerosis was evaluated by the simplified atherosclerosis score of intracranial and extracranial arteries on MRA, according to the scoring method by Uekita et al. [13]. All MRA and MRI findings were reviewed by two investigators (A.T. and H.K.) who were blind to the clinical data. There was 93% of interobserver agreement. In cases where there was disagreement, the investigators reviewed the findings, discussed them and reached a final diagnosis. Total 17 vessels including the common carotid arteries, the extracranial and intracranial portions of the internal carotid arteries and the vertebral arteries, the anterior, middle, and posterior cerebral arteries, and the basilar artery, were evaluated by the cervical and intracranial MRAs. The stenotic lesions were graded virtually from 0 to 2: 0 indicating less than 50% reduction compared to a normally appearing proximal arterial diameter; 1 indicating 50–99% reduction of the arterial diameter and/or unclearly depicted distal artery beyond the stenotic lesion, and 2 indicating an occluded artery. The individual total scores were divided into two scores: an anterior score and a posterior score related to the anterior circulation vessels (common carotid, internal carotid, anterior cerebral, and middle cerebral arteries) or posterior circulation vessels (vertebral, basilar, and posterior cerebral arteries), respectively.

2.3. Ultrasonography and ankle-brachial index

Ultrasonography of carotid artery and ultrasonic duplex Doppler scanning of the lower extremities arteries were performed by using the LOGIQE9, GE-Yokogawa Medical. The stenotic lesions of internal carotid artery were graded by the method of NASCET (North American Symptomatic Carotid Endarterectomy Trial) from 0 to 5: 0 indicating negative, 1 indicating 1–25%, 2 indicating 26–50%, 3 indicating 51–75%, 4 indicating 76–99%, 5 indicating an occluded artery. The stenotic lesions of lower extremities were graded from 0 to 3: 0 indicating negative, 1 indicating mild stenosis, 2 indicating moderate stenosis, 3 indicating severe stenosis. The atherosclerosis of lower extremity arteries were also evaluated by measurement of ankle-brachial index (ABI) by using Colin form BP-203 RPE III, OMRON, Kyoto. A total carotid ultrasonic score (T-CUS) and total extremities ultrasonic score (T-EUS) were estimated as the sum of the right and left ultrasound score grades for the carotid and extremity arteries, respectively. A total ABI (T-ABI) score was estimated as the sum of the right and left ABI.

3. Results

3.1. NS subjects selection

The patients having serum albumin less than 3.0 g/dl were 196 (1.9%) cases of 10,057 patients examined, and the patients having serum cholesterol higher than 250 mg/dl were 1201 (11.3%) cases of 10,619 patients examined. The patients fulfilled both two levels at the same time were only 21 cases. They included 2 cases with negative or equivocal (\pm) by the protein paper kit, 1 case with 1+ (30 mg/dl), 2 cases with 2+ (100 mg/dl), 5 cases with 3+ (300 mg/dl), and 11 cases with 4+ (1000 mg/dl). The 16 patients showed urinary protein 3+ or 4+, namely heavy proteinuria, and they consisted 10 AIS, 4 intracranial hemorrhage, and 2 cerebral venous thrombosis (CVT) (Table 1). Nine patients of these 16 patients showed protein excretion greater than 3.5 g/24 h. In the other 7 patients, the content of protein on urine/24 h was not measured. However, they all had moderate to severe peripheral edema and hyperlipidemia, and the diagnosis of NS was strongly suggested also in these 7 cases.

3.2. Incidence and subtypes of AIS with NS

The 10 cases, 8 males and 2 females, 64.4 ± 8.6 years (Mean \pm SD), were diagnosed as AIS associated with NS. Its incidence rate was 0.09% of all kinds of stroke (total 11,161 patients) and 0.12% of AIS (total 8116 patients) in our hospital. And, its incidence was 5 folds of CVT during the same duration. Their subtypes were 6 large-artery atherosclerosis (LAA), 3 small-vessel occlusion (lacune, SVO), and 1 cardioembolism (CE) by the TOAST criteria [11]. On MRI and MRA, middle cerebral arteries, anterior choroidal artery and internal carotid artery (occlusion) were involved in 6 cases, 1 case and 1 case, respectively. SVOs were located in the supra-tentorial basal ganglia in all 3 cases (Table 2).

3.3. Pathogenesis of NS

NS has not been diagnosed before the admission in any cases of 10 patients. Pathological diagnosis was decided by the histopathological examination of the renal biopsied samples in two cases; amyloid nephropathy due to primary amyloidosis in 1 case and membranoproliferative glomerulonephritis (MPGN) in 1 case. Eight cases were diagnosed as diabetic nephropathy (DN) based on the long history of DM and the exclusion of other pathogenetic diseases, although renal biopsy was not performed (Table 2).

3.4. Evaluation of atherosclerosis

Extent of atherosclerosis was evaluated by using the atherosclerosis score of intracranial and extracranial arteries on MRA. Two cases showed no or one vessel of stenosis, but 8 cases revealed more than two vessels of stenosis. There was no past history of coronary artery disease in all cases, except one case. The ultrasonography of carotid artery

Table 1
The patients of acute stroke.
Abbreviation: TC; total cholesterol, Alb; albumin, UP; urinary protein, TIA; transient ischemic attack, SVO; small-vessel occlusion, LAA; large-artery atherosclerosis, CE; cardioembolism, ICH; intracranial hemorrhage, SAH; subarachnoid hemorrhage, CVT; cerebral venous thrombosis.

	TIA	SVO	LAA	CE	Other	ICH	SAH	Total
Patients n.	660	2196	2547	2023	235	2200	733	10,619
①TC > 250	60 (9.1)	299 (13.6)	373 (14.6)	133 (6.6)	17 (7.2)	242 (11.0)	75 (10.2)	1201 (11.3)
Patients n.	632	2070	2436	1958	202	2055	675	10,057
②Alb < 3.0	3 (0.5)	16 (0.8)	44 (1.8)	66 (3.4)	9 (4.5)	48 (2.3)	8 (1.2)	196 (1.9)
Patients n.	263	886	1123	917	99	1226	477	5002
③Massive UP	8 (3.0)	21(2.4)	40 (3.6)	45 (4.9)	6 (6.1)	118 (9.6)	33 (7.0)	272 (5.4)
① + ② + ③	0	3	6	1	2 (CVT)	4	0	

Download English Version:

<https://daneshyari.com/en/article/3049543>

Download Persian Version:

<https://daneshyari.com/article/3049543>

[Daneshyari.com](https://daneshyari.com)