



Original article

Evaluation of the anti-inflammatory activity of the aqueous and ethanolic extracts of the leaves of *Albizia lebbek* in ratsGirish Gulab Meshram^{a,*}, Anil Kumar^b, Waseem Rizvi^b, C.D. Tripathi^a, R.A. Khan^b^a Department of Pharmacology, Vardhman Mahavir Medical College and Safdarjung Hospital, Inner Ring Road, New Delhi, 110029 Delhi, India^b Department of Pharmacology, Jawaharlal Nehru Medical College, Aligarh Muslim University, Aligarh, 202002 Uttar Pradesh, India

ARTICLE INFO

Article history:

Received 18 September 2014

Received in revised form

5 October 2014

Accepted 23 October 2014

Available online 30 January 2015

Keywords:

Albizia lebbek leaves

Carrageenan

Cotton pellet

Percentage inhibition

Anti-inflammatory

ABSTRACT

Albizia lebbek Benth. (Mimosaceae) is a medicinal tree used to treat several inflammatory ailments in the Indian traditional Ayurvedic system of medicine. The aim of the present study was to evaluate the possible anti-inflammatory activity of the aqueous (AE) and ethanolic (EE) extracts of the leaves of *A. lebbek* to support the ethnopharmacological claims. The study was carried out using Wistar rats (100–150 g). The AE and EE were prepared using the Soxhlet extraction process. The anti-inflammatory activity of the AE and EE of the leaves of *A. lebbek* were studied using carrageenan-induced paw edema and cotton pellet-induced granuloma models. The AE and EE of the leaves of *A. lebbek* at doses of 50, 100, and 200 mg/kg p.o. (oral administration) showed a dose-dependent and significant ($p < 0.05$) inhibition of carrageenan-induced hind paw edema with maximum percentage inhibition (PI) values of 22.34, 30.85, 39.36 and 22.53, 32.98, 42.55, respectively. The AE and EE at doses of 50, 100, 200 mg/kg p.o. significantly ($p < 0.05$) inhibited granuloma formation with PI values of 19.07, 27.57, 38.55 and 23.93, 32.23, 42.33, respectively. The AE and EE of the leaves of *A. lebbek* showed significant ($p < 0.05$) anti-inflammatory activity.

Copyright © 2014, Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Understanding inflammation has always been an enigma for mankind. Something as minor as a bruise or something as major as a myocardial infarction can trigger this phenomenon. The major classes of drugs to suppress inflammation are nonsteroidal anti-inflammatory agents (NSAIDs) and corticosteroids but their toxic adverse effects such as epigastric distress, peptic ulceration, osteoporosis, and iatrogenic Cushing's syndrome have limited their use.^{1,2} Looking at the present scenario, medicinal compounds derived from plant sources such as flavonoids, saponins, alkaloids,

terpenoids, glycosides, and coumarins could provide an excellent fountainhead to develop new anti-inflammatory agents, which could be more efficacious, safer, affordable, and accessible for patients.

Albizia lebbek Benth. (Mimosaceae), commonly known as 'Sirisa' in Sanskrit, is a tall, unarmed, deciduous tree distributed throughout India. The traditional systems of medicine have been utilizing various parts of the tree to treat several inflammatory ailments such as asthma, bronchitis, arthritis, allergies, snake bites, fractures, hemicranias, gingivitis, toothaches, and sinusitis.^{3,4} It is also claimed that the tree is useful in night blindness, cataract, leukoderma, erysipelas, leprosy, tuberculosis, scabies, amoebiasis, syphilis, spermatorrhoea, and piles.^{5,6}

The bark of *A. lebbek* has stolen the limelight by its pleotropic activities such as anti-inflammatory, immunomodulatory, analgesic, antiarthritic, antioxidant, antimalarial, antitumor, and anti-fertility activities, pushing the leaves to the background.^{7–17} Despite the leaves being the most abundant and accessible medicinal part of the tree, possessing anticonvulsive, nootropic, antimicrobial, and antiulcer activities, there are no studies

* Corresponding author. Department of Pharmacology, 6th floor, Vardhman Mahavir Medical College and Safdarjung Hospital, Inner Ring Road, New Delhi, 110029, Delhi, India.

E-mail address: drgirish23@yahoo.co.in (G.G. Meshram).

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.

evaluating their anti-inflammatory activity.^{18–21} Furthermore, phytochemical analysis of the leaves has revealed the presence of potentially bioactive alkaloids, triterpenoid saponins, and tri-O-glycoside flavonoids.^{22–24} This study was hence undertaken to evaluate the possible anti-inflammatory activity of aqueous (AE) and ethanolic (EE) extracts of the leaves of *A. lebbek* in rats at different doses.

2. Materials and methods

2.1. Plant material

The leaves were obtained from the botanical garden of the Aligarh Muslim University (AMU), Aligarh in May 2011 and were authenticated by Dr Wazahat Hussain, a taxonomist, and a voucher specimen (voucher no. 2149) was deposited in the Department of Botany, AMU for further reference. The shade-dried leaves were ground homogeneously using a mixer-grinder and approximately 100 g of the powder subjected to Soxhlet extraction, for 16 hours using 5 L of distilled water and 5 L of 99.9% ethanol (Scientific OEM, Mumbai, India) as solvents. The dark green, semi-solid extracts obtained were made free from the solvents by placing them in an incubator at 60°C for 12 hours. The yield of the AE was 21% and that of the EA was 12.398%. Previous toxicity studies of the leaves of *A. lebbek* did not show any toxicity and behavioral changes in rats up to 2000 mg/kg p.o. (oral administration) dose,²¹ hence doses of 50, 100, and 200 mg/kg p.o. were selected for the present study.

2.2. Animals

Wistar rats (100–150 g) were obtained from the Animal House, Jawaharlal Nehru Medical College (JNMC), Aligarh. They were housed at a temperature of 24 ± 2°C, 12-hour light/dark cycles, 35–60% humidity, in polypropylene cages, and fed a standard rodent diet with water *ad libitum*. Animals were deprived of food but not water 4 hours before the experiment.

2.3. Drugs

Indomethacin (Merck, Bangalore, India), Diclofenac (Reckitt Benckiser, Gurgaon, India), and Carrageenan (Sigma Chemicals, St. Louis, MO, USA) were procured from the respective companies and were used in the study.

2.4. Ethical considerations

Experimental procedures and protocols used in this study were approved by the Institutional Animal Ethics committee of the JNMC and conform to the “Guidelines for care and use of animals in scientific research” (Indian National Science Academy 1998, Revised 2000).

2.5. Carrageenan-induced rat paw edema model

The rats were divided into eight groups ($n = 6$), each receiving distilled water (control), diclofenac 20 mg/kg p.o. (reference standard), and 50, 100, 200 mg/kg p.o. dose of the AE and EE of *A. lebbek*, respectively. Carrageenan (0.1 mL of 1%) was injected into the subplantar tissue of the right hind-paw of each rat. The volume of the carrageenan injected into the foot was measured at 0, 30, 60, 120, and 180 minutes using a plethysmometer (Biodevices, New Delhi, India). The percentage inhibition (PI) at each time interval was calculated²⁵:

$$PI = \frac{(V_t - V_0)_{\text{control}} - (V_t - V_0)_{\text{treated}}}{(V_t - V_0)_{\text{control}}} \times 100$$

V_0 = Mean paw volume at 0 hours

V_t = Mean paw volume at a particular time interval

2.6. Cotton pellet-induced granuloma model

The rats were divided into eight groups ($n = 6$), each receiving distilled water (control), indomethacin 10 mg/kg p.o. (reference standard), and 50, 100, 200 mg/kg p.o. dose of the AE and EE of *A. lebbek*, respectively. Thirty minutes after drug administration, an autoclaved cotton pellet of 10 ± 1.0 mg was aseptically implanted subcutaneously in the back region of the rats while anesthetized with ether (Scientific OEM, Mumbai, India). Extracts were administered once daily for the next 7 days. On Day 8, animals were anesthetized again and cotton pellets (Datt Mediproducts Ltd., New Delhi, India) were removed surgically, freed from the extraneous tissue, and dried in a hot-air oven overnight at 60°C. The dried pellets were weighed and the increment in the dry weight of the pellets was taken as a measure of granuloma formation. The PI of granuloma tissue development was calculated²⁶:

$$PI = \frac{\text{Weight of pellet (control)} - \text{weight of pellet (test)}}{\text{Weight of pellet (control)}} \times 100$$

2.7. Estimation of median effective dose

The median effective dose (ED₅₀) values were estimated using GraphPad Prism software version 5.03 (GraphPad Software Inc., San Diego, CA, USA). The PI values obtained from the carrageenan-induced paw edema and cotton pellet-induced granuloma models were initially normalized to percentage activity assuming that the maximal response (100%) is seen at the dose of 200 mg/kg and the minimal response (0%) is seen at the dose of 0 mg/kg of the AE and EE. The log–dose response curves were then generated using a normalized nonlinear regression curve model, and by interpolation of the log dose (best-fit value) 50% activity was obtained. The antilog of the obtained log dose produced the ED₅₀ value.

2.8. Statistical analysis

Results were expressed as mean ± standard error of the mean (SEM). Statistical analysis was performed using one-way analysis of variance (ANOVA) followed by multiple Tukey’s comparison tests. A p value < 0.05 was considered statistically significant.

3. Results

3.1. Carrageenan-induced paw edema model

The AE and EE of the leaves of *A. lebbek* (50, 100, 200 mg/kg, p.o.) showed a dose-dependent, significant inhibition of carrageenan-induced rat paw edema from 0.5 hours to 3 hours following drug administration, compared to the control group. The maximum PI of paw edema by the AE was observed as 22.34 ($p < 0.05$), 30.85 ($p < 0.05$), and 39.36 ($p < 0.05$) at the doses of 50, 100, 200 mg/kg p.o., respectively. The maximum PI of paw edema by the EE was observed as 25.53 ($p < 0.05$), 32.98 ($p < 0.05$), and 42.55 ($p < 0.05$) at the doses of 50, 100, 200 mg/kg p.o., respectively. Diclofenac 20 mg/kg p.o. showed a maximum PI of 61.70% at 3 hours after its administration (Table 1). The ED₅₀ values of the AE and the EE were 28.91 mg/kg and 27.23 mg/kg, respectively.

Download English Version:

<https://daneshyari.com/en/article/3099696>

Download Persian Version:

<https://daneshyari.com/article/3099696>

[Daneshyari.com](https://daneshyari.com)