




Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.aidm-online.com



CASE REPORT

Cytomegalovirus enteritis in immunocompetent patients: Report of two cases diagnosed using single-balloon enteroscopy



Cheng-Lin Hsieh ^a, Chia-Hung Tu ^a, Min-Horn Tsai ^b,
Cher-Wei Liang ^c, Hsiu-Po Wang ^a, Ping-Huei Tseng ^{a,*}

^a Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan

^b Department of Internal Medicine, National Taiwan University Hospital Hsin-Chu Branch, HsinChu, Taiwan

^c Department of Pathology, National Taiwan University Hospital, Taipei, Taiwan

Received 17 August 2015; received in revised form 27 January 2016; accepted 4 May 2016
Available online 3 June 2016

KEYWORDS

Cytomegalovirus;
Enteritis;
Enteroscopy;
Ileitis;
Immunocompetent;
Jejunitis

Summary Cytomegalovirus (CMV) infection of the gastrointestinal tract involves mostly the colon and rectum and mainly develops in immunocompromised patients. CMV infection in the small intestines has rarely been reported in immunocompetent patients. We report two cases of CMV enteritis that developed in immunocompetent patients and involved the ileum and jejunum, respectively. Both of them were diagnosed with single-balloon enteroscopy (SBE) and further confirmed with histopathology. The first case is a 71-year-old woman with a presentation of obscure gastrointestinal bleeding and severe anemia. Neither esophagogastroduodenoscopy nor colonoscopy identified any active bleeding. SBE and biopsy disclosed multiple scattered ulcers in the distal ileum and histopathology confirmed CMV ileitis. The hemorrhage subsided after conservative medical treatment. The second case is a 59-year-old woman with a presentation of progressive abdominal pain. SBE showed diffuse irregularly-shaped ulcers located from the upper to middle jejunum, and CMV jejunitis was confirmed with endoscopic biopsy and histopathological examination. Antiviral therapy was prescribed and her abdominal pain improved gradually. We discuss the clinical manifestations and management strategies of CMV infection that develops in the small intestines of immunocompetent patients. In addition, we highlight the endoscopic characteristics of CMV enteritis

* Corresponding author. Department of Internal Medicine, National Taiwan University Hospital, Number 7, Chung-Shan South Road, Taipei 100, Taiwan.

E-mail address: pinghuei@ntu.edu.tw (P.-H. Tseng).

and the clinical utilities of SBE in the evaluation of patients with suspected CMV infection of the small intestines.

Copyright © 2016, The Gastroenterological Society of Taiwan, The Digestive Endoscopy Society of Taiwan and Taiwan Association for the Study of the Liver. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Cytomegalovirus (CMV) infection of the gastrointestinal (GI) tract mostly involves the colon and rectum and mainly develops in immunocompromised hosts, such as patients with human immunodeficiency virus (HIV) infection and AIDS, patients who received solid organ or hematopoietic stem cell transplantation, and patients with inflammatory bowel disease [1,2]. The clinical manifestations of CMV infection of the GI tract are nonspecific [3], and the endoscopic findings are also variable, ranging from mucosal erythema, discrete ulceration, pseudo-tumor formation, to bowel perforation [1,4]. Endoscopic biopsy with histopathology evaluation and immunochemical staining remains the gold standard for definitive diagnosis [1].

Recently, CMV infection of the GI tract has been increasingly reported in immunocompetent patients, and the most commonly involved locations are also the colon and rectum [5]. The mortality rate of CMV colitis in immunocompetent patients has been reported to be up to 31.8% [6]. However, CMV infection involving the small intestines in immunocompetent hosts has rarely been reported [7]. Here, we report two cases of CMV enteritis which developed in immunocompetent patients. Both cases were diagnosed with single-balloon enteroscopy (SBE) and confirmed with histopathology.

Case Reports

Case 1

A 71-year-old woman with hypertension, diabetes mellitus (DM), and end stage renal disease (ESRD) under regular hemodialysis was referred to our hospital for obscure gastrointestinal bleeding with intermittent tarry and bloody stool. The physical examination was only remarkable for generalized pallor. Laboratory studies showed a low hemoglobin level of 5.0 g/dL with a mean corpuscular volume of 91.6 fL. Esophagogastroduodenoscopy did not identify any bleeding source and repeated colonoscopy showed only fresh blood in the colon. Intestine-targeted transabdominal ultrasonography disclosed neither mass lesion in the abdomen nor bowel wall abnormality. To investigate a suspected bleeding source in the distal small intestine, we performed SBE (SIF-Q260) with a ST-SB1 splinting tube (Olympus Co., Tokyo, Japan) via the retrograde approach, which showed multiple (~40–50) 0.3–1.0-cm ulcers scattered over the distal ileum (Figure 1). Endoscopic biopsy was done and histopathology showed ulcers with ulcerative debris and granulation tissue. The adjacent villi were mildly

blunted but no cryptic distortion or granuloma was found (Figure 2). CMV ileitis was confirmed with immunohistochemical staining. HIV serology was negative. Supportive care with component therapy was instituted and her bleeding resolved gradually without further antiviral treatment.

Case 2

A 59-year-old woman with hypertension and DM presented with progressive abdominal pain over the upper abdomen. Esophagogastroduodenoscopy, colonoscopy, and transabdominal ultrasonography revealed no significant findings. Contrast-enhanced abdominal computed tomography revealed segmental small bowel wall thickening (~1.0 cm in thickness) with fair mucosal enhancement and engorged mesenteric vessels (Figure 3). The small bowel follow-through showed continuous mucosal irregularity and thickening extending from the proximal jejunum to ileum. SBE (SIF-Q260) with ST-SB1 splinting tube (Olympus Co.) via the antegrade approach was performed and showed diffuse irregularly-shaped ulcers with exudative coating and geographic distribution extending from the upper to middle jejunum (Figure 4). Biopsy was performed and histopathology showed mucosa ulceration with several nuclear inclusion bodies and positive CMV immunoreactivity (Figure 5). CMV jejunitis was confirmed. The serum CMV immunoglobulin-M antibody was negative, immunoglobulin-G antibody was positive, and the serum viral load test detected CMV viremia (2280 copies/mL). Both autoimmune profile and HIV screening test were negative. Antiviral therapy with ganciclovir (250 mg every 12 hours) for 10 days was prescribed, followed by oral valganciclovir (900 mg/d) for 1 month. Her abdominal pain subsided gradually and subsequent serum CMV viral load was undetectable.

Discussion

The diagnosis of CMV infection of the GI tract, especially the small intestines, in immunocompetent hosts is challenging. Among the rarely reported cases in the literature, most present with nonspecific symptoms, including poor appetite, abdominal pain, vomiting, and ileus, and were usually diagnosed with laparotomy for profuse bleeding or bowel perforation [4,7]. Most cases with CMV ileitis over terminal ileum were diagnosed using colonoscopy [2] and only one case of CMV jejunitis was diagnosed with SBE [8]. A high index of clinical suspicion for CMV enteritis and early institution of SBE is necessary to avoid delaying the correct diagnosis.

Download English Version:

<https://daneshyari.com/en/article/3278507>

Download Persian Version:

<https://daneshyari.com/article/3278507>

[Daneshyari.com](https://daneshyari.com)