



## CASE REPORT

# A cerebrovascular stroke following endoscopy for an elderly patient with acute upper gastrointestinal bleeding



Ahmed Gado <sup>a,\*</sup>, Basel Ebeid <sup>b</sup>

<sup>a</sup> Department of Medicine, Bolak Eldakror Hospital, Giza, Egypt

<sup>b</sup> Department of Tropical Medicine, Banysweef University, Banysweef, Egypt

Received 5 December 2014; accepted 4 January 2015

Available online 2 February 2015

### KEYWORDS

Gastrointestinal bleeding;  
Endoscopy;  
Complications;  
Stroke

**Abstract** Elderly people constitute an increasing proportion of those presenting with acute upper gastrointestinal bleeding. Not only in upper gastrointestinal bleeding is advanced age a risk of death, but also exceeding 60 years of age results in an increased risk of cerebrovascular and cardiovascular events. Factors likely to influence the morbidity and mortality associated with endoscopy in an elderly cohort with acute gastrointestinal bleeding include the acuity and severity of bleeding and the presence of comorbid conditions. Audits have shown a surprisingly high incidence of both morbidity and mortality following upper gastrointestinal endoscopy. The following incident is a case report of a cerebrovascular accident following diagnostic upper gastrointestinal endoscopy for an elderly patient with acute upper gastrointestinal bleeding.

© 2015 The Authors. Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Case report

A 67-year-old man presented to the hospital on July 2009 complaining of hematemesis of large amount of coffee ground blood. His medical history consisted of hypertension and diabetes mellitus. He was a smoker and using baby aspirin. His pulse rate was 114 beats per minute and blood pressure 140/90 mmHg. The physical examination was unremarkable.

\* Corresponding author at: Medical Department, Bolak Eldakror Hospital, Bolak Eldakror, Giza, Egypt. Tel.: +20 2 35837644 (residence), +20 1006809363 (mobile); fax: +20 2 35702164/35721225.

E-mail addresses: [agado1954@yahoo.com](mailto:agado1954@yahoo.com), [agado1954@gmail.com](mailto:agado1954@gmail.com) (A. Gado), [bebeid@hotmail.com](mailto:bebeid@hotmail.com) (B. Ebeid).

Peer review under responsibility of Alexandria University Faculty of Medicine.

Relevant laboratory values included hemoglobin of 9 g/dl. Liver and kidney function tests were normal. ECG was normal. Four hours later he developed a second attack of hematemesis. His blood pressure was 130/90 mmHg and pulse rate 130 beats per minute. He received normal saline and one unit blood. Early upper endoscopy was performed five hours after admission. The patient was American Society of Anesthesiologists (ASA) score 2 and initial Rockall score 2. Endoscopy was performed by a staff with attendance of a senior staff. A first degree relative signed the consent. Pre-procedure basic vital signs monitoring using pulse oximetry revealed pulse rate 98 beats per minute and blood pressure 93/47 mmHg. The patient had been sedated with IV midazolam (1 mg) but he was agitated during the procedure which necessitated adding 1 mg midazolam. Supplemental oxygen was administered during endoscopy. The patient was awake during and after the



**Figure 1** CT scan of the brain without contrast.

procedure. Pulse oximetry showed persistent oxygen saturation  $>90\%$  and transient drop of pulse rate to 54 beats per minute during the procedure. Endoscopy revealed normal esophagus and duodenum but the stomach was full of blood and could not be examined. The source of bleeding was not identified so the endoscopist decided to repeat the endoscopy after 24–48 h. Postprocedure vital signs monitoring revealed pulse rate 110 beats per minute and oxygen saturation 100%. During follow up, blood pressure was 130/90 mmHg and pulse rate 120 beats per minute. Twenty-four hours after endoscopy, the patient suffered an ischemic-cerebrovascular stroke leaving him with right sided hemiplegia, aphasia and bedridden. Urgent CT scan of the brain without contrast revealed left fronto-parietal infarction (Fig. 1). ECG was normal. CPK-MB and troponin levels were normal. Hemoglobin level was 5 g/dl. The patient received five units of blood and his pulse rate dropped to 95 beats per minute and hemoglobin recovered to 10 g/dl. For this reason, it was decided to refrain from second endoscopy. The patient received pantoprazole 40 mg per day for the possibility of ulcer, was advised to stop smoking and started physiotherapy. Few days later he received baby aspirin 75 mg per day. The patient was retained in the hospital for two weeks and discharged without diagnosing the cause of bleeding. He was followed up in the outpatient clinic.

A staff meeting was called to analyze why and how this complication had occurred and to consider strategies to minimize the risk of further complications. An elderly hypertensive diabetic patient presented with hematemesis underwent emergency diagnostic endoscopy five hours after admission and developed a cerebrovascular stroke 24 h after endoscopy. The possible cause for the cerebrovascular accident was severe prolonged postbleeding hypovolemia in an elderly patient. The patient had persistent tachycardia ( $>100$  beats per minute) from admission till the occurrence of the stroke. Hemoglobin level was checked only twice on admission and after the stroke (9 g/dl and 5 g/dl, respectively). Hemoglobin level was not checked before the procedure. Hypotension (93/47 mmHg), documented immediately before the procedure, was ignored. Transient bradycardia (54 beats per minute) occurred during the procedure which may be accompanied with hypotension. Blood pressure was not assessed during the procedure. In spite of evidence of active bleeding (blood in stomach and tachycardia), close observation

(pulse rate, blood pressure and hemoglobin level) was not performed for 24 h until the patient had the stroke. After the stroke, assessment of hemoglobin level revealed a low hemoglobin level (5 g/dl). Consequently the patient received massive blood transfusion (five units) which corrected the hemodynamic instability and anemia. Corrective actions were undertaken. In elderly or at risk patients with acute upper gastrointestinal bleeding, intense monitoring and early rapid resuscitation should be performed. Hemoglobin level should be checked every four to six hours. Blood pressure and pulse rate should be monitored before, during and after any endoscopic procedure. An endoscopic procedure may need to be deferred if the patient's blood pressure is too low. The hypotensive patient may require resuscitation with blood and/or IV fluids before it is safe to proceed. Good communication is necessary to improve staff performance.

Four years later the patient presented with an attack of hematemesis and melena. His pulse rate was 130 beats per minute, blood pressure 80/50 mmHg and hemoglobin 7 g/dl. He received four units of blood transfusion. Bleeding stopped spontaneously. Endoscopy was postponed until stabilization of hemodynamic state and hemoglobin level recovered to 11 g/dl. Endoscopy was performed 48 h after admission. The patient had been sedated with IV midazolam (2.5 mg). Supplemental oxygen was administered during endoscopy. Blood pressure, pulse rate and oxygen saturation were monitored before, during and after the procedure without any significant change. The patient was ASA score 3, initial Rockall score 4 and final Rockall score 5. Endoscopy revealed a pre-pyloric gastric ulcer with flat pigmented spot. Biopsy for *Helicobacter pylori* was positive. The patient received a course of 500 mg Clarithromycin, 20 mg Omeprazole and 500 mg Tinidazole; all given twice daily for two weeks. He was discharged few days later without complications. Follow up endoscopy performed after two months revealed normal endoscopic findings and biopsy for *Helicobacter pylori* was negative. The patient was followed in the outpatient clinic until the time of writing this report.

## 2. Discussion

Elderly people (aged over 60 years) constitute an increasing proportion of those presenting with acute upper gastrointestinal bleeding.<sup>1</sup> Advanced age has been consistently identified as

Download English Version:

<https://daneshyari.com/en/article/3431593>

Download Persian Version:

<https://daneshyari.com/article/3431593>

[Daneshyari.com](https://daneshyari.com)