

# The principles of safe laparoscopic surgery

Bruce Levy

Mohammad Mobasheri

## Abstract

The well-established benefits of laparoscopic surgery over open surgery and the rapid technology developments have meant that in just over 30 years laparoscopy has become the first-line modality for performing a host of operative procedures. The objective of this paper is to give a broad overview of laparoscopic surgery, to include: (1) patient positioning and how this aids surgery; (2) methods of establishing pneumoperitoneum, including the open Hassan and closed Veress needle techniques; (3) gas insufflation; (4) maintaining good vision during surgery and how this can be achieved; (5) insertion of additional ports; (6) a description of available laparoscopic instruments and their uses; (7) limitations of laparoscopy; and (8) closure of port sites. Despite its advantages, serious iatrogenic complications have been reported with laparoscopic surgery and the vast majority of these are avoidable. For this reason the paper places a particular emphasis on safety during laparoscopic surgery and steps that should be taken to ensure this.

**Keywords** Keyhole; laparoscopic surgery; laparoscopy; safety

## Introduction

The significant benefits offered by the minimally invasive approach of laparoscopic surgery have consequently seen a rapid development in the principles, practice and technology used in this field. In just over 30 years, the laparoscopic approach has become the first-line approach to many general surgical operations. The first laparoscopic appendicectomy was performed on May 30, 1980 by a German gynaecologist called Kurt Semm. During the early 1980s, news of Semm's laparoscopic appendicectomy spread throughout the world. Erich Mühe was particularly interested in the possibility of performing additional operations using the laparoscopic technique and by 1984, Mühe had worked out the details of an operative laparoscope that he called the 'Galloscope'. On September 12, 1985, he carried out the first laparoscopic cholecystectomy and the concept of 'minimally invasive surgery' was born.<sup>1</sup>

Unfortunately Eric Mühe did not publish his work and consequently was not recognized for performing the first

laparoscopic cholecystectomy. Phillippe Mouret, who was a gynaecologist, documented the first laparoscopic cholecystectomy in 1987.<sup>2</sup> Finally in 1999, Eric Mühe was recognized for his pioneering work and was invited by the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) to give the Storz Lecture that he titled 'The First Laparoscopic Cholecystectomy'.<sup>3</sup>

The widespread adoption of the 'laparoscopic cholecystectomy' resulted in marked improvements in equipment. With increasingly refined technology, it became possible to perform more complex procedures that consequently led to the first report of a laparoscopic colorectal operation being performed in 1991.

The use of laparoscopic surgery with the incorporation of additional programmes such as the 'Enhanced Recovery Program' has resulted in extremely short lengths of postoperative stay. Whereas 10 years ago the average stay following a colorectal resection was 10 to 14 days, papers have been published showing that patients can be discharged safely 23 hours after their surgery.<sup>4</sup>

Despite all the benefits, laparoscopy carries inherent risks. The aim of the paper is to draw attention to these with potential ways of preventing them.

## Principles of safe laparoscopic colorectal surgery

Safe laparoscopic surgery invariably comprises a number of crucial stages, namely:

- positioning of the patient
- gaining access to the abdomen
- insufflation of gas
- maintaining good vision
- placement of additional ports
- use of appropriate laparoscopic instruments and ligation devices
- the limitations of laparoscopy
- closure following laparoscopy.


## Positioning

Initial safe positioning of the patient is necessary. The nature of the operation and region of the abdomen being operated on will influence the position required. Examples are given below.

- Right hemicolectomy and appendicectomy require supine positioning. An adherent mattress such as a large gel or suction mattress is required to prevent patient slippage during head down and right side up positioning, which allows a good view of the caecum, terminal ileum and the ileocolic artery.
- Left-sided colonic resections require the Lloyd Davies position, again with some form of securing mattress, as the head down tilt required is steep as shown in [Figure 1](#). Shoulder supports must be well padded to prevent nerve injury.
- Laparoscopic cholecystectomy requires the patient to be supine and in the reverse Trendelenburg (head up) position, with some left-sided tilt to allow good access to the gallbladder and Calot's triangle.
- Emergency diagnostic laparoscopy, commonly performed to investigate lower abdominal pain in females, requires the Lloyd-Davies position as this allows access to the

**Bruce Levy MBChB (hons) MSc MD FRCS** is a Consultant Colorectal Surgeon at St Richard's Hospital, Chichester, West Sussex, UK. Conflicts of interest: none declared.

**Mohammad Mobasheri BMedSci MBBS MRCS** is a Clinical Research Fellow/Surgical Registrar at Imperial College NHS Trust, London, UK. Conflicts of interest: none declared.



**Figure 1** Table position for steep Trendelenburg.

female pelvic organs should the cause of pain be gynaecological.

### Gaining access to the abdomen

Entry into the abdomen is performed using either an open or closed technique.

- The open 'Hassan' technique can be likened to the now infrequently used 'diagnostic peritoneal lavage' approach. While there are minor surgeon-dependant variations in this approach, the technique itself has become standardized. The approach commences with a small vertical skin incision ( $\pm 1$  cm) below or above the umbilicus depending on the position of the patient's umbilicus in relation to the territory being operated on. The cicatrix of the umbilicus is then elevated using a clip and dissection is performed to expose the abdominal fascia at the base of the cicatrix. The abdominal fascia can then be raised with sutures placed on either side of the midline or with a second clip in order to maximize the distance from the underlying intra-abdominal organs. The fascia and peritoneum is then incised in a controlled and progressive manner in order to maintain a truly open technique as opposed to bluntly pushing in instruments, which can easily cause injury. A 10 mm trocar and port is generally inserted through this opening, though with the advent of smaller scopes 5 mm ports can now be used instead.

Inserting the trocar through the fascia has significant potential for iatrogenic injury and reports of aortic or inferior vena caval injury unfortunately occur more frequently than they should. Damage to the underlying structures can be minimized by elevating the fascia, ensuring the technique is truly open and by pointing trochars down to the pelvis away from the main vessels on insertion.

There are several variations to the described method. Some are due to surgeon choice and others are driven by specific surgical indications.

- Closed method – A Veress needle can be used to create a pneumoperitoneum prior to port insertion. The Veress

needle has a spring-loaded, inner blunt stylet positioned centrally within an outer cannula that has a bevelled end. As the Veress needle is pushed through firm tissue, pressure on the tip pushes the blunt stylet into the shaft of the outer cannula, allowing the outer needle to cut through. As the needle passes through the firm layer, the blunt tipped inner stylet springs out and serves to protect any abdominal viscera from iatrogenic injury by the sharp cannula. Once the Veress needle is in the abdominal cavity, the abdomen is filled with gas before the port is inserted.

Patients that have had previous surgery are prone to adhesion formation and carry an increased risk of visceral injury on entry into the abdomen. Special techniques may be required in such circumstances to permit safe entry. One common technique involves an open cut down at a safe site away from scars, such as Palmers point in the left upper quadrant. An alternative method of entry involves the use of a visual port. The laparoscope is mounted into this port, which is then carefully twisted through the layers of the abdominal wall and into the peritoneal cavity under direct laparoscopic vision.

### Insufflation of carbon dioxide

The abdomen is distended with carbon dioxide in order to create a working space and provide the necessary visualization for the surgery to be performed. Carbon dioxide is used due to its inert, non-flammable properties. In addition, its solubility allows for excretion via the respiratory route. The abdomen is insufflated with carbon dioxide up to a pressure of 12 mm of mercury (mmHg). It is important to be aware that this pneumoperitoneum can cause cardiovascular compromise manifested as hypotension or bradycardia, through a reduction in venous return to the heart. In such a circumstance the pneumoperitoneum should immediately be released and not recommenced until the anaesthetist states that it is safe to do so.

### Good vision

A clear sharp image is essential for safe surgery. The light, which originates from an external light source, is passed to the laparoscope by a fiber-optic cable. The light source can cause serious burns very quickly. It is imperative that the light source does not come in contact with the patient or the drapes when it is on. The safest way to prevent any thermal injury is to switch on the light only after it has been connected to the scope. The laparoscopes used can vary in size and can be either 5 mm or 10 mm in diameter. The 10 mm cameras in general give a better image although the new 5 mm scopes continue to improve.

Camera fog can be very disruptive to the flow of surgery. The fog forms due to condensation on the relatively cold lens. Techniques to avoid fogging include:

- placing the scope in warm water prior to insertion into the abdomen
- using a topical agent to the end of the scope to help reduce condensation.

Laparoscopes come with either a flat end referred to as a zero degree scope or with a bevelled end known as a 30-degree scope. The latter are useful as they have the advantage of being able to view the target area from different angles. This is particularly important in confined working spaces, such as in the pelvis during anterior resection where good views can otherwise be

Download English Version:

<https://daneshyari.com/en/article/3838637>

Download Persian Version:

<https://daneshyari.com/article/3838637>

[Daneshyari.com](https://daneshyari.com)