



Case series

A prospective pilot study on the incidence of post-operative lymphedema in women with endometrial cancer



Elizabeth E. Hopp^{a,*}, Janet L. Osborne^b, Deborah K. Schneider^c, Claudia J. Bojar^d, Denise S. Uyar^d

^a Department of Obstetrics and Gynecology, Medical College of Wisconsin, 9200 West Wisconsin Avenue, Froedtert Medical College Lab Building (FMCLB) 258, Milwaukee, WI 53226, United States

^b Carilion Clinic Gynecologic Oncology, 1 Riverside Circle, Suite 300, Roanoke, VA 24016, United States

^c Department of Surgery, Medical College of Wisconsin, 9200 West Wisconsin Avenue, Suite 3510, Milwaukee, WI 53226, United States

^d Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Medical College of Wisconsin, 9200 West Wisconsin Avenue, Froedtert Medical College Lab Building (FMCLB) 258, Milwaukee, WI 53226, United States

ARTICLE INFO

Article history:

Received 10 October 2015

Received in revised form 19 December 2015

Accepted 23 December 2015

Available online 24 December 2015

Keywords:

Endometrial cancer

Lower-extremity lymphedema

Quality of life

Lymph nodes

Surgical approach

Medical comorbidities

ABSTRACT

To determine the incidence of lower-extremity lymphedema after surgical therapy including lymphadenectomy in endometrial cancer patients using standardized leg measurements. Also, to determine additional risk factors for the development of lymphedema and to study the effect of lymphedema on one's quality of life. In this prospective cohort study, patients with the diagnosis of endometrial cancer who were to undergo definitive surgical management were evaluated pre-operatively and followed post-operatively over the course of two years. Standardized leg measurements were performed by the same individuals at six time-points. Subjects also completed a standardized quality-of-life survey at each time-point. The incidence of lymphedema in 39 women with endometrial cancer using a standardized leg measurement protocol was 12.8% with lymphedema defined as a 20% increase in post-operative leg measurements. There was no significant association between the development of lymphedema and the number of pelvic or para-aortic lymph nodes removed, medical comorbidities, or surgical approach ($p > 0.05$). Of the five patients who met criteria for lymphedema, only one had worsening quality-of-life concerns post-operatively on the FACT-En, version 4, survey. This is the first prospective study using standardized leg measurements to calculate the incidence of post-operative lymphedema in endometrial cancer. Medical comorbidities, surgical approach, number of lymph nodes removed, and location of lymph nodes removed did not appear to affect the development of lymphedema in this cohort. A prospective, multicenter trial is needed to confirm these findings and to further assess the impact of lymphedema on one's quality of life.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Acquired lower-extremity lymphedema occurs after external interruption to the lymphatic system in some gynecologic cancer patients. Endolymphatic pressure increases in the lymphatic vessels and histological changes of the vessels ensue which causes protein-rich fluid to accumulate in the interstitial spaces of the lower-extremity (Mihara et al., 2012). The course of acquired or secondary lymphedema is often chronic and can lead to a worse quality of life in this subset of the gynecologic cancer patient population (Ridner, 2013). The incidence of lymphedema in gynecologic malignancies has been reported as 20% in one systematic review and reported as a range of 6–37% in a retrospective chart review (Cormier et al., 2010; Hareyama et al., 2012). Specifically, patients undergoing lymphadenectomy secondary to endometrial cancer develop lower-extremity edema in 1.2 to 37.8% of cases

(Tada et al., 2009; Abu-Rustum et al., 2006; Todo et al., 2010; Pereira de Godoy et al., 2002).

The risk of lymphedema has previously been correlated with the number of lymph nodes removed. Abu-Rustum et al. (2006) demonstrated that a higher number of lymph nodes removed increase the chance of developing lower-extremity edema. Multiple studies have shown that preservation of the circumflex iliac lymph nodes decreases the incidence of lower-extremity lymphedema (Hareyama et al., 2012; Todo et al., 2010). A retrospective chart review of 286 patients showed that removal of greater than 31 lymph nodes, radiation therapy, and removal of circumflex iliac nodes distal to the external iliac nodes (CINDEIN) increased the risk of lower-extremity lymphedema (Todo et al., 2010).

It is critical to delineate the incidence and risk factors leading to lower-extremity lymphedema as this condition can negatively impact a patient's quality of life physically, mentally, and sexually (Pereira de Godoy et al., 2002; Carter et al., 2013; Beesley et al., 2007). From a recent comprehensive review of the literature, lymphedema associated

* Corresponding author.

E-mail address: ehopp@mcw.edu (E.E. Hopp).

with treatment of gynecologic cancer can cause patients to have self-esteem issues, increased anxiety or fear about their disease, and adversely affect their sexual health (Carter et al., 2013).

The objective of this prospective study was to calculate the incidence of lower-extremity lymphedema after lymphadenectomy in patients with endometrial cancer using standardized leg measurements. This study also examined variables thought to increase the likelihood of lymphedema including medical comorbidities, surgical approach, and number of pelvic and para-aortic lymph nodes removed. We also looked at quality-of-life issues relating to lymphedema.

2. Materials and methods

This prospective cohort study was approved by the Medical College of Wisconsin/Froedtert Hospital Institutional Review Board. To be included, a woman had to have a diagnosis of endometrial cancer and had not yet undergone definitive surgical management. We included subjects regardless of age, race, or body mass index. Women were excluded from the study if they had pre-existing lower extremity edema or were in poor health and could not attend follow-up appointments.

A member of the research team obtained informed consent from the subject. Pre-operatively, information on each subject was collected including age at enrollment, medical comorbidities, and body mass index. Also during the pre-operative period, the subject filled out a quality-of-life survey called the Functional Assessment of Cancer Therapy in Endometrial Cancer Version 4 (FACT-En, version 4). The survey asked about pain in the pelvic area, weight concerns, leg appearance, leg function, and generalized pain of the leg (Fig. 1). The subject then had baseline bilateral leg measurements, which consisted of measuring the width of the leg three inches (7.62 cm) above the lateral malleolus, five inches (12.7 cm) below the patella, and five inches (12.7 cm) above the patella (Fig. 2).

The subject underwent an initial staging surgery performed by a gynecologic oncologist at Froedtert Hospital that included a pelvic and/or para-aortic lymph node dissection, which involved surgical techniques including electrocautery, clips, and sharp dissection. We obtained operative data on the subjects, which included pathology of the tumor, stage, grade, total number of pelvic and para-aortic lymph nodes and surgical approach (laparotomy versus laparoscopy).

The subject was followed for two years post-operatively. The same two nurses took bilateral leg measurements in the exact fashion as performed pre-operatively at three months, six to nine months, one year, and two year time points. At these visits, the subjects were weighed and completed the same quality-of-life survey.

For analysis of the data, subjects who had any type of leg measurements increased by greater than 20% using the standardized leg measurement system by one or two years post-operatively were defined as having “lymphedema.” A chi-square test was performed to look for a significant association between medical comorbidities (congestive

heart failure, diabetes, coronary artery disease, varicose veins, peripheral vascular disease, hypertension) and lymphedema in endometrial cancer. It was also used to determine if a significant association existed between surgical approach and development of lymphedema. A two sample t-test was performed to determine the significance of total pelvic and para-aortic lymph nodes removed in the development of lymphedema. We then qualitatively analyzed the completed quality-of-life surveys.

3. Results

From March 2005 to January 2008, 50 women with endometrial cancer were enrolled in the study. Data collection began in March 2005 and continued until December 2009.

Eleven subjects were removed from the study due to self-withdrawal (two subjects), death (three subjects) and incomplete data in greater than 50% of variables (six subjects).

The remaining 39 women were evaluated at the specified time points post-operatively. The average follow-up visit rate of the 39 subjects at any given time point was 90%.

The incidence of lymphedema in subjects with endometrial cancer as defined by greater than 20% increase in leg measurements post-operatively was 12.8% (5/39 subjects). Of the five subjects who developed lymphedema, three subjects developed lymphedema at one year post-operatively, and two subjects developed lymphedema two years post-operatively. Three subjects developed lymphedema in both legs while two subjects developed lymphedema in one leg. None of the subjects who developed lymphedema had immediate post-operative complications or had recurrence of disease. The five subjects who developed lymphedema had a total of 12 to 41 pelvic and para-aortic lymph nodes removed and had Stage 1A to Stage 1B endometrial cancer. The majority of subjects in the study also had Stage 1A to 1B disease. In patients with lymphedema, the average number of pelvic lymph nodes removed was 14.60 compared to an average number of 15.47 pelvic lymph nodes in those subjects who did not develop lymphedema. In patients with lymphedema, the average number of para-aortic lymph nodes removed was 7.60 compared to an average number of 5.79 para-aortic lymph nodes in those subjects who did not develop lymphedema.

Of the 39 subjects, 44% had significant medical comorbidities previously defined. Twenty-two of 39 subjects (56%) underwent laparotomy and 17 subjects (44%) underwent laparoscopy for the surgical staging procedure. When looking at the covariates in Table 1 of medical comorbidities, surgical approach, and the total number and type of lymph nodes removed, there was no significant association between these covariates and the incidence of lymphedema (p -value >0.05).

When qualitatively looking at worsening quality-of-life issues, the minority of subjects in the study (8/39) had pelvic pain and change in weight, leg pain, leg appearance, or leg function. Table 2 shows that only one of the five patients who developed lymphedema answered “quite a bit” to “very much” to the quality-of-life questions pertaining to lower extremities (Fig. 2) at one year to two years postoperatively.

4. Discussion

We found the incidence of lymphedema in this prospective study of women with endometrial cancer to be 12.8% (5 of 39 patients) over a two-year post-operative interval. We also found that medical history, surgical approach, and the location or number of lymph nodes removed did not influence the development of lymphedema ($p > 0.05$).

To our knowledge, this is the first prospective study looking at the incidence of lymphedema in endometrial cancer using a standardized reporting system of lower leg edema postoperatively. An OVID Medline search was conducted searching the literature from 1950 to the present which contained the terms “lymphedema,” “postoperative,” and “endometrial neoplasms.”

- 1.) I have discomfort or pain in my pelvic area
- 2.) One or both of my legs are swollen and tender
- 3.) I am bothered by a change in weight
- 4.) I have certain parts of my body where I experience significant pain
- 5.) Movement of my leg is painful (Right, Left, Both)
- 6.) I have poor range of movement on this side (Right, Left, Both)
- 7.) My leg on this side feels numb (Right, Left, Both)
- 8.) I have stiffness of my leg in this side

The answer options for all of these questions are a rating of 0 to 4:

- 0 – Not at all
- 1 – A little bit
- 2 – Somewhat
- 3 – Quite a bit
- 4 – Very much

Fig. 1. Questions from the FACT-En, version 4 quality-of-life survey.

Download English Version:

<https://daneshyari.com/en/article/3948512>

Download Persian Version:

<https://daneshyari.com/article/3948512>

[Daneshyari.com](https://daneshyari.com)