



Middle East Fertility Society  
Middle East Fertility Society Journal

www.mefsjournal.org  
www.sciencedirect.com



ORIGINAL ARTICLE

# The effects of serum concentration of androgens, LH and IGF1 in early follicular phase on follicular growth parameters and pregnancy rate



Zahra Raoofi <sup>a,\*</sup>, Farzaneh Hosseini <sup>a,1</sup>, Seyedeh Pegah Parvar <sup>b</sup>,  
Seyedeh Paniz Parvar <sup>c</sup>

<sup>a</sup> Firoozgar Hospital, Iran University of Medical Sciences, Vali-e asr Ave., Tehran 1593748711, Iran

<sup>b</sup> Azad University of Medical Sciences, Iran

<sup>c</sup> Iran University of Medical Sciences, Iran

Received 6 July 2015; revised 25 September 2015; accepted 4 October 2015

Available online 8 January 2016

## KEYWORDS

Androstenedione;  
Dehydroepiandrosterone;  
IGF1;  
LH;  
Ovulation induction;  
Testosterone

**Abstract** Objective: Many studies have showed the role of androgens on the follicular maturation. The present study investigated the effect of serum concentration of androgens, LH and IGF1 in the early follicular phase on the results of the ovulation induction (I/O) and intrauterine insemination (IUI) cycles. Materials and methods: This prospective observational cross-sectional study was carried out in the infertility department of a university hospital in Tehran, Iran. The case's selection was based on the inclusion and exclusion criteria and was nonrandomized. 59 patients under the age of 45 who were candidate for induction ovulation (I/O) or intrauterine insemination were included. The inclusion criteria consist of infertility for at least one year and at least one open tube in HSG. Patients were excluded if they had polycystic ovary syndrome (PCOS) or endometriosis. The serum concentration of androgens (testosterone, dehydroepiandrosterone and androstenedione), LH and IGF1 was measured on the third day of menstruation. Clomiphene and human menopausal gonadotropin (HMG) were drugs of induction ovulation. Human chorionic gonadotropin (HCG) was injected when there was at least one follicle with the size of (18 mm). IUI was done 36 h later for eligible patients and the relation of concentration of androgens, LH and IGF1 with follicular growth parameters and pregnancy rate was analyzed. Results: There was not any statistical significant link between the number and size of follicles with levels of free testosterone, dehydroepiandrosterone, androstenedione, IGF1 and LH. There was not any statistical significant link between the number

\* Corresponding author. Tel.: +98 21 82141201; fax: +98 21 8894.  
E-mail addresses: [Raoofi.z@iums.ac.ir](mailto:Raoofi.z@iums.ac.ir), [drzraoofi@yahoo.com](mailto:drzraoofi@yahoo.com)  
(Z. Raoofi), [farzanehosseini99@yahoo.com](mailto:farzanehosseini99@yahoo.com) (F. Hosseini),  
[pegahprvr@rocketmail.com](mailto:pegahprvr@rocketmail.com) (S.P. Parvar), [panizprvr@gmail.com](mailto:panizprvr@gmail.com)  
(S.P. Parvar).

<sup>1</sup> Tel.: +98 21 82141201; fax: +98 21 8894.

Peer review under responsibility of Middle East Fertility Society.



Production and hosting by Elsevier

<http://dx.doi.org/10.1016/j.mefs.2015.10.002>

1110-5690 © 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of Middle East Fertility Society.  
This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

of follicles in the ovaries and levels of testosterone ( $P = 0.090$  and  $r = 0.223$ ), dehydroepiandrosterone ( $P = 0.642$  and  $r = 0.062$ ) and androstenedione ( $P = 0.526$  and  $r = 0.084$ ), IGF1 ( $P = 0.470$  and  $r = 0.096$ ) and LH ( $P = 0.446$  and  $r = 0.102$ ). There was not any statistical significant link between the mean follicular size with levels of testosterone ( $P = 0.822$  and  $r = 0.03$ ), dehydroepiandrosterone ( $P = 0.733$  and  $r = 0.045$ ) and androstenedione ( $P = 0.526$  and  $r = 0.084$ ), IGF1 ( $P = 0.799$  and  $r = 0.034$ ) and LH ( $P = 0.626$  and  $r = 0.065$ ). Beta Human chorionic gonadotropin (beta-hCG) was positive in 11 patients (18.6%) and negative in 48 patients (81.4%). Serum level of androgen profile, LH and IGF1 in positive BHCG group was lower than negative BHCG group but was not significantly different. Conclusion: It seems that in women who were not affected by PCO, concentration of free testosterone, dehydroepiandrosterone, androstenedione, IGF1 and LH in early follicular phase was not related to follicular growth parameters and pregnancy rate.

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of Middle East Fertility Society. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The androgens as the precursors of estrogens in ovary have the important role in follicular maturation and ovulation. This effect may result in assuming them as the future generation of drugs for induction ovulation. But the exact knowledge of the androgen effects on human follicle maturation does not exist and yet has remained controversial (1,2).

QIN and his colleagues evaluated the baseline level of testosterone during the follicular phase as a predictor of ovarian response and IVF results, and 1260 Chinese women without endometriosis or polycystic ovarian syndrome who underwent IVF cycle, were evaluated. These patients were divided into two groups with limited ovarian reserve (base FSH levels over IU/L 10) (187 patients) and normal ovarian reserve (base FSH levels less than or equal IU/L 10) (1073 people). The results of this study showed that baseline testosterone level in women with limited ovarian reserve is a predictor of large follicles and pregnancy (3).

In a study that was conducted in 2011 by Gleicher and colleagues, he examined the role of androgens in follicular maturation and ovulation in infertile patients. According to the results of this study that reviewed 217 published articles between 2005 and 2011, androgen had an important role in follicular maturation and fertility of women (1). A study conducted by Meldrum and his colleagues examined the role of decreased androgens in the ovarian response to stimulation in older women. Based on these results, increased serum levels of IGF-1, exogenous and local testosterone in the ovaries were associated with increased ovarian response to gonadotropins (4).

Gleicher also investigated the conversion rate of dehydroepiandrosterone in diminished ovarian reserve patients and concluded that women who conceived had a better conversion rate (2). Also because insulin growth factor-1 (IGF-1) is essential for the conversion of androgens to estrogens this study aimed to evaluate the basal concentration of different types of androgens, LH and (IGF-1) in the early follicular phase of the cycle on ovulation induction (I/O) and intrauterine insemination (IUI) results.

## 2. Material and methods

This prospective observational cross-sectional study was carried out in the infertility department of a university hospital

in Tehran, Iran. This study was approved by ethics committee of Iran University of Medical Sciences. The case's selection was based on the inclusion and exclusion criteria and was nonrandomized. All cases provided written informed consent before study registration. 59 patients under the age of 45 who were candidate for induction ovulation (I/O) or intrauterine insemination were included. The inclusion criteria consist of infertility for at least one year and at least one open tube in HSG. Patients were excluded if they had PCOS or endometriosis. Blood samples were collected on the third day of menstruation and the serum level of androgens (testosterone, dehydroepiandrosterone and androstenedione), LH and IGF1 was measured. Also, all the patients underwent transvaginal ultrasound to record and collect the base data. Clomiphene and human menopausal gonadotropin (HMG) were drugs of induction ovulation. Sonographic monitoring was started from the ninth day of the cycle. HCG was injected when there was at least one follicle with the size of (18 mm). IUI was done 36 h later for eligible patients and the records were collected by completing a checklist. The relation of concentration of androgens, LH and IGF1 with follicular growth parameters and pregnancy rate was analyzed by using SPSS software. Chi-2 test and Independent sample *t*-test were used to examine the relation between variables.

## 3. Results

59 patients entered the study that had a mean age of  $32.1 \pm 5.3$  years, with the minimum and maximum ages of 20 and 44 years old. The mean and standard deviation of height, weight and body mass index of patients who entered the study were  $158.3 \pm 13.3$ ,  $66.6 \pm 16.5$  and  $25.4 \pm 4.9$  respectively. The mean and standard deviation of the infertility periods were  $49.1 \pm 36.3$  months that the lowest and highest were 12 and 156 months, respectively. BHCG was positive in 11 patients (18.6%) and negative in 48 patients (81.4%). Table 1 shows the relationship between level of androgen profiles, LH and IGF1 and BHCG results.

There was not any statistically significant link between the number of follicles in the ovaries and levels of testosterone ( $P = 0.090$  and  $r = 0.223$ ), dehydroepiandrosterone ( $P = 0.642$  and  $r = 0.062$ ) and androstenedione ( $P = 0.526$  and  $r = 0.084$ ), IGF1 ( $P = 0.470$  and  $r = 0.096$ ) and LH ( $P = 0.446$  and  $r = 0.102$ ). There was not any statistically

Download English Version:

<https://daneshyari.com/en/article/3966073>

Download Persian Version:

<https://daneshyari.com/article/3966073>

[Daneshyari.com](https://daneshyari.com)