



Short Communication

Energy drink use is associated with alcohol and substance use in eighth, tenth, and twelfth graders

Kathryn Polak ^a, Pamela Dillon ^b, J. Randy Koch ^{a,c}, Willis G. Miller Jr. ^d, Leroy Thacker ^e, Dace Svikis ^{a,f,*}

^a Department of Psychology, Virginia Commonwealth University, Richmond, VA, United States

^b Center for Clinical and Translational Research, Virginia Commonwealth University, Richmond, VA, United States

^c Center for the Study of Tobacco Products, Virginia Commonwealth University, Richmond, VA, United States

^d School of Education, Virginia Commonwealth University, Richmond, VA, United States

^e School of Nursing, Virginia Commonwealth University, Richmond, VA, United States

^f Institute for Women's Health, Virginia Commonwealth University, Richmond, VA, United States

ARTICLE INFO

Article history:

Received 29 March 2016

Accepted 27 June 2016

Available online 29 June 2016

Keywords:

Adolescent health

Caffeine

Energy drinks

Alcohol

Tobacco

Drugs

ABSTRACT

The increasing prevalence of energy drink (ED) use and its link with negative behaviors and adverse health outcomes has garnered much attention. Use of EDs combined with alcohol among college students has been of particular interest. It is unclear if these relationships develop in the context of college, or if similar associations exist in younger individuals. The present study examined associations between ED consumption patterns and other substance use in an adolescent, school-based sample. Participants were $N = 3743$ students attending 8th, 10th or 12th grade in a suburban central Virginia public school system who completed a prevention needs assessment survey in 2012. Chi-square analyses and logistic regressions were used to compare rates of alcohol, tobacco and other drug use across three ED use groups: moderate/heavy (12.6%), light (30.5%), and non-users (57%). Over 40% of the sample reported recent (past month) ED use, with males more likely to report moderate/heavy ED use than females (14.0% and 11.1%, respectively; $p = 0.02$). After adjusting for gender and grade, ED use group predicted lifetime alcohol, tobacco and other drug use (all $p < 0.001$). Moderate/heavy ED users were most likely and ED non-users were least likely to report using each of the 13 substances in the survey, with light ED users intermediate to the other two groups. Moderate/heavy ED users were consistently most likely to report licit and illicit substance use. Additional research is needed to better understand which adolescents are at greatest risk for adverse health behaviors associated with ED use.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

As the popularity of caffeinated energy drink (ED) use continues to increase, so do concerns about possible adverse effects. Many EDs contain substantive amounts of caffeine and heavy caffeine consumption can lead to such adverse effects as agitation/jitteriness, insomnia, tachycardia, and muscle tremors (Reissig et al., 2009; Arria and O'Brien, 2011). Monitoring has proven difficult, however, as many ED products are marketed as conventional foods, which do not require reports of serious adverse events to the FDA.

EDs are particularly popular among college students, with about two-thirds (65.5%) reporting ED use (e.g., Arria et al., 2011). ED use has been linked to heavier drinking, use of illicit and prescription drugs and other risk behaviors (e.g., Arria et al., 2011). Furthermore,

college students who consume EDs combined with alcohol are more likely to use marijuana, meet criteria for alcohol dependence, and engage in other hazardous behaviors (Arria et al., 2010, 2011). ED marketing efforts often focus on young people, and ED consumption has increased among adolescents (Harris and Munsell, 2015; Pennington et al., 2010), with 30 to 50% of teens reporting ED use (Seifert et al., 2011). With increased use has come increased problems, and adolescents make up a significant portion of those negatively impacted by heavy caffeine use. In 2011, nearly 1500 energy-drink-related emergency department visits by individuals aged 12 to 17 were reported (SAMHSA, 2013). In Canada, ED use in adolescents has been linked to alcohol, tobacco and other drug use, as well as sensation-seeking and recent traumatic brain injury (TBI) (Hamilton et al., 2013; Ilie et al., 2015).

Despite increasing prevalence of ED use among adolescents, research has been sparse, with ED use often defined in broad terms (any use, lifetime) that may lack the specificity needed to better understand correlations between ED, other substance use and adverse consequences (Striley and Khan, 2014). Adolescent vulnerability for drug use and problems makes them an important target for substance abuse prevention

* Corresponding author at: Department of Psychology, Institute for Women's Health, Virginia Commonwealth University, 806 W. Franklin St., P.O. Box 842018, Richmond, VA 23284, United States.

E-mail address: dssvikis@vcu.edu (D. Svikis).

efforts. The present study examined associations between ED and other substance use in a sample of 8th, 10th and 12th graders. Specifically, we compared rates of alcohol, tobacco and other drug use across 3 ED consumption groups. We hypothesized that rates of alcohol, tobacco, and other drug use would be highest among the moderate/heavy ED users, followed by light ED users, and finally non-ED users.

2. Method

2.1. Participants

Participants were $N = 3743$ students attending 8th, 10th, or 12th grade in one suburban, central Virginia public school system participating in the 2012 Prevention Needs Assessment Survey (representing over three-fourths (78%) of students in the school district).

2.2. Procedure

Parents of eligible participants were informed about the dates and purpose of the survey via an email. Assurances were given that students and individual schools would not be identified. A waiver of documentation of parental approval was used with passive consent procedures to obtain parental permission for students to complete the survey. On the day of administration, the survey was sent out to randomly selected 8th, 10th, and 12th grade classrooms and all students in attendance whose parents did not opt them out of participation were invited to complete the 45-minute, anonymous, paper-and-pencil survey during the school day. Prior to survey administration, students were reminded that the survey was anonymous and participation was voluntary.

Survey content focused on school experiences, peer interactions, family influences, community environment, risky behaviors, and substance use, including ED, alcohol, tobacco, and other drugs. The survey was developed by an organization dedicated to community-based prevention activities. Prior to administration, survey content was reviewed and approved by the school board, as well as the central Virginia school system and Institutional Review Board.

2.3. Measures

Measures included demographic, ED, alcohol, tobacco, and other drug use variables from the anonymous survey. The ED use item, "On how many occasions (if any) have you had an energy drink (e.g., Monster, Red Bull) during the past 30 days?" (with response options: 0, 1–2, 3–5, 6–9, 10–19, 20–39, and 40+) was used to categorize students into one of three past month ED use groups: non-users (0 times, $N = 1693$; 57%), light ED users (1–5 times, $N = 905$; 30.5%), and moderate/heavy ED users (6 or more times, $N = 374$; 12.6%). For all other substances (see Table 1), participants were labelled "users" if they reported using that particular substance 1 or more times (lifetime).

2.4. Data analysis

Demographic (grade and gender) and other substance use data were compared across the 3 ED use groups using chi-square analyses. Then, logistic regression was used to predict alcohol, tobacco, and other drug use across the three ED use groups. Repeated contrasts were performed for the ED use variable, comparing non-ED users to light ED users, moderate/heavy ED users to non-users, and light ED users to moderate/heavy users. All analyses were carried out using SPSS version 21. For all reported analyses, Bonferroni corrections were applied to reported p -values and confidence intervals to adjust for the multiple comparisons and maintain an overall 0.05 family-wise error rate. Items left blank or with multiple responses were excluded from analyses involving that specific item.

3. Results

3.1. Sample demographics

The sample included $N = 1447$ (38.7%) 8th graders, $N = 1225$ (32.7%) 10th graders, and $N = 1071$ (28.6%) 12th graders. Mean age was 15.4 years ($SD = 1.7$) and 48.3% were female. The sample was predominantly White (59%), followed by African American (33.2%), Hispanic (10.9%), Native American (5.8%), and Asian (5.3%).

3.2. ED use by grade and gender

3.2.1. Any ED use (lifetime)

The percent of adolescents who consumed EDs at least once increased from 64.5% in 8th grade to 73.5% in 10th grade, and 77.4% in 12th grade ($p < 0.01$). There were no gender differences, with nearly three-fourths of both males and females reporting ED use (70.2% vs 71.4%, respectively; NS).

3.2.2. Any ED use (past month)

The percent of adolescents who consumed EDs at least once in the past month did not differ by age/grade level (45.5% vs 40.5% vs 41.8%, respectively; $p = 0.06$). However, males were more likely to report past month use than females (45.0% vs 41%, respectively; $p = 0.03$).

3.2.3. Moderate/heavy ED use (past month)

Moderate/heavy ED use rates also did not vary significantly between 8th, 10th, and 12th graders (13.2%, 11.3%, and 13%, respectively; NS). However, males were more likely to report moderate/heavy ED use than females (14.0% vs 11.1%; $p = 0.02$). Also, male 12th graders were more likely to report moderate/heavy ED use (18.3%) than male 10th and 8th graders (10.4% and 13.5%, respectively; $p < 0.01$). No such relationship was found for females ($p = 0.07$).

Table 1
Alcohol, tobacco, and other drug use by ED use group in one Central Virginia public school system in 2012.

Other substance use (Lifetime)	Overall rate (%)	Moderate/heavy ED use (6+ times) (%)	Light ED use (1–5 times) (%)	Non-ED Use (%)	χ^2 Value (adjusted p -value) ^a
Alcohol	43.6%	63.0%	55.9%	32.9%	159.17 (<0.001)
Cigarettes	28.1%	52.1%	34.0%	19.2%	186.92 (<0.001)
Marijuana	28.7%	48.4%	34.6%	20.0%	148.45 (<0.001)
LSD	5.1%	14.3%	5.3%	3.0%	78.04 (<0.001)
Cocaine	2.7%	9.3%	3.1%	1.1%	78.01 (<0.001)
Inhalants	11.5%	23.6%	14.5%	7.3%	89.97 (<0.001)
Methamphetamine	1.4%	5.5%	1.3%	0.5%	56.18 (<0.001)
Amphetamines	10.0%	20.1%	13.4%	5.7%	87.84 (<0.001)
Sedatives	8.1%	21.7%	8.0%	5.0%	114.38 (<0.001)
Tranquilizers	4.9%	13.5%	5.7%	2.5%	80.02 (<0.001)
Prescription Narcotics	7.8%	17.3%	10.2%	4.3%	81.99 (<0.001)
Heroin	1.1%	5.0%	1.0%	0.4%	53.83 (<0.001)
Ecstasy	3.5%	8.5%	5.4%	1.8%	47.60 (<0.001)

^a Bonferroni correction applied.

Download English Version:

<https://daneshyari.com/en/article/4202311>

Download Persian Version:

<https://daneshyari.com/article/4202311>

[Daneshyari.com](https://daneshyari.com)