


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Case Report

Computed tomography identification of an exophytic colonic liposarcoma

Chung Kuo Chou MD^{a,*}, Sung-Ting Chen MD^b

^a Department of Radiology, Yuan's General Hospital, No. 162, Cheng-Kung 1st Rd, Kaohsiung, Taiwan 80249, Republic of China

^b Department of Pathology, Yuan's General Hospital, Taiwan, Republic of China

ARTICLE INFO

Article history:

Received 29 February 2016

Received in revised form

7 June 2016

Accepted 15 June 2016

Available online 18 July 2016

Keywords:

Liposarcoma

Colon

CT

ABSTRACT

It may be difficult to ascertain the relationship between a large intra-abdominal tumor and the adjacent organs if they are close together. In the current case, a definitive preoperative diagnosis of an exophytic colonic tumor was obtained by the demonstration of obtuse angles between the tumor and colon and by distinct recognition of the mucosa-submucosa of the colonic wall on computed tomography; the accuracy of this preoperative diagnosis was subsequently confirmed by pathologic findings.

© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Liposarcoma is the most common type of soft-tissue sarcoma. It usually occurs in the retroperitoneum and then in the extremities. However, it rarely involves the gastrointestinal tract. In the colon [1–12], it may originate in the submucosa as an intraluminal tumor or in the subserosa as an exophytic mass. In the current case, we present 2 computed tomography (CT) manifestations useful in ascertaining the exophytic nature of the tumor's colon origin: (1) obtuse angles between the tumor and colon and (2) identification of the normal mucosa and submucosa of the colonic wall.

Case report

A 62-year-old male patient presented to the surgical department with the complaint of a palpable mass in the right lower abdomen, which he had been aware of for 2 weeks. He denied having experienced local pain or tenderness, tarry or bloody stool, tenesmus or constipation. The patient received a CT examination that showed a homogeneous soft-tissue tumor, about 14 × 13 × 9 cm in size and abutting the anterior surface of the ascending colon on precontrast scans (Fig. 1). The angles between the tumor and the colon were obtuse, an appearance suggestive of a colonic origin of the

Competing Interests: The authors have declared that no competing interests exist.

* Corresponding author.

E-mail address: wushungxian@kimo.com (C.K. Chou).

<http://dx.doi.org/10.1016/j.radcr.2016.06.005>

1930-0433/© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – (A) Precontrast axial image. A homogeneous soft-tissue tumor (T) was abutting on anterior surface of the ascending colon (C). An obtuse angle (arrows) was noted in both sides between the tumor and colon. This appearance suggested that the tumor was growing out from the colon, instead of compressing from outside toward the colon. (B) Postcontrast axial image. The tumor (T) was heterogeneously enhanced. The submucosa (arrow) was distinct between the inner mucosa (arrowhead) and outer muscularis propria and tumor. This differentiation excluded the possibility of an intraluminal tumor. (C) A magnified view of the rectangle in (B). Arrow, submucosa; arrowhead, mucosa.

tumor. No fat-density tissue was found. An intravenous contrast medium injection showed a heterogeneous enhancement of the tumor. The hypodense parts, not obviously enhanced, had a density of about 15-20 HUs. The fat-density submucosa was readily distinguished between the tumor and the inner mucosa (Figs. 1 and 2). This differentiation indicated that the tumor was most likely originating from a layer beneath the submucosa. The inhomogeneous texture was interpreted to be due to different tumor components and necrosis. The patient was admitted for further treatment. A subsequent colonoscopic examination revealed an external compression of the ascending colon without mucosal lesion or tumor. Based on a preoperative CT diagnosis of a gastrointestinal stromal tumor (GIST) of the ascending colon, the patient received a right hemicolectomy. The gross specimen showed an exophytic soft solid tumor with a myxoid to gelatinous consistency (Fig. 3). The microscopic examination revealed a myxoid liposarcoma in the subserosa with focal penetration through the serosa of the colon. The muscular layer was not invaded. The patient recovered uneventfully and was discharged 10 days later. Despite receiving adjuvant chemotherapy, the patient ultimately succumbed to a widespread intra-abdominal metastasis 14 months later.

Discussion

Primary liposarcoma very rarely occurs in the colon. The reported CT appearance of liposarcoma of the colon included an intra-abdominal fatty-tissue mass [3], a slightly enhanced solid mass with a cystic lesion located in the ascending colon [8], and a large macrolobulated peripherally enhanced mass without radiologic hint of colonic invasion or obstruction [9]. The tumor locations of these 3 cases were all in the subserosa, and none showed CT evidence of colon obstruction. The difficulty in determining the tumor origin may have been because the tumors were closely adhered to the colon. It might have easily been thought to be an intra-abdominal tumor adjacent to the colon. However, in the present case, 2 CT manifestations strongly suggested the possibility of an exophytic tumor originating from a layer beneath the mucosa of the colon. Obtuse angles between the colon and the tumor, which implied an intrinsic tumor bulging outward from the colonic wall instead of an extrinsic tumor compressing inward, were the first CT manifestation. This appearance is similar to an obtuse angle between an extrapulmonary mass and the lung parenchyma [13]. When an extrapulmonary mass bulges toward the lung parenchyma, the pleura is lifted

Download English Version:

<https://daneshyari.com/en/article/4247887>

Download Persian Version:

<https://daneshyari.com/article/4247887>

[Daneshyari.com](https://daneshyari.com)