



Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# The importance of determining surgical indications in cases of lung cancer and interstitial pneumonia with multiple intrapulmonary lymph nodes



Yuki Nakajima<sup>a,\*</sup>, Hirohiko Akiyama<sup>a</sup>, Hiroyasu Kinoshita<sup>a</sup>, Maiko Atari<sup>a</sup>, Mitsuro Fukuhara<sup>a</sup>, Hiroshi Sakai<sup>b</sup>, Hidetaka Uramoto<sup>a</sup>

<sup>a</sup> Division of Thoracic Surgery, Saitama Cancer Center, 780 Komuro, Ina, Kita-adachi-gun, Saitama 362-0806, Japan

<sup>b</sup> Division of Thoracic Oncology, Saitama Cancer Center, 780 Komuro, Ina, Kita-adachi-gun, Saitama 362-0806, Japan

## ARTICLE INFO

## Article history:

Received 25 August 2015

Received in revised form

25 November 2015

Accepted 25 November 2015

Available online 28 November 2015

## Keywords:

Intrapulmonary lymph node

Interstitial pneumonia

Multiple intrapulmonary nodule

Thoracoscopic surgery

## ABSTRACT

**INTRODUCTION:** When considering treatment for pulmonary cancer complicated by interstitial pneumonia, the greatest problem is acute exacerbation. We report two cases of pulmonary cancer complicated by interstitial pneumonia in which multiple intrapulmonary lymph nodes were noted.

**CASE PRESENTATIONS:** Cases 1 and 2 were 76- and 62-year-old males, respectively. Both cases were diagnosed as pulmonary cancer complicated by interstitial pneumonia, and upon chest CT, it was noted that, in addition to the primary lesions, there were multiple nodule shadows under the pleura in the lung lobe, which required identification in order to rule out lung metastasis. In Case 1, the pulmonary nodules were first resected during surgery and a swift diagnosis was made, determining them to be intrapulmonary lymph nodes, which were then operated on curatively. In Case 2, the patient underwent thoracoscopic observation during surgery, the shadows were diagnosed visually as intrapulmonary lymph nodes, and curative surgery was implemented.

**DISCUSSION AND CONCLUSIONS:** The greatest problem in treating pulmonary cancer complicated with interstitial pneumonia is acute exacerbation, wherein, in the absence of any surgical indications, alternative treatment is limited. Thus, contra-indicating surgery for a patient due to a diagnosis of metastasis within the lungs, based only on nodule images, should be avoided. If nodules are noted in the area of the pleura, the possibility exists that these could be intrapulmonary lymph nodes, along with metastasis within the lung, and thoracoscopic surgery should be implemented proactively while keeping these in mind.

© 2015 The Authors. Published by Elsevier Ltd. on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

When treating pulmonary cancer complicated by interstitial pneumonia, it is critical to determine whether or not surgery is indicated [1]. Furthermore, the greatest problem when considering treatment for pulmonary cancer complicated by interstitial pneumonia is acute exacerbation, which presents an added risk in cases of surgery, anti-cancer drug treatment and radiotherapy. At present, radiotherapy and many anti-cancer drugs are prohibited as treatments for interstitial combined pulmonary cancer [2]. As such, in cases of pulmonary cancer complicated by interstitial pneumonia, surgery is the only treatment that offers hope of a cure. In cases such as those in this study, where multiple nodules are found in other lung lobes, the authors consider how to assess the sur-


gical indications for pulmonary cancer complicated by interstitial pneumonia, and whether determination is possible using imaging diagnosis alone. The authors report their experience of two cases of pulmonary cancer complicated by interstitial pneumonia in which multiple lung nodes were identified by proactive thoracoscopy, and were subjected to simultaneous curative surgery. Herein, we report these cases along with observations from the published literature.

## 2. Presentation of cases

### 2.1. Case 1

The patient was a 76-year-old man in whom abnormal shadows were observed on a chest X-ray examination. His medical history included hypertension and hyperlipidemia. He had a smoking history of 20 cigarettes per day for the past 46 years. He worked as a doll maker. Auscultation of the chest resulted in the discovery of fine crackle sounds on both upper dorsal sides. Blood biochemistry

\* Corresponding author. Fax: +81 48 722 1129.  
E-mail address: [yuki@nms.ac.jp](mailto:yuki@nms.ac.jp) (Y. Nakajima).



**Fig. 1.** Image data from Case 1: (A) Left lung S1 + 2 lung cancer (arrow); (B) Left lung S9 nodules; Biopsied nodule; (C) Right lung S10 nodule; (D) Left lung S9 nodule from the pathological specimen. Inside, anthracosis can be seen alongside the intrapulmonary lymph nodes.

tests showed KL-6: 622 U/ml, SPD: 79.4 ng/ml, LDH: 215 IU/l, CEA: 6.9 ng/ml, SCC: 1.8 ng/ml, and CYFRA: 4.0 ng/ml. Blood gas analysis showed pH 7.419, pCO<sub>2</sub>: 36.2 mmHg, and pO<sub>2</sub>: 71.6 mmHg, sO<sub>2</sub>: 93.9%. Respiratory function testing demonstrated VC: 3.27 l, %VC: 111, FEV1: 2.70, % FEV1: 135, and FEV%: 82.8. A chest X-ray examination showed a shadow with pale nodules measuring 20 mm in size with unclear margins in the left upper lung. Both lower pulmonary areas exhibited reticular shadows. A chest contrast CT examination indicated a node shadow measuring 19 mm in size in the left upper lobe S3. Both lower lobes had five nodules measuring from 6 to 12 mm in size. Both pulmonary areas had emphysematous lesions. Dorsal interstitial lesions were also identified in the lower lobes (Fig. 1A–C). Bronchoscopic examination including exfoliative cytodiagnosis from the B3c area resulted in a positive finding for non-small cell lung cancer (NSCLC). Based on the findings of the above examinations, the patient was diagnosed as having pulmonary cancer complicated by interstitial pneumonia, cT1aN0M0, Stage IA or cT4N0M0, Stage III B. In order to determine the surgical indications, nodules from the left lung lower lobe were thoracoscopically biopsied and a swift diagnosis was thus obtained during surgery, based on the policy that if the findings did not indicate lung metastasis, curative surgery would be performed; whereas, if they did indicate metastasis, surgery would not be performed. As blood gas analysis indicated a low pO<sub>2</sub> level, reduction surgery was performed. Thoracoscopic observations of the lungs confirmed that the intrapulmonary lymph nodes were located directly below the lower lobe pleura. The nodules observed on CT images were biopsied after S9 pulmonary wedge resection, and were immediately examined to afford a swift diagnosis; which showed that they were intrapulmonary lymph nodes with no malignancy. Upper division segmentectomy of the left lung and mediastinal lymph node dissection were subsequently performed. The surgery took 201 min and 50 g of blood were lost. Ultimately, the pathological results demonstrated adenocarcinoma, pT1bN0M0,

Stage IA. A left S9 nodule shadows indicated no malignant intrapulmonary lymph nodes (Fig. 1D). The post-surgical progress was good, with the patient leaving the hospital eight days after surgery. At 25 months following surgery, the patient is alive and has had no recurrence.

## 2.2. Case 2

The patient was a 62-year-old male who worked in the transportation field. Abnormal shadows were identified during a chest X-ray examination. His medical history included surgery for stomach cancer. He had a smoking history of 20 cigarettes per day for the past 43 years. Regarding his family history, his father had suffered from pulmonary fibrosis, and his mother had a history of stomach cancer and stroke. Chest auscultation revealed the presence of bilateral fine crackle sounds on the upper dorsal sides. Blood biochemistry findings indicated KL-6: 832 U/ml, SPD: 189 ng/ml, LDH: 228 IU/ml, CEA: 5.2 mg/ml, SCC: 1.4 ng/ml, and CYFRA: 1.8 ng/ml. Blood gas analysis showed pH 7.453, pCO<sub>2</sub>: 39.0 mmHg, pO<sub>2</sub>: 98.9 mmHg, and sO<sub>2</sub>: 97.2%. Respiratory function testing indicated VC: 4.20, %VC: 119.3, FEV1: 2.96, %FEV1: 106, FEV%: 72.3, and %DLCO: 129.7. A chest X-ray examination showed increasing density in the area overlapping the pulmonary artery in the right lung. Both sides showed reticular shadows in the lower pulmonary area. Chest contrast CT examination demonstrated the presence of a shadow measuring 29 mm in size and the presence of nodules with a cavity located directly below the right lung lower lobe S6 pleura. Within the same lung lobe, seven nodule shadows measuring between 2 and 3 mm in size were noted on the other side. Lymph node #11i had expanded to 16 mm. Both sides indicated reticular shadows in the lower dorsal lobe, with interstitial lesions (Fig. 2A–C). Bronchoscopic examination with exfoliative cytodiagnosis demonstrated a positive finding for NSCLC. Based on the above findings, the patient was diagnosed as having pulmonary

Download English Version:

<https://daneshyari.com/en/article/4288594>

Download Persian Version:

<https://daneshyari.com/article/4288594>

[Daneshyari.com](https://daneshyari.com)