



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Arthroscopic treatment of synovial chondromatosis in the ankle joint



Ahmet Ozmeric^a, Nevres Hurriyet Aydogan^a, Onur Kocadal^{a,*}, Talip Kara^a,
Murad Pepe^a, Serap Gozel^b

^a Department of Orthopaedics, Traumatology, Ankara Training, Research Hospital, Ankara, Turkey

^b Department of Pathology, Ankara Training and Research Hospital, Ankara, Turkey

ARTICLE INFO

Article history:

Received 6 May 2014

Received in revised form 22 October 2014

Accepted 23 October 2014

Available online 29 October 2014

Keywords:

Synovial chondromatosis

Osteochondromatosis

Ankle

Arthroscopic treatment

ABSTRACT

INTRODUCTION: Synovial chondromatosis is characterized by the presence of metaplastic cartilage nodules originating from the synovia, bursa and tendon sheaths. Although it is extremely rare in the ankle joint, malignant transformation is possible. The choice of treatment is usually open surgery for excision of loose bodies and synovectomy. Limited data is available concerning arthroscopic approaches.

PRESENTATION OF CASE: A 28-year-old male patient was evaluated for pain and swelling of the right ankle joint. Based on the findings of physical examination and radiographic investigations, arthroscopic surgery was performed due to ankle impingement syndrome. A diagnosis of synovial osteochondromatosis was made following the pathological survey.

DISCUSSION: Synovial chondromatosis is slowly progressive and is considered to be a self-limiting situation. Treatment strategies are decided on according to the patient's complaints, age and disease stage. Open or arthroscopic surgery can be performed. Some advantages of arthroscopic surgery are wide visualization areas, easy access to areas difficult to reach, lower morbidity, no necessity for casting and immobilization, early rehabilitation and quick recovery period.

CONCLUSION: In conclusion, arthroscopic management can be successful in selected patients with synovial osteochondromatosis localized to the ankle joint.

© 2014 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Synovial chondromatosis is a disease with unknown etiology, originating from synovia and characterized by the presence of metaplastic cartilaginous nodules in the synovial cavities, bursa or tendon sheaths.¹ The disease is called synovial osteochondromatosis when the cartilaginous tissue is ossified. Ossification is not seen in approximately 45% of patients.² The disease is commonly seen in men and between the 3rd and 5th decades of life.³ Although the exact etiology is not known; embryologic remnants, infections and trauma are thought to play a role.⁴ The knee, hip and elbow joints are frequently reported to be involved by the condition. However, shoulder and ankle joints are involved extremely rarely.^{1,5–8}

The disease is classified in 3 stages and evaluated according to following criteria: the early stage with intrasynovial differentiation

without loose bodies, the transitional stage by intrasynovial cartilaginous nodules with loose bodies and late stage with multiple loose bodies.⁹ The treatment decision is made according to the patient's age, symptoms and the disease stage.

Nowadays, the arthroscopic approach is frequently preferred for ankle pathologies. The main advantages of the arthroscopic approaches are decreased morbidity, synchronous visualization and treatment feature for intra and extra articular pathologies. The hypertrophic synovia and multiple loose bodies are typical arthroscopic findings.¹⁰

In this case report, we presented an arthroscopically managed adult patient with anteriorly localized right ankle chondromatosis and discussed the potential benefits of arthroscopic surgery.

2. Presentation of case

A twenty-eight year old male patient was admitted to our hospital with decreased range of motion, swelling and increased pain during movement in the right ankle joint. He had experienced progressive pain with reduced range of motion for a year. He had no history of trauma, systemic inflammatory disease or family history

* Corresponding author at: Ankara Training, Research Hospital, Uluçanlar, Ankara, Turkey. Tel.: +90 5323364878.

E-mail address: onurkocadal@gmail.com (O. Kocadal).



Fig. 1. The lateral plain radiographic image of the ankle joint. Multiple calcified foci are seen on the anterior aspect of the ankle joint.

of bone or joint diseases. The physical examination revealed that he had mild tenderness around the anterior ankle joint on palpation with palpable loose bodies. The plantar flexion and dorsiflexion angles were 25° and 5° , respectively. He had increased pain during dorsiflexion. No sign of instability appeared in the ankle joint. Multiple nodules 3–9 mm in diameter with calcifications were located at the anterior aspect of the right ankle on the plain anteroposterior and lateral X-ray images (Fig. 1). Magnetic resonance imaging (MRI) revealed multiple calcified well-circumscribed loose bodies at the same location and synovitis in the ankle joint (Fig. 2). The laboratory tests were within the normal limits and the patient was



Fig. 2. The loose bodies are demonstrated on the anterior aspect of the ankle joint in the sagittal T1-weighted MRI.




Fig. 3. Arthroscopic view of the anterior ankle compartment showing multiple loose bodies.

scheduled for arthroscopic surgery with the diagnosis of anterior impingement syndrome due to right ankle synovial chondromatosis.

The ankle joint was entered via anteromedial and anterolateral arthroscopic portals during spinal anesthesia and tourniquet application. Multiple loose bodies and hypertrophic synovia around the anterior ankle joint were seen (Fig. 3). Arthroscopic partial synovectomy and excision of loose bodies were performed (Fig. 4). The portals were closed primarily after drain insertion. A compressive bandage was applied. The drain was removed in the 1st postoperative day and the active and passive range of motion exercises was started. The patient was allowed partial weight bearing with crutches and at the 2nd week he was mobilized with full weight.

There were multiple cartilaginous loose bodies, with the biggest and smallest dimensions of $0.9\text{ cm} \times 0.7\text{ cm} \times 0.5\text{ cm}$ and $0.4\text{ cm} \times 0.3\text{ cm} \times 0.2\text{ cm}$ in the permanent pathology report respectively (Fig. 5).

The patient's dorsiflexion and plantar flexion degrees were 25° and 30° , respectively, at the end of the 11th postoperative month. No complications were diagnosed in the follow-up period with no recurrence on the plain X-ray images and MRI.



Fig. 4. The macroscopic appearance of the loose bodies after arthroscopic excision.

Download English Version:

<https://daneshyari.com/en/article/4289310>

Download Persian Version:

<https://daneshyari.com/article/4289310>

[Daneshyari.com](https://daneshyari.com)