

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Journal of Asia-Pacific Biodiversity

journal homepage: <http://www.elsevier.com/locate/japb>Journal of
Asia-Pacific
Biodiversity

Short communication

First report of *Vexillum zelotypum* Reeve 1845 (Costellariidae: Neogastropoda) in Korea

Tsukasa Waki, Ronald G. Noseworthy, Kwang-Sik Choi*

School of Marine Biomedical Science (BK21 PLUS), Jeju National University, Jeju, Korea

ARTICLE INFO

Article history:

Received 4 December 2015

Received in revised form

4 January 2016

Accepted 4 January 2016

Available online 8 January 2016

Keywords:

Costellariidae

Gastropoda

intertidal

Jeju

Vexillum zelotypum

ABSTRACT

The costellariid gastropod *Vexillum zelotypum* Reeve 1845a is reported for the first time on Jeju Island. Four Costellariidae species are now part of the island's mollusk fauna. Comparisons are made with the original description and descriptions in the literature; some taxonomic comments are also made. The addition of new record to the island's fauna may be the result of previous insufficient sampling, global warming, and possible changes in the warm currents flowing northward to Japan and Korea.

Copyright © 2016, National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA). Production and hosting by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Jeju Island, 33.10–33.50 North, 126.10–127.0 East, is Korea's largest and southernmost island with an area of about 1,800 km². It is situated about 80 km south of the Korean peninsula. Although cooler water currents, particularly from the eastern area of the Yellow Sea, exert some influence, the warm Tsushima Current, a branch of the Kuroshio flowing from the southwest, gives the island a warm maritime climate with mild winters and hot, humid summers, and an average yearly temperature of approximately 16°C (Chung 2007; Chen 2009). Jeju Island has a rich mollusk fauna, with over 1000 species presently recorded. This fauna is probably a combination of warm temperate (from central Japan to the Yellow Sea) and tropical and subtropical species (from southern Japan to Indo-West Pacific) (Noseworthy et al 2007).

The second author has conducted extensive surveys to enumerate the mollusk fauna, with special attention to its biodiversity, biogeography, and ecology. As a result of those surveys, several species new to the island fauna have been reported (Noseworthy and Choi 2010; Noseworthy et al 2012; Noseworthy et al 2014).

The family Costellariidae, which belongs to the Neogastropoda, consists of about 300 species, and most occur in tropical to

temperate zones. At present, three species have been reported from Korea (Noseworthy et al 2007; Noseworthy and Choi 2010). Here we report the discovery of the costellariid *Vexillum zelotypum* Reeve 1845a, the first record of this species for Korea.

Materials and methods

A mollusk survey was conducted at Yerae-dong, on the south coast of Jeju Island, on September 10, 2015 (Figure 1). The sampling area, a moderately high-energy coast, faces the East China Sea, and is composed mainly of rather flat shelves of basalt interspersed with tidepools of various sizes and depths.

One dead specimen of *V. zelotypum* Reeve 1845a was found under a rock in one of the tidepools. The specimen was washed, dried, and registered at the National Marine Biodiversity Institute of Korea. As the specimen was dead, no anatomical examination could be made. Identification of the specimen was made using the English translation of the original description written in Latin (Reeve 1845a), Reeve (1845b), and the first illustration of the type specimen (Reeve 1845c; Figure 2B), and Okutani (2000).

Systematic account

Order Neogastropoda Wenz, 1938

Family Costellariidae MacDonald, 1860

Genus *Vexillum* Röding, 1798

* Corresponding author. Tel.: +82 64754 3422.

E-mail address: skchoi@jejunu.ac.kr (K.-S. Choi).

Peer review under responsibility of National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA).




Figure 1. Survey site of *Vexillum zelotypum* Reeve 1845a.

***Vexillum zelotypum* Reeve 1845a** (Figure 2A)

Mitra zelotypa Reeve 1845a: 60; Reeve 1845b: 271–272; Reeve 1845c: pl. 38, Figure 325.

Costellaria zelotypa: Adams 1864: 200; Higo and Goto 1993: 261; Okutani 2000: 563, pl. 280, no. 47.

Turricula (Pusia) zelotypa: Tryon, 1882: 182, pl. 54, Figure 558.

Vexillum zelotypa: Cernohorsky, 1970: 55.

Vexillum (Costellaria) zelotypa: Turner, 2001: 69.

Type locality. Not designated; “A very distinct species from Mr. Cuming’s collection, concerning which he has no information as to its locality.” (Reeve 1845c)

Material examined. Jeju Island, Seogwipo-shi, Yerae-dong, 33°14′23.42″ North, 126°23′49.22″ East; September 10, 2015 (1 adult specimen).

Dimensions. Length 21.0 mm; width 10.3 mm.

Description. Shell fusiform; body whorl inflated, especially at shoulders. Strong axial ribs over-ridden by spiral riblets, creating a fenestrated (window-like) sculpture. Ashy-gray with pale red–brown or yellowish band. Axial interstices somewhat darker in color; spire may be spotted with rusty brown. Aperture pinkish and

narrow; columella and aperture sometimes spotted or streaked with purple. Columellar lip with three or four rather strong plicae (Figure 2A).

Habitat. Rock and gravel bottom, from subtidal zone to depth of 30 m (Okutani 2000).

Distribution. Korea (Jeju, new record), From Kii Peninsula, Japan to Indo-Pacific (Okutani 2000).

Discussion

Although somewhat worn, the Yerae-dong specimen agrees well with the English version of the original description of *Mitra zelotypa* in overall shape and sculpture (Reeve 1845b). Reeve (1845b) mentions the longitudinal ribbing which is “strongly, clearly cancellately ridged”, as well as four prominent columellar lirae, and also the pale violet aperture. He displays only the dorsal aspect of the type specimen in his illustration (Reeve 1845c; Figure 2B). The Yerae-dong specimen is less elongate, with a slightly angulate periphery and with only three columellar folds, but exhibits the well-developed cancellate sculpture. Furthermore, this specimen, although somewhat worn, also possesses traces of the

Download English Version:

<https://daneshyari.com/en/article/4394983>

Download Persian Version:

<https://daneshyari.com/article/4394983>

[Daneshyari.com](https://daneshyari.com)