



Taxonomy and systematics

A new species of *Characithecium* (Monogenea: Dactylogyridae) from external surface and gills of two species of *Astyanax* (Ostariophysi: Characidae) in southern Brazil

Especie nueva de Characithecium (Monogenea: Dactylogyridae) de superficie externa y branquias de dos especies de Astyanax (Ostariophysi: Characidae) en el sur de Brasil

Moisés Gallas*, Cláudia Calegari-Marques, Suzana B. Amato

Departamento de Zoología, Instituto de Biociencias, Universidade Federal do Rio Grande do Sul, Caixa Postal 15014, 91501-970 Porto Alegre, Rio Grande do Sul, Brazil

Received 17 August 2015; accepted 29 February 2016

Available online 6 August 2016

Abstract

A new species of *Characithecium*, a genus of monogenean parasites of Neotropical characids, is described from specimens collected on the body surface and gills of *Astyanax* aff. *fasciatus* and *Astyanax jacuhiensis* from Lake Guaíba, State of Rio Grande do Sul, Brazil. The new species is similar to *Characithecium costaricensis*, but differs by having the haptor separated from the body by a long peduncle, and by having an accessory piece composed of 2 subunits in the male copulatory complex – 1 ventral pincer-shaped at the distal end, and 1 dorsal with an expanded rod-shaped proximal end and the distal end containing 3 elongated projections–, and the dorsal bar usually presenting a small median elevation in the anterior margin. This study increases the number of species of *Characithecium* to 7 and expands our knowledge on the diversity of monogenean parasites of characids.

All Rights Reserved © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: Characids; Dactylogyrid; Ectoparasites; Lake Guaíba; Monogeneans

Resumen

Una especie nueva de *Characithecium*, género que comprende especies de parásitos monogénicos de carácidos neotropicales, se describe a partir de especímenes colectados de la superficie corporal y las branquias de los peces *Astyanax* aff. *fasciatus* y *Astyanax jacuhiensis* en el lago Guaíba, estado de Rio Grande do Sul, Brasil. Esta especie nueva es similar a *Characithecium costaricensis* pero se diferencia por tener el haptor separado del cuerpo por un pedúnculo largo y por presentar una estructura accesoria compuesta por 2 subunidades en el complejo copulador masculino –una ventral con forma de pinza en el extremo posterior y otra dorsal con forma de barra en el extremo anterior y posterior, la cual tiene 3 prolongaciones alargadas–. Además, la barra dorsal usualmente presenta una pequeña elevación media en el margen anterior. Este estudio aumenta el número de especies de *Characithecium* a 7 y mejora nuestro conocimiento sobre la diversidad de parásitos monogénicos en carácidos. Derechos Reservados © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: Carácidos; Dactylogyridae; Ectoparásitos; Lago Guaíba; Monogéneos

* Corresponding author.

E-mail address: mgallas88@gmail.com (M. Gallas).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

Introduction

Characithecium Mendoza-Franco, Reina, & Torchin, 2009 was proposed to include *Urocleidoides costaricensis* (Price & Bussing, 1967) Kristky & Leiby, 1972, a species considered as *incertae sedis* by Kritsky, Thatcher, and Boeger (1986) due to the presence of overlapping gonads, mid-ventral vagina, ventral anchor larger than the dorsal, and ventral bar showing a median posterior projection (Mendoza-Franco et al., 2009).

Rossin and Timi (2014) revised the diagnosis of *Characithecium* to include 4 new species collected in Argentina and 1 new combination. The genus currently comprises 6 species: *Characithecium chascomusensis* (Suriano, 1981) Rossin & Timi, 2014, *Characithecium costaricensis* (Price & Bussing, 1967) Mendoza-Franco, Reina, & Torchin, 2009, *Characithecium longianchoratum* Rossin & Timi, 2014, *Characithecium robustum* Rossin & Timi, 2014, *Characithecium quadratum* Rossin & Timi, 2014 and *Characithecium chelatum* Rossin & Timi, 2014.

Urocleidoides astyanacis Gioia, Silva & Artigas, 1988 was described from specimens parasitizing the gills of *Astyanax fasciatus* (Cuvier, 1819) and *Astyanax scabripinnis* (Jenyns, 1842) in Brazil (Gioia, Cordeiro, & Artigas, 1988). This species was also recorded in *Astyanax altiparanae* Garutti & Britski, 2000 by Azevedo, Madi, and Ueta (2007). However, Mendoza-Franco et al. (2009) considered *U. astyanacis* as a junior synonym of *C. costaricensis*. This study describes a new species of *Characithecium* found on the body surface and gills of *A. aff. fasciatus* and *Astyanax jacuhiensis* (Cope, 1894) from Lake Guaíba, State of Rio Grande do Sul, Brazil.

Material and methods

A total of 70 specimens of *A. aff. fasciatus* and 60 of *A. jacuhiensis* were collected with seine nets between March 2012 and November 2013 in Lake Guaíba, municipalities of Guaíba (30°08'28"S, 51°18'53"W) and Barra do Ribeiro (30°17'11"S, 51°18'01"W), State of Rio Grande do Sul, Brazil. Individual fishes were stored in separate plastic bags and kept under refrigeration until necropsy. Each fish had its external surface scraped with a knife to remove mucus, scales and to detach the monogeneans. The gills were removed, put in a jar and shaken at least 50 times with formalin solution 1:4,000. The monogeneans found were processed and the description of their terminology and measurements was conducted according to Gallas, Calegario-Marques, and Amato (2014). Measurements are shown in micrometers (μm) and represent the range followed by the mean, the standard deviation, and the sample size (in parenthesis).

Line drawings were made with a drawing tube using a Nikon E-200 microscope, scanned and prepared using CorelDraw X4® and Adobe's Photoshop® CS2. Ecological parameters follow Bush, Lafferty, Lotz, and Shostak (1997). Holotype and paratypes were deposited in the 'Coleção Helmintológica do Instituto Oswaldo Cruz' (CHIOC), Rio de Janeiro, State of Rio de Janeiro, Brazil, and voucher specimens were deposited in the 'Coleção Helmintológica do Laboratório de Helmintologia'

(CHDZ), Departamento de Zoologia, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, State of Rio Grande do Sul, Brazil. Hosts were identified following Bertaco and Lucena (2010). Representative specimens of the hosts were deposited in the 'Coleção Ictiológica', Departamento de Zoologia, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, State of Rio Grande do Sul, Brazil.

Description

Characithecium triprolatum n. sp. (Figs. 1–9)

Description (based on 22 specimens): Dactylogyridae, Ancyrocephalinae. Body 322–555 (426 ± 87 ; $n = 11$) long, 60–122 (88 ± 20 ; $n = 11$) wide at gonads level. Moderately developed cephalic lobes, 1 pair terminal and the other lateral; cephalic glands clustered in 2 groups in the region of the pharynx. Two pairs of eyes, the anterior pair slightly closer than the posterior one; few accessory granules in the cephalic area. Spherical pharynx, 15–25 (20 ± 3 ; $n = 11$) in diameter; caeca without branches forming a cyclocoel posterior to the gonads. Haptor separated from the body by a peduncle, 55–125 (73 ± 22 ; $n = 10$) long, 35–60 (51 ± 7 ; $n = 10$) wide. Ventral anchor containing an elongate superficial root and a short deep root, both lacking protuberances; slightly straight shaft and elongate point. Ventral anchor 32–40 (37 ± 3 ; $n = 12$) long, base 15–22 (18 ± 3 ; $n = 12$) wide. Ventral bar 5–10 (7 ± 1 ; $n = 10$) long, 17–25 (21 ± 2 ; $n = 10$) wide, showing a regular anterior margin and a posterior margin containing a median projection and expanded ends. Dorsal anchor containing an elongate superficial root and a short deep root both lacking protuberances; slightly straight shaft and elongate point. Dorsal anchor 22–35 (28 ± 4 ; $n = 12$) long, base 10–15 (12 ± 2 ; $n = 12$) wide. Dorsal bar 7.5–12 (10 ± 2 ; $n = 12$) long, 22–35 (27 ± 3 ; $n = 12$) wide, U-shaped, usually showing a small elevation on the anterior margin and a regular posterior margin. Hooks with protruding thumb and dilated shank. Hook pairs 1 and 5, 10–12 (12 ± 1 ; $n = 20$) long, filamentous hook (FH) loop 2–5 (3 ± 1 ; $n = 20$) long. Hook pairs 2, 3, 4, 6, and 7, 15–22 (18 ± 2 ; $n = 56$) long, filamentous hook (FH) loop 4–10 (6 ± 1 ; $n = 56$) long. Slightly overlapped gonads. Testis posterior to the ovary, 22–40 (29 ± 7 ; $n = 5$) long, 10–20 (13 ± 3 ; $n = 8$) wide. Copulatory complex composed of male copulatory organ (MCO) and an accessory piece containing 2 subunits, the dorsal one apparently articulated directly to the MCO. MCO showing a counterclockwise ring coil, ring diameter 12–17 (14 ± 2 ; $n = 8$); base differentiated, possibly fused to the proximal portion of the dorsal subunit of the accessory piece. Accessory piece containing a pincer-shaped ventral subunit on the distal portion, 20–35 (26 ± 4 ; $n = 10$) long; dorsal subunit containing 3 elongations that arise from a rod-shaped proximal portion on the distal portion, 27–45 (36 ± 7 ; $n = 11$) long. One prostatic reservoir, posterior to the base of the MCO. Seminal vesicle present, a dilation of vas deferens; vas deferens looping left caeca, anterior to ovary. Slightly sclerotized mid-ventral vaginal opening. Seminal receptacle present, anterior to ovary. Ovary 37–75 (56 ± 10 ; $n = 9$) long, 12–30 (19 ± 6 ; $n = 9$) wide.

Download English Version:

<https://daneshyari.com/en/article/4461216>

Download Persian Version:

<https://daneshyari.com/article/4461216>

[Daneshyari.com](https://daneshyari.com)