



Research note

Diaphanocephalus galeatus (Nematoda: Diaphanocephalidae), parasite of *Salvator merianae* (Squamata: Teiidae) in southern Brazil

Diaphanocephalus galeatus (Nematoda: Diaphanocephalidae), parásito de *Salvator merianae* (Squamata: Teiidae) en el sur del Brasil

Thainá Dutra Vieira*, Fabiana Fedatto-Bernardon, Gertrud Müller

Laboratório de Parasitologia de Animais Silvestres, Departamento de Microbiologia e Parasitologia, Instituto de Biologia, Universidade Federal de Pelotas, Campus Universitário, caixa postal: 354, CEP 96010-900, Pelotas, Rio Grande do Sul, Brazil

Received 4 August 2015; accepted 22 January 2016

Available online 24 May 2016

Abstract

The gastrointestinal tract of 20 Argentine black and white tegu *Salvator merianae*, was examined in search of nematodes. The diaphanocephalid nematode *Diaphanocephalus galeatus* was identified with a prevalence of 80%. This work contributes to the knowledge of the host's helminth diversity, being the first record in southern Brazil, and presents unpublished photographic records of the species. All Rights Reserved © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: Argentine Black and White tegu; Reptiles; Nematodes; Parasitological indexes

Resumen

Se revisó el tubo gastrointestinal de 20 tegus argentinos *Salvator merianae* en busca de nematodos. Se encontró el diafanocefálico *Diaphanocephalus galeatus* con una prevalencia del 80%. Este trabajo contribuye al conocimiento de la diversidad de helmintos del huésped, siendo el primer registro de la especie en el sur de Brasil y presenta registros fotográficos inéditos de la especie. Derechos Reservados © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: Tegu argentino; Reptiles; Nematodos; Índices parasitológicos

Parasites are among the most diverse and highly specialized organisms; they act as regulators of host populations, maintaining the integrity and stability of ecosystems (Brooks & Hoberg, 2001; Windsor, 1995). Knowledge of endoparasites of wildlife is important for understanding the ecology, natural history, life cycle, and evolution of both parasite and host species (Silva et al., 2008). Studies on lizards in South America demonstrate

that these reptiles act as definitive hosts for several helminth species, mainly nematodes (Ávila & Silva, 2010).

Salvator merianae (Duméril & Bibron, 1839) (Squamata: Teiidae), known as the Argentine black and white tegu or teiú, is considered one of the largest South American lizards. It is found in eastern Bolivia, Argentina, Uruguay, and Brazil (Rio Grande do Sul, Mato Grosso do Sul, Goiás, Pará, Pernambuco, Amazonas, Maranhão, and introduced in the Fernando de Noronha archipelago) (Embert, Fitzgerald, & Waldez, 2010; Loebmann & Quintela, 2009).

This lizard species has a diurnal habit, living in fields, rocky areas, and forests near watercourses. It is omnivorous, with a

* Corresponding author.

E-mail address: thainadutravieira@hotmail.com (T.D. Vieira).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

varied diet consisting of fruits, insects, snails, fish, amphibians, reptiles, rodents, adult birds, nestlings, and eggs (Achaval & Olmos, 2003).

Regarding the helminths parasites of *S. merianae* in South America, until now 6 nematodes species have been recorded: *Cruzia travassosi* Kalil & Vogelsangi, 1932 (Kathlaniidae); *Physaloptera retusa* Rudolphi, 1819 (Physalopteridae); *Diaphanocephalus galeatus* (Rudolphi, 1819) (Diaphanocephalidae); *Spinicauda spinicauda* Olfers, 1819 (Heterakidae); *Physaloptera tupinambae* Pereira, Alves, Rocha, Lima, & Luque, 2012 (Physalopteridae); *Physaloptera bainae* Pereira, Alves, Rocha, Lima, & Luque, 2014 (Physalopteridae); and the cestode *Oochoristica* sp. (Cyclophyllidae) (Ávila & Silva, 2010, 2011; Pereira, Alves, Rocha, Lima, & Luque, 2012, 2014).

In southern Brazil there is no helminthological information related to *S. merianae*. Therefore, the aim of this study is to identify gastrointestinal nematodes inhabiting the intestinal tract of this lizard from municipalities of Rio Grande do Sul State, recording their parasitological indexes: prevalence, abundance and mean intensity of infection.

Twenty *S. merianae* were necropsied, 17 collected (already dead) on highways of southern Brazil from the municipalities of Pelotas (31°46'19" S, 52°20'33" W), Capão do Leão (31°46'3" S 52°26'55" W), and Morro Redondo (31°35'18" S, 52°37'47" W) under permission ICMBio N° 38913-1 issued to Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) (N° 38913-1); the remaining 3 animals were donated by Núcleo de Reabilitação da Fauna Silvestre and Centro de Triagem de Animais Silvestres of Universidade Federal de Pelotas (NURFS-CETAS/UFPel).

The gastrointestinal tracts were sectioned into esophagus, stomach, and small and large intestines. These were opened and washed with a water jet through a 150 µm sieve, and the resulting contents and mucous were examined under an Olympus SZ stereomicroscope. The nematodes were removed, quantified, fixed in AFA, and stored in alcohol 70° GL. They were then mounted on semi-permanent slides with Aman's lactophenol, and viewed on an Olympus CX21 microscope for morphological and morphometric identification according to Freitas and Lent (1938) and Anderson, Chabaud, and Willmott (2009).

The measurements were expressed in millimeters (the mean and standard deviation shown in parentheses) and the photographs taken on an Olympus BX 41 microscope with attached camera system. The parameters calculated were prevalence, abundance, and mean intensity of infection according to Bush, Lafferty, Lotz, and Shostak (1997).

Voucher specimens were deposited in Coleção de Helminthos do Laboratório de Parasitologia de Animais Silvestres do Instituto de Biologia, Universidade Federal de Pelotas (CHLAPASIL/UFPel) (N° 605-611), Rio Grande do Sul, Brazil; Coleção Helminológica do Instituto Oswaldo Cruz (CHIOC) (N° 6735-6737), Rio de Janeiro, Brazil; and Coleção de Invertebrados Museo de La Plata (MLP-He) (N° 6843), La Plata, Argentina.

Diaphanocephalus galeatus (Rudolphi, 1819) Railliet & Henry, 1909 (Fig. 1).

Characterization. Based on 96 specimens (10 males and 10 adult females were measured). Male: total body length 2.69 – 4.5 (3.48 ± 0.58); width (in the half of body) 0.3 – 0.46 (0.39 ± 0.056); length of oral capsule 0.24 – 0.28 (0.26 ± 0.01); width of oral capsule 0.24 – 0.32 (0.28 ± 0.02); esophagus length 0.39 – 0.48 (0.41 ± 0.02); esophagus width (maximum width) 0.17 – 0.27 (0.21 ± 0.02); nervous ring not visible; distance from the excretory pore to cephalic end 0.31 – 0.48 (0.38 ± 0.05); length of the largest spicule 0.58 – 0.84 (0.68 ± 0.08); length of the smaller spicule 0.4 – 0.49 (0.44 ± 0.03); length gubernaculum 0.18 – 0.51 (0.28 ± 0.12); length genital cone 0.29 – 0.3 (0.29 ± 0.007) (visualized in only 2 specimens). Female: total body length 2.19 – 5.62 (4.28 ± 1.01); width (in the half of body) 0.31 – 0.51 (0.44 ± 0.06); length of oral capsule 0.29 – 0.38 (0.32 ± 0.03); width of oral capsule 0.3 – 0.39 (0.34 ± 0.03); esophagus length 0.42 – 0.57 (0.49 ± 0.04); esophagus width (maximum width) 0.19 – 0.30 (0.23 ± 0.03); nervous ring not visible; distance from the excretory pore to cephalic end 0.4 – 0.62 (0.5 ± 0.09); distance from vulvar lip until the rear end 0.52 – 1.2 (0.89 ± 0.24); length of the vagina 0.37 – 1.1 (0.70 ± 0.28); length of ovejector 0.3 – 0.43 (0.36 ± 0.04); distance from the anus to end of tail 0.085 – 0.18 (0.12 ± 0.02); length of eggs 0.06 – 0.087 (0.083 ± 0.10); width of eggs 0.03 – 0.055 (0.041 ± 0.005) (50 eggs were measured).

Diaphanocephalus galeatus was found parasitizing the stomach, small, and large intestine of *S. merianae*. Three hundred forty-six specimens were collected (195 females and 151 males) with a prevalence of 80%, mean abundance 17.3, and mean intensity of infection 21.6 per host. The presented photographic records are unpublished, so they are of important scientific value, considering that Freitas and Lent (1938) presented only illustrations in the description of the parasite.

In Brazil, the nematode was registered previously parasitizing *S. merianae* and *Tupinambis teguixin* (Linnaeus, 1758). In *S. merianae* this nematode species was reported in the states of Mato Grosso (Cuiabá, Chapada dos Guimarães, Poconé), Mato Grosso do Sul (Aquidauana, Serra da Bodoquena, Corumbá and its Nhecolândia district), and Pernambuco (specifically, Fernando de Noronha archipelago where the species was introduced) (Ávila & Silva, 2011). In *T. teguixin*, *D. galeatus* have been collected in the states of Rio de Janeiro, São Paulo, Mato Grosso do Sul, and Rio Grande do Sul, as well as in Argentina and Bolivia (Freitas & Lent, 1938; Vicente, Rodrigues, Gomes, & Pinto, 1993). An additional host species *Tupinambis rufescens* (Günther, 1871) was recorded in Argentina (Spinelli, Fiorito, & Stiebel, 1992).

The morphology and morphometry of males of *D. galeatus* found in this study agree with those described by Freitas and Lent (1938) in *T. teguixin*; however, females studied herein had slightly larger measurements (length and width of the esophagus, distance from the excretory pore to cephalic end, distance from vulvar lip until the rear end, and vaginal length) compared to those presented by the authors (length and width of the esophagus [0.368–0.400 and 0.216–0.240], distance from the excretory pore to cephalic end [0.368–0.416], distance from vulvar lip until the rear end and [0.520–0.872] and vaginal length [0.192–0.304]).

Download English Version:

<https://daneshyari.com/en/article/4461274>

Download Persian Version:

<https://daneshyari.com/article/4461274>

[Daneshyari.com](https://daneshyari.com)