

Author's Accepted Manuscript

The relation of childhood trauma to suicide ideation in patients suffering from obsessive-compulsive disorder with lifetime suicide attempts

Vahid Khosravani, Zoleikha Kamali, Razieh Jamaati Ardakani, Mehdi Samimi Ardestani



PII: S0165-1781(17)30272-X
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.05.032>
Reference: PSY10525

To appear in: *Psychiatry Research*

Received date: 17 February 2017
Revised date: 19 April 2017
Accepted date: 19 May 2017

Cite this article as: Vahid Khosravani, Zoleikha Kamali, Razieh Jamaati Ardakani and Mehdi Samimi Ardestani, The relation of childhood trauma to suicide ideation in patients suffering from obsessive-compulsive disorder with lifetime suicide attempts, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.05.032>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The relation of childhood trauma to suicide ideation in patients suffering from obsessive-compulsive disorder with lifetime suicide attempts

Vahid Khosravani ^{a,*}, Zoleikha Kamali ^b, Razieh Jamaati Ardakani ^c, Mehdi Samimi Ardestani ^d

^aClinical Research Development Center of Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

^bUniversity Technology Malaysia (UTM), Johor Bahru, Malaysia.

^cDepartment of Psychology, Ardakan University, Ardakan, Yazd, Iran.

^dDepartments of Psychiatry, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

*Corresponding author. E-mail address: vahid.psy@gmail.com

Abstract

The aim of this study was to evaluate the relations of childhood trauma (CT) and obsessive-compulsive (OC) symptom dimensions to suicide ideation in patients with obsessive-compulsive disorder (OCD). Seventy OCD outpatients with lifetime suicide attempts and 60 controls were included. Participants completed the Scale for Suicide Ideation (SSI), the Childhood Trauma Questionnaire-Short Form (CTQ-SF), the Yale-Brown Obsessive Compulsive Scale (Y-BOCS), the Dimensional Obsessive-Compulsive Scale (DOCS) and the Depression Anxiety Stress Scales (DASS-21). Among OCD patients, 97.1% had current suicide ideation. OCD patients revealed higher scores on CT, suicide ideation, depression and anxiety than controls. The CT history of sexual abuse (SA) and OC symptom dimension of unacceptable thoughts explained suicide ideation. It was concluded that SA and unacceptable thoughts may contribute to high suicidality and have important implications for the assessment and treatment of suicide risk in OCD patients with lifetime suicide attempts.

Keywords: Childhood trauma, Obsessive-compulsive disorder, Suicide ideation, Lifetime suicide attempt

Download English Version:

<https://daneshyari.com/en/article/4933186>

Download Persian Version:

<https://daneshyari.com/article/4933186>

[Daneshyari.com](https://daneshyari.com)