

Author's Accepted Manuscript

The prevalence and outcomes of exposure to potentially traumatic stressful life events compared across patients with bipolar disorder and unipolar depression

Stacey McCraw, Gordon Parker



PII: S0165-1781(16)31974-6
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.06.070>
Reference: PSY10617

To appear in: *Psychiatry Research*

Received date: 23 November 2016
Revised date: 17 May 2017
Accepted date: 18 June 2017

Cite this article as: Stacey McCraw and Gordon Parker, The prevalence and outcomes of exposure to potentially traumatic stressful life events compared across patients with bipolar disorder and unipolar depression, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.06.070>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The prevalence and outcomes of exposure to potentially traumatic stressful life events compared across patients with bipolar disorder and unipolar depression.

Stacey McCraw, BScience (hons), MPsych (Clin)^{a,b}, Gordon Parker, M.D., Ph.D^{a,b*}

^a*School of Psychiatry, the University of New South Wales, Sydney, NSW, Australia.*

^b*The Black Dog Institute, Sydney, NSW, Australia.*

*Corresponding author: Black Dog Institute, Prince of Wales Hospital, Randwick 2031, Sydney, Australia. s.mccraw@unsw.edu.au

Abstract

This study examined the profile of stressful life experiences in patients with unipolar depression (UP) compared to bipolar disorder (BP), including their subsequent psychological impact and affirmation of post-traumatic stress symptoms. We studied 747 tertiary patients diagnosed with either a UP (n= 413) or BP condition (n = 334) according to a structured research interview and psychiatrist assessment. An online assessment collected additional study variables. Results showed that despite being younger in age, the BP group were more likely to have experienced 8 of the stressful life events, including childhood and adulthood abuse. However, both groups judged the consequential 'impact' of those events similarly. The BP group was significantly more likely than the UP group (45% vs. 36%) to affirm exposure to an extremely stressful event across their lifetime, resulting in 26.3% of the BP and 14.5% of the UP group affirming DSM-IV criteria for lifetime PTSD. The onset of DSM-IV-defined post-traumatic symptoms tended to occur in adulthood for both groups, with trauma often following the onset of the mood disorder. Thus, BP patients displayed greater vulnerability toward traumatic experiences and anxiety disorders, but reported a similar psychological impact in response to each stressor, compared with the UP group.

Keywords: mood disorders; depression; mania; trauma; comorbidity; assessment

Introduction

Download English Version:

<https://daneshyari.com/en/article/4933222>

Download Persian Version:

<https://daneshyari.com/article/4933222>

[Daneshyari.com](https://daneshyari.com)