

Author's Accepted Manuscript

The Columbia-Suicide Severity Rating Scale:
Associations between Interrupted, Aborted, and
Actual Suicide Attempts among Adolescent
Inpatients

Ryan M. Hill, Claire E. Hatkevich, Iram Kazimi,
Carla Sharp



PII: S0165-1781(17)30546-2
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.06.014>
Reference: PSY10561

To appear in: *Psychiatry Research*

Received date: 28 March 2017
Revised date: 30 May 2017
Accepted date: 5 June 2017

Cite this article as: Ryan M. Hill, Claire E. Hatkevich, Iram Kazimi and Carla Sharp, The Columbia-Suicide Severity Rating Scale: Associations between Interrupted, Aborted, and Actual Suicide Attempts among Adolescent Inpatients, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.06.014>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

RUNNING HEAD: INTERRUPTED AND ABORTED ATTEMPTS

The Columbia-Suicide Severity Rating Scale: Associations between Interrupted, Aborted, and
Actual Suicide Attempts among Adolescent Inpatients

Ryan M. Hill, PhD^{a*}, Claire E. Hatkevich, MA^b, Iram Kazimi, MD^a, & Carla Sharp, PhD^b

* Address correspondence to:

Ryan M. Hill
Instructor, Department of Psychiatry and Behavioral Sciences
The University of Texas Health Science Center at Houston
1941 East Road, Suite 2316
Houston, TX 77054
Ryan.M.Hill@uth.tmc.edu
Office: 713-486-2706

ABSTRACT

The Columbia—Suicide Severity Rating Scale divides suicide attempt behaviors into actual, interrupted, and aborted attempts, but limited data have been reported regarding associations between interrupted, aborted, and actual attempts. This study provided initial data on the ability of interrupted and aborted attempts to estimate the frequency of actual suicide attempts. Participants were adolescent psychiatric inpatients (59.9% female), 12-17 years (mean=14.73, SD=1.62). Results suggest that interrupted and aborted suicide attempts are associated with the frequency of actual suicide attempts, controlling for suicidal ideation and depressive symptoms. Future research should evaluate whether interrupted and aborted attempts prospectively predicting actual suicide attempts.

Keywords: risk assessment, suicide-related behaviors, incremental validity

Download English Version:

<https://daneshyari.com/en/article/4933233>

Download Persian Version:

<https://daneshyari.com/article/4933233>

[Daneshyari.com](https://daneshyari.com)