

## Author's Accepted Manuscript

Quality of life in individuals with attenuated psychotic symptoms: Possible role of anxiety, depressive symptoms, and socio-cognitive impairments

Tsutomu Takahashi, Yuko Higuchi, Yuko Komori, Shimako Nishiyama, Mihoko Nakamura, Daiki Sasabayashi, Yumiko Nishikawa, Tomiki Sumiyoshi, Michio Suzuki



PII: S0165-1781(17)30922-8

DOI: <http://dx.doi.org/10.1016/j.psychres.2017.08.024>

Reference: PSY10739

To appear in: *Psychiatry Research*

Cite this article as: Tsutomu Takahashi, Yuko Higuchi, Yuko Komori, Shimako Nishiyama, Mihoko Nakamura, Daiki Sasabayashi, Yumiko Nishikawa, Tomiki Sumiyoshi and Michio Suzuki, Quality of life in individuals with attenuated psychotic symptoms: Possible role of anxiety, depressive symptoms, and socio-cognitive impairments, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.08.024>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Quality of life in individuals with attenuated psychotic symptoms: Possible role of anxiety, depressive symptoms, and socio-cognitive impairments

Tsutomu Takahashi<sup>a,\*</sup>, Yuko Higuchi<sup>a</sup>, Yuko Komori<sup>a</sup>, Shimako Nishiyama<sup>a</sup>, Mihoko Nakamura<sup>a</sup>, Daiki Sasabayashi<sup>a</sup>, Yumiko Nishikawa<sup>a</sup>, Tomiki Sumiyoshi<sup>b</sup>, Michio Suzuki<sup>a</sup>

<sup>a</sup> *Department of Neuropsychiatry, University of Toyama Graduate School of Medicine and Pharmaceutical Sciences, Toyama, Japan*

<sup>b</sup> *Department of Clinical Epidemiology, Translational Medical Center, National Center of Neurology and Psychiatry, Japan*

\*Corresponding author: Tsutomu Takahashi, M.D.

Department of Neuropsychiatry, University of Toyama Graduate School of Medicine and Pharmaceutical Sciences, 2630 Sugitani, Toyama 930-0194, Japan

Tel: +81-76-434-2281

Fax: +81-76-434-5030

E-mail: tsutomu@med.u-toyama.ac.jp

### Abstract

Individuals with Clinical High-Risk state for Psychosis (CHR-P) are reported to exhibit impaired quality of life (QOL) similar to that observed in schizophrenia, but its determinants remain unclear. We investigated the QOL of 33 subjects with CHR-P, 45 patients with schizophrenia, and 63 healthy subjects using the Quality of Life Scale (QLS). The CHR-P and schizophrenia groups were administered the Brief Assessment

Download English Version:

<https://daneshyari.com/en/article/4933513>

Download Persian Version:

<https://daneshyari.com/article/4933513>

[Daneshyari.com](https://daneshyari.com)