



Short communication

## A robust chemo-enzymatic lactone synthesis using acyltransferase from *Mycobacterium smegmatis*

A. Drożdż<sup>a</sup>, U. Hanefeld<sup>b</sup>, K. Szymańska<sup>c</sup>, A. Jarzębski<sup>c,d</sup>, A. Chrobok<sup>a,\*</sup><sup>a</sup> Silesian University of Technology, Department of Chemical Organic Technology and Petrochemistry, Krzywoustego 4, 44-100 Gliwice, Poland<sup>b</sup> Gebouw voor Scheikunde, Biokatalyse, Afdeling Biotechnologie, Technische Universiteit Delft, Julianalaan 136, 2628BL Delft, The Netherlands<sup>c</sup> Silesian University of Technology, Department of Chemical and Process Design, Strzody 7, 44-100 Gliwice, Poland<sup>d</sup> Institute of Chemical Engineering, Polish Academy of Sciences, Bałtycka 5, 44-100 Gliwice, Poland

## ARTICLE INFO

## Article history:

Received 4 March 2016

Received in revised form 29 March 2016

Accepted 31 March 2016

Available online 13 April 2016

## Keywords:

Acyltransferase

Baeyer-Villiger oxidation

Lactones

Hydrogen peroxide

## ABSTRACT

The new application of acyltransferase, isolated from *Mycobacterium smegmatis* for the chemo-enzymatic Baeyer-Villiger oxidation of cyclic ketones to lactones was demonstrated. Acyltransferase exhibited high activity, and high stability under harsh reaction conditions, like oxidation with 60% aq. H<sub>2</sub>O<sub>2</sub> at 45 °C. This paves the way to a novel robust chemo-enzymatic method for lactone synthesis with high yields.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Baeyer-Villiger (BV) oxidation of cyclic ketones remains one of the most important protocols for the synthesis of lactones [1] with applications in the synthesis of antibiotics, steroids, pheromones and polymers [2].

Biocatalysis offers an environmentally friendly alternative for typical BV transformations which make use of organic percarboxylic acids [1]. The oxidation of ketones to lactones can be carried out using either Baeyer-Villiger monooxygenases (BVM) [3,4] or hydrolases such as *Candida antarctica* lipase B (CALB) [5–8] and esterases [9]. In the first case, using BVM, the enzyme catalyses the oxidation of ketones with oxygen and nicotinamide adenine dinucleotide phosphate NADPH as a source of electrons [3,4] to obtain lactones with enantioselectivities up to 100%. However, as BVMs are relatively expensive and poorly stable, the second option is preferred, in which the enzyme boosts the *in-situ* oxidation of long- or medium-chain carboxylic acids or ethyl acetate with hydrogen peroxide to generate peracids used to oxidise ketones to lactones in the second (chemical) step [5–8]. Clearly, this single-pot chemo-enzymatic approach is far more elegant and attractive commercially.

The search for enzymes that are active and stable in this reaction was crucial and the results were already reported in several papers. Hydrogen peroxide and the peracid generated during the reaction are

inactivating reagents for enzymes. To improve the enzyme stability and to allow the effective recycling of enzymes the immobilization techniques can be used [10]. The most studied was Novozyme-435, *i.e.* CALB immobilized on acrylic resin [5]. But very efficient cross-linked enzyme aggregates (CLEAs) with perhydrolase, in a chemo-enzymatic reaction have been also proposed [7]. Our previous studies demonstrated very good activity of CALB immobilized on siliceous materials with organosilanes terminated with alkyl groups [6] or in ionic liquids [8].

The acyltransferase, more recently isolated from *Mycobacterium smegmatis* (MsACT), appeared to demonstrate in the presence of hydrogen peroxide a perhydrolysis: hydrolysis activity ratio 50-fold greater than the common lipases (CALB included) [11]. It was immobilized and used as a paint additive, catalyzing peracid formation even under those conditions [12,13]. Therefore, MsACT emerges as a natural candidate to replace lipases also in the chemo-enzymatic Baeyer-Villiger oxidation of cyclic ketones to lactones.

In this light, we deemed it important to test the MsACT performance in this reaction and to compare it with those shown by free CALB or immobilized as Novozyme-435. Needless to say that the reaction under study is of a major practical significance.

## 2. Experimental

## 2.1. Materials

30% and 60% aq. H<sub>2</sub>O<sub>2</sub> and UHP were purchased from Acros Organics. Cyclic ketones and native CALB [5,000 LU/G] were purchased from

\* Corresponding author.

E-mail addresses: [u.hanefeld@tudelft.nl](mailto:u.hanefeld@tudelft.nl) (U. Hanefeld), [katarzyna.szymanska@polsl.pl](mailto:katarzyna.szymanska@polsl.pl) (K. Szymańska), [andrzej.jarzebski@polsl.pl](mailto:andrzej.jarzebski@polsl.pl) (A. Jarzębski), [anna.chrobok@polsl.pl](mailto:anna.chrobok@polsl.pl) (A. Chrobok).

Sigma Aldrich. Novozyme-35 was donated by Novozymes. Enzyme MsAcT was produced and the activity determined as described earlier [14].

## 2.2. General method for Baeyer-Villiger oxidation

The ketone (0.25 mmol) and 0.5 ml of ethyl acetate (5.09 mmol) were introduced into a 25 ml round-bottom flask and the contents of the flask was shaken. Next, 4 mg of MsAcT was introduced, and 60% aq.  $\text{H}_2\text{O}_2$  (0.50 mmol) was added dropwise. The flask was sealed with a septum and mixed in a thermostated shaker ( $\pm 0.5^\circ\text{C}$ ) with orbital stirring at 250 rpm at  $35^\circ\text{C}$  for 2 h to 5 days, depending on the reaction rate. Periodically, 10  $\mu\text{l}$  of the sample diluted with 0.7 ml of dichloromethane was collected during the reaction to monitor the progress of the reaction utilising GC (Perkin Elmer Clarus 500 chromatograph with SUPELCOWAX™ 10 column (30 m  $\times$  0.2 mm  $\times$  0.2  $\mu\text{m}$ ) with *n*-decane as an external standard.


The structure of the products were confirmed using GC–MS analysis (Agilent Gas Chromatograph 7890C equipped with a HP-5 MS column (30 m  $\times$  0.25 mm  $\times$  0.25  $\mu\text{m}$ ; MS Agilent 5975C, EI ionization 70 eV, and the results were compared to NIST/EPA/NIH Mass Spectral Library.

## 3. Results and discussion

As can be seen from Scheme 1 the applied chemo-enzymatic method of lactone synthesis involves MsAcT as the biocatalyst of peracid formation, hydrogen peroxide as oxidant and ethyl acetate as both the peracid precursor and solvent. As a model ketone 2-methylcyclohexanone was used.

The experiments were performed under the reaction conditions recommended for this reaction ( $25^\circ\text{C}$ , molar ratio of ketone to 30% aq.  $\text{H}_2\text{O}_2$  1: 2) [6]. At first, they aimed at discriminating the regions of specific kinetic control of the chemo-enzymatic Baeyer-Villiger reaction. They were made by varying the amount of MsAcT in the range of 2–7 mg, while keeping the amount of a model ketone constant (0.25 mmol; 2-methylcyclohexanone). For a fixed value of molar ratio of the ketone to 30% aq.  $\text{H}_2\text{O}_2$  (1:2) and an excess of ethyl acetate the rate of lactone formation at  $25^\circ\text{C}$  appeared to depend on the biocatalyst content, provided it was  $\leq 4$  mg (Fig. 1). Those findings delineated the region of effective control of ketone oxidation by the created peracid (4–7 mg of MsAcT). Since differences in the reaction courses carried out using 4 and 7 mg of the enzyme per 0.25 mmol of ketone appeared to be small, therefore all further studies were performed using 4 mg of the enzyme.

At this stage we also checked the influence of ethyl acetate and the possibility to replace its excess with a buffer. These studies showed that the 5.09 mmol of ethyl acetate (0.5 ml) per 0.25 mmol of ketone (approximately, only 5% of ethyl acetate is consumed for the reaction) is the most effective and that the addition of the buffer has insignificant influence on the reaction rate (Fig. 2). The latter phenomenon could be explained by an extensive hydrophobicity of MsAcT active center [11, 13–16]. In all cases, with the addition of buffer or not, the reaction system was always biphasic according to the use of water solution of  $\text{H}_2\text{O}_2$  and the creation of water molecule as a by-product in the reaction.



Scheme 1. Model chemo-enzymatic Baeyer-Villiger oxidation studied in this work.




Fig. 1. Effect of MsAcT content on the BV oxidation of 2-methylcyclohexanone (0.25 mmol) with 30% aq.  $\text{H}_2\text{O}_2$  (0.50 mmol) in ethyl acetate (5.09 mmol, 0.5 ml) at  $25^\circ\text{C}$ .

The effect of the buffer was also checked for two forms of CALB lipase; a native (liquid) and its immobilized form - Novozyme 435. As can be seen from Fig. 3 the presence of water appeared to exert a strong negative effect on their activity, regardless the biocatalyst form. This could be ascribed to two factors: (i) specific structure of the lipase, especially the presence of hydrophobic polypeptide chain (lid or flat), isolating its active centre from the reaction medium [17], (ii) hydrolysis of ethyl acetate which lowered the pH and this brought the reaction to a halt.

Further studies aimed to determine the effect of oxidising agent (30 and 60% aq.  $\text{H}_2\text{O}_2$ , urea hydrogen peroxide UHP). They showed that the most reactive is 60% aq.  $\text{H}_2\text{O}_2$  (Fig. 4). A twofold molar ratio of 60% aq.  $\text{H}_2\text{O}_2$  to the ketone was large enough to ensure optimum kinetics, i.e. very similar in value to that obtained using a fourfold excess of 30% aq.  $\text{H}_2\text{O}_2$ . It is noteworthy, that the idea of using 60% aq.  $\text{H}_2\text{O}_2$  was also aimed to probe the MsAcT performance under harsh reaction conditions, since the exposure of CALB to high concentration of aq. hydrogen peroxide resulted in its complete deactivation [18]. Anhydrous UHP appeared to be poorly reactive under these reaction conditions. The efforts for the isolation of enzyme after the reaction were unsuccessful. The use of enzyme immobilization methods may produce improvements in the enzyme performance, as was already described in the literature. [19,20].




Fig. 2. The influence of the solvent on the BV oxidation of 2-methylcyclohexanone (0.25 mmol) with 30% aq.  $\text{H}_2\text{O}_2$  (0.50 mmol) and MsAcT (0.004 g) at  $25^\circ\text{C}$ .

# دانلود مقاله



<http://daneshyari.com/article/49771>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات