



Short Communication

Trends in average days' supply of opioid medications in Medicaid and commercial insurance

Ali Bonakdar Tehrani^a, Rachel Mosher Henke^a, Mir M. Ali^{b,*}, Ryan Mutter^c, Tami L. Mark^d^a Truven Health Analytics, An IBM Company, United States^b Substance Abuse & Mental Health Services Administration, Center for Behavioral Health Statistics & Quality, 5600 Fishers Lane, Rockville, MD 20852, United States^c Substance Abuse & Mental Health Services Administration, Center for Behavioral Health Statistics & Quality, Rockville, MD, United States^d RTI International, Washington D.C., United States

HIGHLIGHTS

- We examined trends in average days' supply of opioid medications for Medicaid and commercially insured population.
- Average days' supply for all drugs except morphine increased for both Medicaid and commercially insured population.
- More outreach education is needed to convince providers that opioids should be prescribed cautiously.

ARTICLE INFO

Keywords:

Days' supply
Opioid
Medicaid
Commercial insurance

ABSTRACT

Objectives: To calculate trends in adult average days' supply for six commonly prescribed opioids: hydrocodone, hydromorphone, morphine, oxycodone, oxycodone, oxycodone, and tapentadol to assess whether physicians changed prescribing practices at the time of the intensifying epidemic.

Methods: We used 2005–2015 Truven Health MarketScan Commercial Claims and Encounters data to measure trends in opioid average days' supply among commercially insured individuals and 2005–2014 MarketScan Multi-State Medicaid data to measure trends in opioid average days' supply among Medicaid beneficiaries.

Results: For Medicaid, we found an increase in days' supply for all drugs except morphine. The largest percentage increase was for oxycodone, which increased 4.5 days (37%). Opioid days' supply for individuals with commercial insurance exhibited similar but steeper trends. The largest increase was also for oxycodone, which increased 6 days (56%). Between 2013 and 2015, when the opioid epidemic had begun to be widely publicized, there was no decline in the median days supplied for any of the opioids.

Conclusions: Our results find that days' supply of opioids are increasing despite public health campaigns and media attention on the risks of opioid prescribing. More effective interventions to curb opioid prescribing are needed to reverse these trends.

1. Introduction

Increasing abuse of opioid pain medications and mortality due to opioid overdose is a major public health concern (Dart, Surratt, Cicero, et al., 2015). In 2014, there were about 19,000 opioid-related deaths, representing a 20% increase from 2013 (CDC, 2016). Although there has not been a dramatic change in the amount of pain that Americans report over the past two decades, the number of opioid prescription drugs sold in the United States has nearly quadrupled since 1999 (CDC, 2016; Chang, Daubresse, Kruszewski, & Alexander, 2014; Daubresse, Chang, Yu, et al., 2013). An estimated 20% of patients presenting to

physician offices with noncancer pain symptoms or pain-related diagnoses (including acute and chronic pain) receive an opioid prescription (Daubresse et al., 2013). In 2012, health care providers wrote 259 million prescriptions for opioid pain medication (Paulozzi, Mack, & Hockenberry, 2014).

In March 2016, the CDC issued opioid prescribing guidelines recommending that opioids should not be the first-line therapy for chronic pain because of their risks and limited evidence of long-term efficacy (Dowell, Haegerich, & Chou, 2016). Further, the CDC recommended that opioid prescribing for acute pain should not be provided “just in case” and should be limited to the expected duration of

* Corresponding author at: Substance Abuse & Mental Health Services Administration, Center for Behavioral Health Statistics & Quality, 1 Choke Cherry Road, Rockville, MD 20852, United States.

E-mail address: mir.ali@samhsa.hhs.gov (M.M. Ali).

<http://dx.doi.org/10.1016/j.addbeh.2017.08.005>

Received 6 April 2017; Accepted 10 August 2017

Available online 19 August 2017

0306-4603/ Published by Elsevier Ltd.

pain severe enough to require opioids (typically 3–7 days) to minimize unintentional initiation of long-term opioid use.

Guidelines for the number of pills prescribed (or days supplied) are intended to minimize unintentional or intentional diversion, which is a greater issue when there are leftover pills. In an effort to reduce the risks of addiction and diversion, states have implemented limits on the number of days supplied under a prescription (CDC, 2015). For example, Connecticut recently passed legislation limiting prescriptions for pain medications to a 7-day supply except for certain circumstances, such as chronic pain, cancer pain, and palliative care (Connecticut General Assembly, 2016).

Despite this legislation and the awareness that the provision of lengthy prescriptions may lead to abuse and diverted pills, little is known about the trends in days' supply of prescribed opioids. Reducing the number of pills prescribed is an important component of addressing the opioid epidemic, making it important to track trends in the number of days supplied for opioid medications. In this study, we use claims data to calculate trends in the average adult days' supply for six commonly prescribed opioids: hydrocodone, hydromorphone, morphine, oxycodone, oxymorphone, and tapentadol.

2. Materials and methods

2.1. Databases and sample population

We used two Truven Health MarketScan® Databases for this analysis: 1) 2005–2015 Commercial Claims and Encounters Database, which includes insurance claims from employees and their dependents covered by large, self-insured employers and by regional health plans; 2) 2005–2014 Multi-State Medicaid Database, which contains the pooled healthcare experience of approximately 6 million Medicaid enrollees. The sample population was limited to adults aged 18 to 64, excluding people who were diagnosed with cancer. Individuals dually eligible for Medicare and Medicaid were excluded from the Medicaid analysis. The Medicaid sample was 30% male and 70% female, and the average age was 40.5 years. The Commercial sample was 42% male and 58% female, with an average age of 46.2 years.

2.2. Opioid drugs

We identified all oral or tablet forms of prescriptions filled for hydrocodone, hydromorphone, oxycodone, oxymorphone, tapentadol, and morphine by adults aged 18 to 64, excluding prescriptions to patients diagnosed with cancer and those with invalid information. These medications were chosen because (1) they are the most frequently prescribed opioid pain medicines; (2) they were available throughout the time period, thereby allowing trends to be examined; and (3) they were oral pills for which days supplied is relevant. We included combination drugs, which we grouped according to the opioid ingredient. For example, the hydrocodone group included hydrocodone/acetaminophen, hydrocodone/ibuprofen, hydrocodone/homatropine, and hydrocodone/chlorpheniramine.

2.3. Outcomes

The outcome measured was average days' supply of the six opioid drug groups. Medians and ranges also were calculated. For a sensitivity analysis, we limited the Medicaid sample to states that continuously contributed data to Medicaid MarketScan. The results of the sensitivity analyses, medians, and ranges are reported in the Appendix.

2.4. Statistical analysis

Two-sample *t*-tests were used to compare average days' supply in the final year for which data are available (i.e., 2014 or 2015 depending on




Fig. 1. Average days' supply for opioid medications in adults with Medicaid, 2005–2014. Source: Truven Health MarketScan Multi-State Medicaid data, 2005–2014. Notes: The Medicaid states included in MarketScan varied by year. The average sample size for each drug-year is presented in parentheses: hydrocodone (1,090,000), hydromorphone (31,000), morphine (90,000), oxycodone (865,000), oxymorphone (10,500), and tapentadol (6200). The difference in average days' supply in 2014 compared to the baseline year (2005) is statistically significant ($p < 0.05$) for all drugs.

insurance) to 2005. All analyses were conducted using SAS 9.4 (Cary, NC, USA).

3. Results

3.1. Medicaid

For Medicaid (Fig. 1), we found an increase in days' supply for all drugs except morphine. Over the study period, the largest percentage increase was for oxycodone days supplied, which increased 37%. The average days supplied increased for hydrocodone, hydromorphone, oxymorphone, and tapentadol by 29%, 16%, 12%, and 24%, respectively. However, average days' supply of morphine decreased by 3%. In terms of the absolute change in days' supply, oxycodone, hydrocodone, hydromorphone, oxymorphone, and tapentadol increased by 4.5, 4, 2.5, 3, and 5 days, respectively. The average days' supply of morphine decreased by one day. After 2013, when the opioid epidemic had begun to be widely publicized, there was no decline in the median days supplied for any of the opioids (Appendix, Fig. A1).

3.2. Commercial

The average opioid days' supply for patients covered by commercial insurance (Fig. 2) exhibited a similar but steeper increase. Over the study period, the largest percentage increase was for oxycodone, which increased by 56%. Hydrocodone, hydromorphone, oxymorphone, and tapentadol increased by 33%, 40%, 32%, and 40%, respectively. Average days' supply of morphine decreased by 4%. In terms of absolute change in days' supply, oxycodone, hydrocodone, hydromorphone, oxymorphone, and tapentadol increased by 6, 3.5, 5, 7, and 7.5 days, respectively. The average days' supply of morphine decreased by one day. Between 2013 and 2015, when the opioid epidemic had begun to be widely publicized, there was no decline in the median days supplied of any of the opioids, and one of the most widely prescribed (hydrocodone) increased (Appendix, Fig. A2).

The sensitivity analysis was conducted using the same states over time (Appendix, Fig. A3). Results revealed similar trends, with days' supply increasing for all drugs except oxymorphone. The largest increase was for oxycodone, which increased 66% (an increase of 7 days) over the study period. Average days' supply of hydrocodone, hydromorphone, morphine, and tapentadol increased by 52%, 28%, 1%, and 30%, respectively. The average days' supply of oxymorphone did not change. The absolute change in days' supply of hydrocodone,

Download English Version:

<https://daneshyari.com/en/article/5037654>

Download Persian Version:

<https://daneshyari.com/article/5037654>

[Daneshyari.com](https://daneshyari.com)