



Short communication

Cigarette smoking quit rates among adults with and without alcohol use disorders and heavy alcohol use, 2002–2015: A representative sample of the United States population



Andrea H. Weinberger^{a,b}, Misato Gbedemah^c, Renee D. Goodwin^{c,d,e,*}

^a Ferkauf Graduate School of Psychology, Yeshiva University, Bronx, NY 10461 USA

^b Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY, 10461, USA

^c Department of Epidemiology and Biostatistics, Graduate School of Public Health and Health Policy, The City University of New York, New York, NY, USA

^d Institute for Implementation Science and Population Health, The City University of New York, New York, NY, USA

^e Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA

ARTICLE INFO

Keywords:

Smoking

Cessation

Alcohol use disorders

Heavy alcohol use

Epidemiology

ABSTRACT

Background: While the overall smoking quit rate has increased over time, it is not known whether the quit rate has also increased among persons with alcohol use disorders (AUDs) or heavy alcohol use (HAU). The current study examined quit rates among adults with and without AUDs and HAU over a 12-year period in a representative sample of US adults.

Methods: Data were drawn from the National Household Survey on Drug Use, an annual cross-sectional study of US persons. Quit rate (i.e., the rate of former smokers to ever smokers) was calculated annually from 2002 to 2014 (for HAU) and 2015 (for AUD). Time trends in quit rates by AUD/HAU status were tested using linear regression.

Results: The prevalence of past-month cigarette smoking was much higher for persons with, compared to without, AUDs (38% vs. 18%) and HAU (49% vs. 19%). In the most recent data year, the quit rate for persons with AUDs was approximately half that of persons without AUDs (26% versus 49%) and for persons with HAU was less than half that of persons without HAU (22% versus 48%). Over time, the smoking quit rate increased for persons with and without AUDs/HAU and the rate of increase was greater for persons with AUDs/HAU. Yet, quit rates for persons with AUDs and HAU remained much lower than persons without AUDs and HAU.

Conclusions: It may be beneficial for public health and clinical efforts to incorporate screenings and treatment for tobacco use into programs for adults with AUDs and HAU.

1. Introduction

Tobacco use is responsible for enormous societal burden in the United States (US) and is a leading cause of mortality around the world (USDHHS, 2014; WHO, 2012). Cigarette smoking appears substantially more common among adults with alcohol use disorders (AUDs) and heavy alcohol use (HAU) and both AUD and HAU appear to be associated with lower rates of smoking cessation (Falk et al., 2006; Jiang et al., 2015; Weinberger et al., 2016). For example, cross-sectional data from US adults suggest lower quit rates among adults with AUDs compared to those without psychiatric or substance use disorders (Smith et al., 2014) and among heavy drinkers compared to lighter drinkers or abstainers (Dawson, 2000). The majority of studies on quitting among adults with AUDs and HAU have estimated quit rate at a

single time period cross-sectionally. It therefore remains unclear whether quit rates for adults with AUDs and HAU have increased over time and in a manner that is comparable to quit rates among those without AUDs/HAU. Understanding the degree to which all groups—especially those vulnerable to continued cigarette use—are benefitting equally from tobacco control efforts can inform next steps in terms of the potential need for additional or more targeted public health and clinical efforts for persons with AUDs and HAU.

The purpose of the current study was to examine the quit rate among adults with and without AUDs and HAU over a 12-year period in a representative sample of US adults. The first aim was to compare the quit rate among adults with and without AUDs and HAU in the most recent data year. The second aim was to examine trends in quit rates from 2002 to 2014/2015 among adults with and without AUDs and

* Corresponding author at: Department of Epidemiology and Biostatistics, Graduate School of Public Health and Health Policy, The City University of New York, New York, NY, USA.
E-mail addresses: andrea.weinberger@einstein.yu.edu (A.H. Weinberger), misato.gbedemah@qc.cuny.edu (M. Gbedemah), renee.goodwin@sph.cuny.edu (R.D. Goodwin).

HAU.

2. Material and methods

2.1. Study population

Study data were drawn from The National Survey on Drug Use and Health (NSDUH) public data portal (<http://www.icpsr.umich.edu/> <http://www.icpsr.umich.edu/>) for the years 2002 to 2015. The NSDUH provides annual cross-sectional national data on the use of tobacco, other substance use, and mental health in the US and is described in depth elsewhere (SAMHSA, 2014). A multistage area probability sample for each of the 50 states and the District of Columbia was conducted to represent the male and female civilian non-institutionalized US population aged 12 and older. The datasets from each year were concatenated, adding a variable for the survey year. Because the assessment of HAU changed when data were collected in 2015, data are presented for HAU from 2002 to 2014 and for AUDs from 2002 to 2015. For this study, the analyses were restricted to participants who were 18 years old or older and who met criteria for lifetime smoking (unweighted $n = 222,967$) from 2002 to 2015.

2.2. Measures

2.2.1. Demographics

Demographic variables were categorized as follows: age (18 to 20 years old as reference, 21 to 25 years old, 26 to 34 years old, 35 to 49 years old, 50 to 64 years old, 65 years or older), gender (male as reference, female), total annual family income (< \$20,000 as reference, \$20,000 to \$49,999, \$50,000 to \$74,999, \$75,000 or more), education (less than high school as reference, high school graduate, some college, college graduate), race/ethnicity (non-Hispanic White as reference, non-Hispanic Black, non-Hispanic Native American/Alaskan Native, non-Hispanic Native Hawaiian/Other Pacific Islander, non-Hispanic Asian, non-Hispanic more than one race, Hispanic) and marital status (married as reference, widowed, separated or divorced, never married).

2.2.2. Alcohol use variables

Past-year AUDs (alcohol dependence or alcohol abuse) were assessed in each annual survey by computer-assisted interviewing instrumentation based on DSM-IV criteria (APA, 1994). Alcohol abuse was defined as meeting one or more of four alcohol abuse criteria in the past 12 months and not meeting criteria for alcohol dependence in the past 12 months. Alcohol dependence was defined as meeting three or more of seven alcohol dependence criteria. Respondents who met criteria for past-year alcohol abuse or dependence were classified as having a past-year AUD. Detailed alcohol abuse and dependence criteria were described in NSDUH 2015 Codebook (Appendix D Recoded Substance Dependence and Abuse Variable Documentation section; SAMHSA, 2015). Past-month HAU was defined for all participants (i.e., men and women) as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days. AUD and HAU were not mutually exclusive categories.

2.2.3. Smoking status and quit rate

Respondents were classified as ever smokers if they reported having smoked 100 or more lifetime cigarettes. Respondents were classified as former smokers if they reported having smoked 100 or more lifetime cigarettes and no cigarette smoking in the past year. The Quit Rate was calculated as the rate of former to ever smokers and is considered a measure of total cessation in a population.

2.3. Statistical analyses

All analyses were performed incorporating the NSDUH sampling weights and controlling for the complex clustered sampling using SAS-

callable SUDAAN Version 11.0.1 (RTI International, Research Triangle Park, NC; <http://www.rti.org/sudaan/http://www.rti.org/sudaan/>). First, demographic characteristics were examined by AUD status in the most recent data year. Chi-square tests were then performed to assess the statistical significance of associations between demographics and AUD status. Second, the weighted number of former and ever smokers by past-year AUD status (AUD, no AUD) were calculated for each year from 2002 to 2015. Quit rates were then calculated as the ratio of former to ever smokers for those with and without AUDs. Time trends in the quit rates by AUD status were tested using linear regression with continuous year as the predictor for the linear time trend. These analyses were conducted twice: first with no covariates (unadjusted) and then adjusting for age, gender, income, education, race/ethnicity and marital status. To examine whether time trends in quit rate differed by AUD status, linear regressions included the 2-way interaction of survey year X AUD status (past-year AUD versus no past-year AUD). All analyses described above were repeated with HAU in place of AUD.

3. Results

3.1. Analytic sample

The prevalence of past month cigarette smoking among persons with AUDs was approximately twice that among persons without AUDs in 2015 (38% vs. 18%) and more than twice as common among persons with HAU—with almost half reporting current smoking—compared with those without HAU in 2014 (49% vs. 19%; see Supplemental Table 1¹ for demographics and smoking status by AUD/HAU).

3.2. Quit rate by AUD status

For each year of the study, the quit rate was lower for adults with AUDs compared to adults without AUDs (see Fig. 1). In 2002 and 2015, the quit rate among those with AUDs was approximately half that of those without AUDs (2002: 20% versus 47%; 2015: 26% versus 49%). Controlling for demographics, there was a significant increase in the quit rate from 2002 to 2015 for both those with AUDs (adjusted beta = 0.00601, 95% CI = 0.00599-0.00603, $p < 0.001$) and without AUDs (adjusted beta = 0.00292, 95% CI = 0.00291-0.00293, $p < 0.001$). There was a significant interaction such that rate of increase was significantly more rapid among those with AUDs relative to those without AUDs ($F = 18366$, $df = 1$, $p < 0.001$).

3.3. Quit rate by HAU status

For each year of the study, the quit rate was lower among adults with HAU compared to those without HAU (see Fig. 2). In 2002 and 2014, the quit rate among adults with HAU was less than half of that among adults without HAU (2002: 20% versus 46%; 2014: 22% versus 48%). Controlling for demographics, there was a significant increase in the quit rate from 2002 to 2014 for both adults with HAU (adjusted beta = 0.00484, 95% CI = 0.00482-0.00485, $p < 0.001$) and without HAU (adjusted beta = 0.00263, 95% CI = 0.00262-0.00263, $p < 0.001$). There was a significant interaction such that rate of increase in quit rate was more rapid among those with HAU compared to those without HAU ($F = 9784$, $df = 1$, $p < 0.001$).

4. Discussion

This study was the first to use representative data from the US population to examine quit rates over time among adults with and without AUDs/HAU. Between 2002 and 2014/2015, the smoking quit rate increased for adults with and without AUDs and HAU and the rate of increase in quitting over time was more rapid among persons with AUDs/HAU relative to persons without AUDs/HAU. Yet, quit rates for adults with AUDs and HAU were consistently half those of adults

Download English Version:

<https://daneshyari.com/en/article/5119887>

Download Persian Version:

<https://daneshyari.com/article/5119887>

[Daneshyari.com](https://daneshyari.com)