



Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: [www.elsevier.com/locate/tetlet](http://www.elsevier.com/locate/tetlet)

## Synthesis, structure and reactivity of a macrocyclic imine: aza-[13]-macrodilides

Jun Ma, Raghu Vannam, Daniel W. Terwilliger, Mark W. Peczuł\*

Department of Chemistry, University of Connecticut, 55 N. Eagleville Road, U3060, Storrs, CT 06269, USA

### ARTICLE INFO

#### Article history:

Received 30 April 2014

Revised 15 May 2014

Accepted 16 May 2014

Available online xxx

#### Keywords:

Macrocyclic

Imine

Synthesis

Structure

### ABSTRACT

The in situ synthesis and subsequent reactions of macrocyclic imine **2** are reported. The imine was trapped with cyanotrimethylsilane to give  $\alpha$ -amino nitrile aza-[13]-macrodilides in a 1:1 ratio of diastereomers. A crystal structure of the *syn*  $\alpha$ -cyano nitrile diastereomer, **7a**, provided insights into the lack of selectivity in reactions of **2** relative to macrocyclic alkene **1**. Reactions to functionalize the *syn* diastereomer **7a** are also reported.

© 2014 Elsevier Ltd. All rights reserved.

Macrocycles occupy an important niche among organic compounds because they share features with both small molecules and larger ones.<sup>1</sup> The number of atoms in the ring, the presence and positioning of planar units and the stereogenic centers within it all contribute to the overall shape or topology of the macrocycle.<sup>2,3</sup> We have been focusing recently on how these factors interact in the context of some [13]-macrodilides such as **1** (Fig. 1). Planar units such as esters, amides, and alkenes, for example, help to rigidify macrocycles and thereby reduce the number of low-energy conformations available to them. In the context of **1**, the ester and alkene units simplified its structure to three planar units and one 'hinge' atom.<sup>2b,c</sup> Further, the planar units must orient themselves in a non-coplanar fashion that creates a fold to the ring much like the twist of *E*-cyclooctene. Macrocyclic **1** exhibits planar chirality when homochiral starting materials are used; we have shown that the absolute configuration of key stereogenic centers govern the planar chirality<sup>4</sup> of [13]-macrodilides.<sup>2</sup>

Here we evaluate the substitution of an imine for the alkene unit in **1** (e.g., **2** in Fig. 1).<sup>2,5</sup> Specifically, we thought that the imine functionality, from the  $\alpha$ -carbon through the imine carbon and nitrogen to the carbon attached to the nitrogen, would act as a four-atom planar unit (blue atoms in Fig. 1). The investigation aimed to illustrate the imine as another planar unit that would rigidify the [13]-macrodilide structure and also switch the reactivity from that of a nucleophile to an electrophile. We were also motivated by the beneficial new properties associated with





Figure 1. [13]-Macrodilide **1** and aza-[13]-macrodilide analog **2**.

the incorporation of a nitrogen atom into a macrocycle.<sup>6–9</sup> Analysis of products, for example, aza-[13]-macrolide **7** (Scheme 1) that arise from additions to macrocyclic imine **2** would also provide information about the structure of **2**.<sup>10</sup> The structure and reactivity of imine **2** were of interest because we considered it an aza-analog of **1**. We report the in situ synthesis of macrocyclic imine **2** and its subsequent functionalization to novel aza-[13]-macrodilides.

The synthesis of imine **2** followed a strategy similar to the synthesis of **1**.<sup>2</sup> Sequential, chemoselective acylations of racemic 1,3-butanediol—first with pentenoic acid then with *N*-Boc- $\beta$ -alanine—gave diester **3** in 80% over two steps (Scheme 1). Ozonolysis of the terminal alkene of **3** then provided aldehyde **4** (86%). It should be noted that the choice of aldehyde and Boc protected amine groups for the precursor to the macrocycle was made after earlier attempts with alternative protecting group schemes (i.e., Cbz protected amine with the aldehyde) were unsuccessful. With compound **4** in hand, the following steps were done sequentially in one pot in the successful sequence: (i) TFA deprotection of the

\* Corresponding author. Tel.: +1 860 486 1605; fax: +1 860 486 2981.

E-mail address: [mark.peczuł@uconn.edu](mailto:mark.peczuł@uconn.edu) (M.W. Peczuł).



**Scheme 1.** Synthesis aza-[13]-macrodilolides **5** and **7** via macrocyclic imine **2**.

Boc carbamate; (ii) 20-fold dilution and dehydration using  $\text{MgSO}_4$ <sup>11</sup> and (iii) neutralization by addition of triethylamine. At the outset we had hoped to characterize and potentially isolate imine **2** because there is literature precedence for the formation of other macrocyclic imines.<sup>8,12</sup> Attempts at isolation of **2** were unsuccessful, however. We ultimately resorted to trapping the imine directly with nucleophiles to confirm that it had in fact formed. Sodium cyanoborohydride reduction in methanol<sup>14</sup> did not result in the expected aza-[13]-macrodilolide product **5**; instead, intramolecular amidation and transesterification led to a ring contracted  $\gamma$ -lactam **6** (Scheme 1, inset). The structure of **6** strongly suggested that **5**, and consequently imine **2** itself, were intermediates in the ring contraction (See Supporting information). Hydride reduction using  $\text{NaBH}(\text{OAc})_3$  did provide aza-[13]-macrodilolide **5**, albeit in only 6% yield as a mixture with other unidentified products.

Addition of cyanotrimethylsilane (TMSCN) after in situ imine formation, on the other hand, yielded cyano aza-[13]-macrodilolides **7** as a ~1:1 mixture of diastereomers, **7a** and **7b** (Scheme 1 and Fig. 2) in a combined 51% yield. It should be noted that each diastereomer of product **7** is itself racemic because the 1,3-butane diol used to start the synthesis was racemic. The yield for the formation of **7** was calculated based on *N*-Boc aldehyde **3** as starting material and formally included four steps.

The *syn* diastereomer, **7a**, was a crystalline solid. Data from X-ray crystallography revealed the structure depicted in Figure 2. Compound **7a** has features that are reminiscent of the X-ray structure collected for **1** and provides clues about macrocyclic imine **2**. First, each ester creates a four atom planar unit that reduces the number of freely rotating bonds in the backbone of the macrocycle. Second, the stereogenic centers at C2 and C9 both put the groups attached to them (methyl at C2 and cyano at C9) in pseudo equatorial positions. Finally, the cyano moiety at C9 is exocyclic to the macrocycle; that is, it points outside the ring. Epoxidations on chiral [13]-macrodilolides akin to **1** were highly diastereoselective.<sup>2a</sup> In those examples, the  $\pi$ -system of the alkene was shown to react exclusively through its exterior face. The position of the cyano group on the periphery of the macrocycle in **7a** suggests that the  $\alpha$ -cyano amine arose by nucleophilic attack on a planar imine also from the exterior face.

The structural data from **7a** suggest that imine **2** largely resembles [13]-macrodilolide **1** (Fig. 2). The difference is that the imine of

**2** acts as an electrophile rather than as a  $\pi$  nucleophile as does the alkene in **1**. Conversion of **2** to a ~1:1 mixture of diastereomers **7** shows that there is an absence of diastereofacial selectivity for reactions of this  $\pi$ -system, however. This result is in contrast to other compounds where macrocyclic diastereocontrol was operative.<sup>2a,13</sup> The dichotomy indicates a further difference in the nature of macrocycles **1** and **2**. The lack of stereoselectivity for TMSCN addition to **2** could be explained by at least four different models: (i) Imine **2** could take up a rigid geometry that is different than the configuration of **1** in Fig. 2 that would allow attack of nucleophiles from either diastereoface. This is unlikely based on the structural requirements of **1** and **2**. Specifically, we have only observed the formation of *E* alkenes (e.g., **1**) which suggests that the *E* configuration of the  $\pi$ -system is best accommodated based on the architecture of the 13-membered ring macrocycle. (ii) The  $\alpha$ -amino nitrile products could equilibrate after they are formed and the reaction is under thermodynamic control. Epimerization of  $\alpha$ -amino nitriles is known but under conditions that are more robust (either acidic or basic) than those used in the formation of **7a** and **7b**.<sup>14,15</sup> As reported here, imine **2** arose by TFA deprotection of a Boc group and then the acid was neutralized by one equivalent of TEA prior to addition of TMSCN. Moreover, exposure of **7a** to di-isopropylethylamine for 12 h at 70 °C showed no epimerization. (iii) The macrocyclic imines could interconvert via an amination intermediate. Such a process could interconvert an *E* imine to *Z* or it could interconvert alternate *E* imines (Fig. 3A). Although we cannot rule it out, the *Z* imine is unlikely based on the ring size and the rigidifying units. The interconversion of *E* imines, however, is on equal footing as the proceeding imine rotational model. (iv) The imine unit is freely rotating within the [13]-macrodilolide. We argue that the lack of selectivity is due to facile rotation of the imine unit in **2** in contrast to the stationary alkene unit of **1** (Fig. 3B and C). In this scenario, rapid rotation of the imine exposes each diastereoface outward toward solution and therefore makes each available for attack by nucleophiles. The nearly 1:1 ratio of diastereomeric products is consistent with the model. Similar bond lengths for both the alkene and the imine imply that the removal of the hydrogen is sufficient to allow facile rotation of the planar unit.<sup>16</sup> Although somewhat speculative, the rotational model is consistent with the structures and reactivity we have observed for this group of macrocycles.

Our next objective was to modify the newly prepared aza-[13]-macrodilolides using **7a** as the example compound. Alkylation of

Download English Version:

<https://daneshyari.com/en/article/5269298>

Download Persian Version:

<https://daneshyari.com/article/5269298>

[Daneshyari.com](https://daneshyari.com)