Triglyceride increase in the core of high-density lipoproteins augments apolipoprotein dissociation from the surface: Potential implications for treatment of apolipoprotein deposition diseases

Shobini Jayaraman, Jose Luis Sánchez-Quesada, Olga Gursky

PII: S0925-4439(16)30254-X
DOI: doi: 10.1016/j.bbadis.2016.10.010
Reference: BBADIS 64576

To appear in: *BBA - Molecular Basis of Disease*

Received date: 18 August 2016
Revised date: 13 October 2016
Accepted date: 16 October 2016

Please cite this article as: Shobini Jayaraman, Jose Luis Sánchez-Quesada, Olga Gursky, Triglyceride increase in the core of high-density lipoproteins augments apolipoprotein dissociation from the surface: Potential implications for treatment of apolipoprotein deposition diseases, *BBA - Molecular Basis of Disease* (2016), doi: 10.1016/j.bbadis.2016.10.010

Triglyceride increase in the core of high-density lipoproteins augments apolipoprotein dissociation from the surface: Potential implications for treatment of apolipoprotein deposition diseases

Running title: Triglyceride-lowering therapies may help alleviate apoA-I amyloidosis

Authors: Shobini Jayaraman[1], Jose Luis Sánchez-Quesada[2], Olga Gursky[1]

Affiliation:

[1] Department of Physiology & Biophysics, Boston University School of Medicine, Boston USA

[2] Cardiovascular Biochemistry Group, Biomedical Research Institute IIB-Sant Pau, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain.

Corresponding author: Dr. Olga Gursky

Department of Physiology & Biophysics, Boston University School of Medicine, W321, 700 Albany St. Boston MA 02118. E-mail: gursky@bu.edu Tel : 617-638-7894 FAX : 617-638-4041