

Oxidized cholesteryl esters and inflammation[☆]

Soo-Ho Choi^a, Dmitri Sviridov^b, Yury I. Miller^{a,*}

^a Department of Medicine, University of California, San Diego, La Jolla, CA 92093, USA

^b Baker IDI Heart and Diabetes Institute, Melbourne, VIC 3004, Australia

ARTICLE INFO

Article history:

Received 29 March 2016

Received in revised form 8 June 2016

Accepted 24 June 2016

Available online xxxxx

Keywords:

Cholesterol

Cholesteryl ester

Oxidation

TLR4

MD-2

Atherosclerosis

ABSTRACT

The oxidation hypothesis of atherosclerosis proposes that oxidized LDL is a major causative factor in the development of atherosclerosis. Although this hypothesis has received strong mechanistic support and many animal studies demonstrated profound atheroprotective effects of antioxidants, which reduce LDL oxidation, the results of human clinical trials with antioxidants were mainly negative, except in selected groups of patients with clearly increased systemic oxidative stress. We propose that even if reducing lipoprotein oxidation in humans might be difficult to achieve, deeper understanding of mechanisms by which oxidized LDL promotes atherosclerosis and targeting these specific mechanisms will offer novel approaches to treatment of cardiovascular disease. In this review article, we focus on oxidized cholesteryl esters (OxCE), which are a major component of minimally and extensively oxidized LDL and of human atherosclerotic lesions. OxCE and OxCE-protein covalent adducts induce profound biological effects. Among these effects, OxCE activate macrophages via toll-like receptor-4 (TLR4) and spleen tyrosine kinase and induce macropinocytosis resulting in lipid accumulation, generation of reactive oxygen species and secretion of inflammatory cytokines. Specific inhibition of OxCE-induced TLR4 activation, as well as blocking other inflammatory effects of OxCE, may offer novel treatments of atherosclerosis and cardiovascular disease. This article is part of a Special Issue entitled: Lipid modification and lipid peroxidation products in innate immunity and inflammation edited by Christoph J. Binder.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Low-density lipoprotein (LDL), which gradually accumulates in the vascular wall in humans and in hypercholesterolemic animal models, is highly susceptible to oxidative damage due to its complex lipid-protein composition and a large number of polyunsaturated fatty acyl (PUFA) chains. Strong evidence for the presence of oxidized LDL (OxLDL) in human atherosclerotic lesions provided a scientific rationale for the “oxidation hypothesis of atherosclerosis,” stating that OxLDL is a major causative factor in the development of atherosclerosis [1,2]. This hypothesis generated widespread enthusiasm for antioxidants as a therapeutic approach to delay the development of atherosclerosis and reduce the incidence of cardiovascular events. Indeed, many animal studies demonstrated profound atheroprotective effects of various

antioxidants [2]. However, the results of human clinical trials with antioxidants were mainly negative, except in selected groups of patients with clearly increased systemic oxidative stress, such as patients on hemodialysis or diabetics with haptoglobin 2-2 genotypes associated with higher hemoglobin-mediated oxidative stress [3–5].

If reducing lipoprotein oxidation in the vessel wall is difficult to achieve in the majority of human populations, can deleterious effects of OxLDL be contained? In our opinion, deeper understanding of the biological effects of OxLDL and its components may identify specific targets for future therapy, offering better focused and, potentially, personalized treatment of cardiovascular disease (CVD).

We and others focus on inflammatory responses to OxLDL because inflammation in atherosclerotic lesions is believed to be a major contributor to plaque vulnerability to rupture, which causes acute cardiovascular and cerebrovascular events. The concept of host-derived, damage-associated molecular patterns (DAMPs) helps understand why OxLDL becomes inflammatory [6]. It appears that DAMPs share structural motifs with microbial pathogen-associated molecular patterns and activate pattern-recognition, innate immune receptors. Ensuing inflammatory responses can be harmful if unconstrained and prolonged, such as in atherosclerosis, a life-long, chronic inflammation of large arteries leading to CVD. OxLDL epitopes detected in human vulnerable plaques by specific antibodies become increasingly more prominent as lesions progress and rupture [7]. These epitopes are particularly

Abbreviations: 12/15-LO, 12/15-lipoxygenase; 9-ON, 9-oxononanoate; CE, cholesteryl esters; CVD, cardiovascular disease; DAMPs, damage-associated molecular patterns; HPODE, hydroperoxyoctadecadienoate; LDL, low-density lipoprotein; MD-2, myeloid differentiation-2; mLDL, minimally oxidized LDL; OxCE, oxidized CE; OxLDL, oxidized LDL; OxPL, oxidized phospholipid; PUFA, polyunsaturated fatty acid or acyl; TLR4, toll-like receptor-4.

[☆] This article is part of a Special Issue entitled: Lipid modification and lipid peroxidation products in innate immunity and inflammation edited by Christoph J. Binder.

* Corresponding author.

E-mail address: yumiller@ucsd.edu (Y.I. Miller).

<http://dx.doi.org/10.1016/j.bbalip.2016.06.020>

1388–1981/© 2016 Elsevier B.V. All rights reserved.

Please cite this article as: S.-H. Choi, et al., Oxidized cholesteryl esters and inflammation, *Biochim. Biophys. Acta* (2016), <http://dx.doi.org/10.1016/j.bbalip.2016.06.020>

prominent in advanced coronary and carotid lesions in macrophage-rich areas, lipid pools, the necrotic core and in ruptured plaques. The presence of OxLDL in clinically relevant human lesions provides a strong rationale to use OxLDL epitopes as biomarkers in plasma and as a molecular imaging target in atherosclerotic plaques for clinical applications.

Among OxLDL components, oxidized phospholipids (OxPL) and oxysterols were the major focus of much research in the field, resulting in identification of molecular structures, receptors and signaling pathways leading to inflammatory responses in vascular cells [8–13]. Less attention was given to oxidized cholesteryl esters (OxCE). In this article, we review studies of OxCE, with the goal to describe remarkable features of CE oxidation and OxCE-activated pathways and responses and to demonstrate the importance of containment of OxCE-induced inflammation in atherosclerosis.

2. Cholesteryl ester oxidation and its prevalence in atherosclerotic plaque

Cholesterol esterified with a fatty acyl chain is the form in which cholesterol is transported within LDL from liver to the periphery and stored in lipid droplets inside the cell. Even though CE reside in the hydrophobic core of LDL, CE with PUFA chains are readily oxidized, via enzymatic and free radical mechanisms. Inducers and mechanisms of LDL oxidation have been extensively investigated and reviewed elsewhere [14]. These include reactions catalyzed by 12/15-lipoxygenase (12/15-LO), myeloperoxidase, nitric oxide synthases and NADPH oxidases, as well as those mediated by transition metals, heme and hemoglobin [15]. Several studies suggest that CE is a preferential substrate for 12/15-LO [16,17]. In an *in vitro* reaction of LDL oxidation by rabbit 15-LO, a close homolog of human 15-LO and mouse 12/15-LO, even when the LDL particle was loaded with free linoleic acid, cholesteryl linoleate constituted the major 15-LO substrate [16]. Remarkably, intracellular OxCE hydrolysis and subsequent incorporation of an oxidized fatty acyl chain into PL often is the source of OxPL in the cell [17]. The authors used wild type and 12/15-LO-deficient murine peritoneal macrophages and radioisotope labeled cholesteryl linoleate and cholesteryl arachidonate to demonstrate that both intracellular CE and the CE in LDL are effectively oxidized by macrophage 12/15LO and that the oxidized fatty acyls originated in OxCE can then be found as part of OxPL molecules.

Specific 12/15-LO products of CE oxidation can be further oxidized in free radical-mediated reactions, forming numerous and complex isoprostane OxCE products, with up to 6 oxygen atoms inserted in the molecule of cholesteryl arachidonate [18,19]. Only the PUFA chain of the CE undergoes oxidation, whereas the sterol remains unmodified. Prominent among these polyoxygenated CE products were molecules with a bicyclic endoperoxide group [18,19], such as cholesteryl (9,11)-epidioxy-15-hydroperoxy-(5Z,13E)-prostadienoate shown in Fig. 1 (abbreviated as BEP-CE for the presence of bicyclic endoperoxide and hydroperoxide groups).

Importantly, specific OxCE molecules first identified in test-tube oxidation reactions were also found in cellular systems, in vascular lesions of experimental animals and in plasma and atherosclerotic plaques isolated from human CVD patients. Cholesteryl hydroperoxyoctadecadienoate (HPODE) was a major oxidized lipid in LDL incubated with activated human monocytes [20]. Cholesteryl HPODE can further decompose to form cholesteryl 9-oxononanoate (9-ON), a core aldehyde, which was found in human atherosclerotic lesions and can react with ϵ -aminogroup of lysines and form covalent adducts with proteins [21], particularly in small, dense LDL under oxidative conditions [22].

Earlier studies estimated that 2% of total CE and 30% of cholesteryl linoleate are oxidized in human plaques [23,24]. A recent study employing advanced mass spectrometry techniques identified many OxCE species in human atherosclerotic lesions and quantified that, on average, 23% of cholesteryl linoleate, 16% of cholesteryl arachidonate and 12% of cholesteryl docosahexaenoate were oxidized [25]. In another mass spectrometry study, OxCE comprised from 11% to as much as




Fig. 1. Oxidized cholesteryl esters. Cholesteryl arachidonate has 4 unsaturated bonds in the fatty acyl chain, which makes it susceptible to oxidation in general and a preferable substrate for 12/15-lipoxygenase (12/15-LO). Initial 12/15-LO enzymatic oxygenation of cholesteryl arachidonate is often followed by free radical oxidation, resulting in a variety of polyoxygenated CE molecules. The example shown is a product with bicyclic endoperoxide and hydroperoxide groups (BEP-CE, or cholesteryl (9,11)-epidioxy-15-hydroperoxy-(5Z,13E)-prostadienoate), which has specific biological activity as illustrated in Fig. 2.

92% of the CE-PUFA pool in human plaques [26]. Among cholesteryl arachidonate oxidation products, BEP-CE was detected in human plasma from CVD patients and in human atherosclerotic lesions [27], as well as in experimental studies with hypercholesterolemic mice and zebrafish [6,28]. Admittedly, it is difficult to establish if OxCE in human atherosclerotic plaque are mainly associated with lipoproteins retained in the extracellular matrix or reside intracellularly. Human atherosclerotic lesions contain OxCE not only in a free lipid form, but also as covalent adducts to proteins, including apoB-100, as detected with a monoclonal antibody raised against proteins modified with cholesteryl 9-ON [29].

3. Biological effects of OxCE

It was noticed that OxLDL inactivates PDGF, and cholesteryl HPODE, likely forming covalent adducts with PDGF, was identified as a lipid in OxLDL responsible for this effect [30]. Hydrogen peroxide and short fatty acid hydroperoxides did not inactivate PDGF. Cholesteryl HPODE also inactivated bFGF and TGF- β but not EGF [30]. Both cholesteryl HPODE and cholesteryl 9-ON activated PKC and ERK1/2 in endothelial cells and resulted in expression of connecting segment-1 and enhanced adhesion of monocytes to endothelial cells [31]. Cholesteryl 9-ON induced expression of both TGF- β and TGF- β receptor type I in human U937 promonocytic cells, the effect mediated by ERK1/2 and potentially involved in sustaining vascular remodeling in atherosclerosis [32]. Cholesteryl HPODE was the active component of moderately oxidized LDL that activated PPAR α -dependent expression of CD36 in human monocyte-derived macrophages, whereas PL-esterified HPODE and 7-ketocholesterol did not produce this effect [33].

There is a mechanism to detoxify OxCE in circulation, which involves CETP-mediated exchange of OxCE between LDL and HDL [34] and selective uptake of HDL-associated OxCE by liver via an SR-BI-mediated mechanism [35]. The selective uptake of OxCE was 2–3 fold higher compared with selective uptake of non-oxidized CE, and high-cholesterol feeding of rats significantly decreased OxCE uptake by liver parenchymal cells but dramatically increased OxCE uptake by Kupffer cells [35].

4. OxCE activation of TLR4/MD-2

Work from our laboratory demonstrated that minimally oxidized LDL (mmLDL), produced by LDL oxidation with 15-LO-expressing cells, induced profound cytoskeleton changes in macrophages, including actin polymerization, cell spreading, membrane ruffling and macropinocytosis

Download English Version:

<https://daneshyari.com/en/article/5508502>

Download Persian Version:

<https://daneshyari.com/article/5508502>

[Daneshyari.com](https://daneshyari.com)