



Influence of two ovulation-inducing agents on the pituitary response and follicle blood flow in mares



Y.L. Boakari ^{a, b}, J.C. Ferreira ^{c, d, *}, H.S. Canesin ^a, D.L. Thompson Jr. ^d, F.S. Lima ^c,
J.C.F. Pantoja ^a, C. Meira ^a

^a Department of Animal Reproduction and Veterinary Radiology, School of Veterinary Medicine and Animal Science, UNESP, Botucatu, São Paulo, Brazil

^b Gluck Equine Research Center, Department of Veterinary Science, University of Kentucky, Lexington, United States

^c Department of Veterinary Clinical Medicine, College of Veterinary Medicine, University of Illinois, Urbana, IL, USA

^d School of Animal Sciences, Louisiana State University Agricultural Center, Baton Rouge, LA, United States

ARTICLE INFO

Article history:

Received 6 February 2017

Received in revised form

24 May 2017

Accepted 30 May 2017

Available online 8 June 2017

Keywords:

LH

Deslorelin

hCG

Follicle vascularity

Doppler ultrasonography

ABSTRACT

The objective of the current study was to evaluate the effects of deslorelin and hCG, two ovulation-inducing therapies, on LH surge and follicle vascularity in mares. Thirty mares were either treated with 1.5 mg IM of deslorelin, 2,500 IU IV of hCG or 2 mL IM of NaCl 0.9% (GnRH, hCG and Saline groups, respectively). Power-flow Doppler examination and blood collection were performed every hour during the first 12 hours after treatment (H0) and every six hours between hours 12 (H12) and 30 (H30) after treatment. Moreover, endpoints were evaluated every hour through the last six hours before ovulation (OV-6 to OV-1). In GnRH group, plasma LH concentration progressively increased ($P < 0.001$) during the first 6 hours after treatment and remained high ($P > 0.1$) until OV-1. A significant increase in LH concentrations was first detected ($P < 0.05$) at 24 hours after treatment in hCG group, while no changes ($P > 0.1$) on LH levels were found during H0-H30 and between OV-6 and OV-1 in the Saline group. Independent of the treatment, significant variations on the percentage of the follicle wall with Doppler signals were not observed ($P > 0.1$) throughout the entire experiment. A weak correlation between the preovulatory follicle vascularity and the plasma LH concentration was found in GnRH, hCG and Saline groups ($r = +0.29, +0.29$ and -0.23 , respectively; $P < 0.0001$). These results described for the first time the immediate and continuous pituitary response to ovulation-inducing therapy with injectable deslorelin. Moreover, spontaneous and induced ovulations were not preceded by an increased follicle vascularity, which differs from previous reports in large animals.

Published by Elsevier Inc.

1. Introduction

Determining the most appropriate moment to perform natural or artificial mating is essential to maximize conception rates through the breeding season in mares [1]. However, the long physiologic duration of the equine estrus, when compared to other domesticated livestock [2–4], makes it difficult to accurately predict the time of ovulation [5]. Currently, the use of ovulation-inducing therapies [6] and the visualization of impending ovulation indicators by B-mode ultrasound examination [7] has been applied to ensure breeding at the appropriate time.

In mares, ovulation-inducing therapies with deslorelin or hCG

stimulate the final maturation and the rupture of the follicular wall within 48 h [1,8,9]. Deslorelin, a synthetic GnRH agonist [1,10], acts on the adenohypophysis inducing the pulsatile release of endogenous LH and the subsequent ova maturation [11]. Differently, hCG binds to follicular LH receptors due to its structural similarity to the equine gonadotropin molecule [12,13], prompting an analogous LH-activity [14].

Transrectal Doppler ultrasonography has been recently used to study the blood flow of the follicle wall in mares [15,16], cattle [17,18], ewes [19], canines [20], and women [21]. Changes in the ovarian vascular network during estrus allow the delivery of essential components for follicular maturation and steroidogenesis [22]. Therefore, Doppler ultrasonography has the potential to be an efficient real-time method for *in vivo* evaluation of the functional status of preovulatory follicles. However, the current literature remains unclear about the follicle hemodynamics during the

* Corresponding author. 1008 W Hazelwood Dr, Urbana, IL 61802, USA.

E-mail address: jair@illinois.edu (J.C. Ferreira).

impending ovulation phase in mares. Initial studies described an increased follicle vascularity through the preovulatory period [16,23] followed by an abrupt decrease of the Doppler signals in the follicle wall during the last four hours before ovulation [7]. In contrast, the occurrence of ovulation apparently was not affected by the absence of changes on the follicular vascularity in mares treated with hCG [24] or recombinant equine LH [15]. Moreover, the effect of synthetic GnRH agonist agents on the blood flow of dominant follicles has not been described in mares.

Studies with cattle and mares have suggested a correlation between the LH surge and the blood-flow of follicles after ovulation-inducing therapy [25–28]. Considering the distinct biologic activity of hCG and GnRH analogs on final follicular maturation, we have hypothesized that the hemodynamics of dominant follicles are dependent on the ovulation-inducing agent. Therefore, the primary purpose of the present study was to characterize the influence of treatment with hCG and deslorelin on the vascularity of preovulatory follicles in mares. The specific goals were: a) To describe the relationship between changes in the percentage of follicle wall with power-flow signals and the concentrations of LH during the impending ovulation phase, and b) to evaluate the effectiveness of Doppler technology for predicting spontaneous and induced ovulations in mares.

2. Material and methods

2.1. Animals and experimental groups

Thirty cycling mixed breed mares with 5–15 years of age, weighting 250–380 kg, were used for the present study. Mares were handled in accordance with the Guide for Care and Use of Agricultural Animals in Agricultural Research and Teaching (protocol number 139/2012). The study was conducted during a single mid-breeding season (from January to March) in the Reproduction Center of the Department of Animal Reproduction and Veterinary Radiology at the São Paulo State University, Brazil (Latitude 22° 53' 09" and Longitude 48° 26' 4"). All mares were maintained on grass hay, pelleted feed and trace mineralized salt with free access to water. The age of mares was estimated from dental characteristics [29]. Score for body condition was evaluated according to Henneke et al. (1983) and for all mares, it remained high (score \geq 7) during the experiment.

B-mode ultrasonography examination was done once a day to monitor follicular development. Mares showing a pre-ovulatory follicle \geq 35 mm of diameter associated with endometrial edema were assigned into three experimental groups. The first group (n = 10) received GnRH (1.5 mg of Deslorelin, IM); the second group (n = 10) received hCG (2500 IU of hCG, IV); and third group (n = 10) received a Saline (1.5 mL of NaCl 0.9%, IM) and served as a control group in our experiment. Only mares that ovulated between 30 and 48 h after treatment were used in the GnRH and hCG groups. All mares from the Saline group had spontaneous ovulation. Mares with double ovulations or anovulatory hemorrhagic follicles were not used in the study. The moment immediately before treatment was considered H0.

2.2. Data collection

Blood collection and Doppler ultrasonography examination were performed to characterize the changes in plasma LH concentration and vascularity of the preovulatory follicle. Data collection was performed every one hour during the first 12 h of the study and every six hours from hours 12 (H12) to 30 (H30) after

treatment. Additionally, endpoints were evaluated hourly from H30 until the detection of a corpus luteum (CL) in mares treated with GnRH or hCG. In the Saline group, mares were scanned with B-mode ultrasonography every 4 h from H30 to the visualization of impending ovulation signals within 12 h. After that, endpoints were evaluated hourly, and mares that ovulated in less than 10 h were included in the study. Impeding ovulation was predictable by the combination of a thick and echogenic granulosa, decreased turgidity, loss of spherical shape, detached granulosa segments and echoic spots in the antrum [30–33].

2.3. Doppler ultrasonography examination

Vascularity of the preovulatory follicles was estimated using a pulsed-wave color Doppler ultrasound instrument (SONOACE PICO, Medison Brazil Ltda) equipped with a linear endocavitary transducer of 5–9 MHz (LV5-9CDn, 60 mm). Doppler settings were maintained constant during the entire study. Power-flow mode function was used to display blood-flow signals from the vessels of the follicle wall. The entire follicle was scanned in a slow and continuous motion. Scans of 1 min were recorded in a portable computer equipped with a video capture device (Pinnacle Studio 9, Ottawa, ONT, Canada). The follicular vascularity of each film was posteriorly estimated by two independent operators blind to treatment or ovulation. A similar subjective methodology has been used for studying blood-flow of follicles in mares [16,34] and heifers [18]. The vascularity of preovulatory follicles was scored considering the percentage (0%–100%) of follicle wall circumference with Doppler signals during the film clip analysis, as described previously for large animals [7,18,24,34].

Supplementary video related to the Doppler ultrasonography examination of the follicle can be found at <http://dx.doi.org/10.1016/j.theriogenology.2017.05.032>

2.4. Plasma LH concentration

Blood samples were collected immediately before each Doppler ultrasonography examination by jugular venipuncture into heparinized tubes for the measurement of plasma LH concentration. Blood was centrifuged (1500 \times g/10 min) and plasma was stored at -20° C until assayed.

Hormonal assays were performed in the School of Animal Sciences at Louisiana State University Agricultural Center, Baton Rouge, USA. Plasma samples were assayed by radioimmunoassay, as previously validated for mares [35]. The double-antibody assay was based on anti-equine chorionic gonadotropin primary antiserum generated in a rabbit and radio-iodinated ovine LH tracer. The intra- and inter-coefficient of variation were 3.2% and 7.0%, respectively. The sensitivity of the assay was 0.4 ng/mL.

2.5. Statistical analysis

The first part of the data was normalized to the moment immediately before treatment (Hour 0) and ended at Hour 30. The last six hours before the CL visualization (OV-6 to OV-1) were used for retrograde statistical analysis of the ovulation-impending phase. A Shapiro-Wilk test was performed to visualize the normal distribution of the data. Mixed-model analysis for repeated measures (SAS PROC MIXED-Version 9.2 SAS Institute, Inc, Cary, NC) was used to compare the means of each response variable between groups and time and their interaction. Post hoc analyses were conducted using a Tukey test. No statistical differences between operators, among hours or an operator-by-hour interaction were

Download English Version:

<https://daneshyari.com/en/article/5523098>

Download Persian Version:

<https://daneshyari.com/article/5523098>

[Daneshyari.com](https://daneshyari.com)