



Short communication

Pathology and genetic findings in a rare case of *Mycobacterium caprae* infection in a sow

Benedetta Amato^a, Teresa Maria Capucchio^b, Elena Biasibetti^b, Elena Mangano^a,
Beatrice Maria Boniotti^c, Lodovica Maria Pacciarini^c, Sergio Migliore^a, Maria Vitale^{a,*},
Michele Fiasconaro^a, Vincenzo Di Marco Lo Presti^a

^a Istituto Zooprofilattico Sperimentale della Sicilia, via S. Andrea 96, Barcellona Pozzo di Gotto, ME, 98051, Italy

^b Dipartimento di Scienze Veterinarie – Università degli Studi di Torino, Largo Braccini 2, Grugliasco, TO 10195, Italy

^c National Reference Centre for Bovine Tuberculosis, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, via Bianchi 7/9, Brescia, 25124, Italy

ARTICLE INFO

Keywords:

Bovine tuberculosis
Mycobacterium caprae
Isolation from mammary glands
Rare spoligotype
Anatomo-pathological descriptions

ABSTRACT

Bovine tuberculosis, a reemerging zoonosis in diverse ecological scenarios, has been reported in the autochthonous Nebrodi black pig breed population used for meat production in Italy. During a routine abattoir inspection in 2013, 24 of 299 carcasses (8%) of Nebrodi black pigs presented tuberculosis-like lesions at pathologic examination. *Mycobacterium bovis* was isolated from 23 animals and *M. caprae* from a 3-year-old sow. The sow showed severe diffuse lesions involving the visceral organs, right coxofemoral joint, and mammary glands. Isolation of *M. caprae* from mammary glands is uncommon, with only one other case involving a sow reported so far; however, *Mycobacteria* infection of the mammary glands may be transmitted from lactating sows to piglets, contributing to the spread and maintenance of bovine tuberculosis in swine. Genotyping analysis showed *M. caprae* spoligotype SB0866 and profile 4,1,5,4,4,11,4,2,4,3,8,7 MIRU-VNTR (mycobacterial interspersed repetitive units-variable number of tandem repeats). The worldwide prevalence of this spoligotype is very low. The finding of severe, diffuse tuberculous lesions strongly suggests that Nebrodi black pigs are susceptible for *Mycobacterium* spp. and that they might act as a distributor for these microorganisms. Since natural ecosystems with multiple contacts among different livestock species and wild animals are very common in Mediterranean regions, current surveillance and eradication plans for bovine tuberculosis will need to be extended to other potential reservoir species in regions where extensive and traditional breeding systems are operated.

1. Introduction

Bovine tuberculosis (bTB) is a neglected, though reemerging zoonosis with broad implications for livestock production, agricultural trade, and public health (Menin et al., 2013). *Mycobacterium caprae*, a species within the *Mycobacterium tuberculosis* complex first identified in 2003 (Aranaz et al., 2003), has been isolated mainly in Europe and with increasing frequency in cattle in Spain (Rodríguez et al., 2011). A recent report on comparative genomics analysis suggests that *M. caprae* is one of the *M. bovis*-related mycobacteria that have evolved to infect not only goats and sheep but also other hosts such as wild boar, red deer, cattle, and humans, with high lesion scores (De La Fuente et al., 2015). Swine are receptive animals for infection by several species of *Mycobacterium* primarily through the oral route probably due to their omnivorous nature and habit of eating infected carrion (Gortazar et al.,

2003; Nugent et al., 2015). Though generally considered as bTB spillover hosts, there is evidence that wild boars might act as bTB reservoir hosts in Spain (Martin-Hernando et al., 2007; Vicente et al., 2006). Because domestic pigs and wild boars are known to be infected by *M. bovis* and

M. caprae and because both can infect a wide range of animal species, current eradication programs targeting only *M. bovis* in cattle have been expanded in many countries to address *M. caprae* infection in a manner similar to *M. bovis* (Rodríguez-Campos et al., 2014).

While the prevalence of bTB has gradually declined thanks to cattle health surveillance systems in many regions of Italy, controlling the disease remains a major problem in Sicily, particularly in natural areas like the Nebrodi National Park where there is extensive farming of mixed livestock and contact between domestic and wild animals is frequent. A recent outbreak of bTB has been reported in a fallow deer

* Corresponding author at: Istituto Zooprofilattico Sperimentale della Sicilia, Via Gino Marinuzzi 3, 90129 Palermo.
E-mail addresses: marvitus@yahoo.com, maria.vitale@izssicilia.it (M. Vitale).

(*Dama dama*) herd (Amato et al., 2016) that shares the same geographical area as the Nebrodi black pig (*Sus scrofa*), an autochthonous breed reared in free or semi-free roaming conditions. The pigs roam wooded areas for most of their lives, foraging mainly on acorns roots, bulbs, and fruit. They are captured shortly before slaughtering and are fattened by feeding with small amounts of grain, bran, and barley. Nebrodi black pig has been recently identified as an epidemiological reservoir for bTB (Di Marco et al., 2012). Here we report a rare case of *M. caprae* infection involving the mammary glands and other tissues of an adult sow. The anatomico-pathological features and the genotyping characterization of the isolated strain are described.

2. Materials and methods

An abattoir survey was carried out in the first half of 2013 in the province of Messina (Italy) to examine pathological signs of bTB infection in Nebrodi black pigs and to further address their role as a potential reservoir host for tuberculosis infection. The lymph nodes of the head and internal organs (lungs, liver, spleen, stomach, intestines, and respective lymph nodes) of 299 slaughtered Nebrodi black pigs were examined for macroscopic lesions. Lesions were sampled for tuberculosis suspicion when caseous, calcified lesions were observed. For the purposes of this study, additional histopathology and bacteriological analyses were performed to confirm the diagnosis in the cases of abnormal pathology findings. For microbiological examination, tissue samples were processed and cultured in liquid and solid media (Middlebrook 7H9 broth and Löwenstein-Jensen medium) according to the official culture protocol (OIE Manual, 2010). For histological examination, samples were fixed in 10% neutral buffered formalin and routinely processed to obtain 4- μ m sections stained with hematoxylin and eosin and Ziehl-Neelsen stain. Molecular characterization and genotyping of the isolates were performed by spoligotyping and mycobacterial interspersed repetitive units-variable number of tandem repeats (MIRU-VNTR) analysis (ETRA-E, VNTR 2163a, 2163b, 4052, 3155, 1895, 3232, and MIRU-26), as described elsewhere (Boniotti et al., 2009).

3. Results

Twenty-four of the 299 carcasses (8%) presented bTB-like lesions at gross examination. Macroscopic examination revealed localized granulomatous lesions affecting almost exclusively the lymph nodes of the head, thorax, and abdomen (submandibular, retropharyngeal, tracheobronchial, gastric, intestinal lymph nodes). Of particular interest were the severe generalized lesions of the tonsils, mammary glands, and right coxofemoral joint noted in a 3-year-old sow. The lesions of the lymph nodes of the head, thorax, and abdomen usually appeared as caseous necrotic-calcified tubercles (< 1 cm in diameter) or larger lesions (> 1 cm in diameter) encased in a thick fibrotic capsule. The lungs, liver, and spleen showed features consistent with protracted and disseminated tuberculosis. Small, translucent gray nodules with a pale yellow core, detectable in both surface and section, were observed in the tonsils. Multiple granulomas of different size containing yellowish-green exudate were present in three mammary glands (Fig. 1). A voluminous mass (10 cm) was observed in the right coxofemoral joint (Fig. 2). Transversal bone sections showed thickening of the joint capsule with extensive areas of necrosis and inflammation. X-ray examination revealed bone remodeling of the trochanter and ischial osteolysis (Fig. 3). Histological examination confirmed that lesions appeared as classic granulomas containing a variable number of multinucleated giant cells, macrophages, and lymphocytes, with minimal or abundant calcifications at the center. Occasional neutrophils in and around the necrotic core were observed, together with numerous acid-fast bacilli (Fig. 4). Four isolates obtained from the coxofemoral joint, lungs, iliac lymph nodes, and mammary glands were identified as *M. caprae*. MIRU-VNTR typing showed spoligotype SB0866 and MIRU-



Fig. 1. Multiple disseminated granulomas involving the mammary gland of a sow infected by *M. caprae*. A high-resolution version of this slide for use with the Virtual Microscope is available as eSlide: VM03976.



Fig. 2. Right coxofemoral joint of a sow infected by *M. caprae*. Granulomas with a caseous-necrotic center and severe fibroblastic reaction in the joint capsule. A high-resolution version of this slide for use with the Virtual Microscope is available as eSlide: VM03977.




Fig. 3. X-ray showing bone remodeling of the trochanter and ischial osteolysis in the right coxofemoral joint of a sow infected by *M. caprae*.

VNTR profile 4,1,5,4,4,11,4,2,4,3,8,7.

4. Discussion

The bTB infection rate is about 4% in cattle (Italian Reference Centre for Bovine Tuberculosis) and 7% in pigs in Sicily (Di Marco et al., 2012). In the Nebrodi Park ecosystem, traditional farming practices with extensive mixed breeding (cattle, pigs, sheep and goats, poultry, horses) can facilitate contact between infected and uninfected

Download English Version:

<https://daneshyari.com/en/article/5545401>

Download Persian Version:

<https://daneshyari.com/article/5545401>

[Daneshyari.com](https://daneshyari.com)