



## Full Length Article

## Taurine abated subacute dichlorvos toxicity

M.G. Akande\*, U.S. Ahmed

Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Abuja, Federal Capital Territory 900001, Nigeria



## ARTICLE INFO

## Keywords:

Taurine  
Dichlorvos  
Malondialdehyde  
Antioxidant enzymes  
Oxidative stress

## ABSTRACT

The aim of the study was to determine the effects of taurine in rats exposed to subacute dichlorvos toxicity. Fifty rats were weighed and assigned into five groups of ten rats each. The groups received: distilled water, soya oil (1 ml/kg), taurine (50 mg/kg), dichlorvos (10 mg/kg) and the combination treatment group received taurine first and then dichlorvos 30 min later. The treatments were administered once daily by oral gavage for 4 weeks. The rats were euthanized and blood samples were collected after the termination of the study. Serum samples were analysed for malondialdehyde concentration and activities of antioxidant enzymes (superoxide dismutase and catalase). Dichlorvos increased malondialdehyde concentration and reduced the activities of superoxide dismutase and catalase. There was attenuation of malondialdehyde concentration and improvement of activities of superoxide dismutase ( $P = 0.0273$ ) and catalase ( $P < 0.0001$ ) in rats treated with taurine. It is postulated that taurine ameliorated dichlorvos-induced oxidative stress through the reduction of malondialdehyde concentration and the enhancement of activities of antioxidant enzymes.

## 1. Introduction

Pesticides are chemicals that are widely used in agricultural, domestic and public health settings for the control of pests. There are evidences of the link between exposure to pesticides and the incidence of cancer, Parkinson's disease, Alzheimer's disease, multiple sclerosis, diabetes, aging, cardiovascular and chronic renal disease [1,2]. Dichlorvos (2,2-dichloro-vinyl dimethyl phosphate, DP) is an organophosphate insecticide that is used for the protection of greenhouse plants, fruits and vegetables against mushroom flies, aphids, spiders, mites, caterpillars, thrips and white flies [3]. The insecticide is commonly used on cattle, sheep, poultry and around livestock edifices for pest control [4]. It is applied as an anthelmintic in dogs, swine and horses [5]. In Nigeria, DP is widely used by farmers for agricultural pest control [6]. The routes of exposure to DP include ingestion, inhalation and dermal exposure [7]. It is known that organophosphorus compounds, including DP, elicit neurotoxicity in mammals by inhibiting acetylcholinesterase activity [8].

Oxidative stress has been identified as an important molecular mechanism of toxicity of DP [3]. It is defined as an imbalance between the production of free radicals and the antioxidant system of the body [9]. Enzymatic and non-enzymatic antioxidants scavenge free radicals and reactive oxygen species (ROS) in biological systems [10]. Taurine (2-aminoethanesulphonic acid, TA) is a conditionally essential amino acid with high water solubility and it is an end product of cysteine and methionine metabolism [11]. TA is an efficacious non-enzymatic

antioxidant that protects many organs in the body against toxicity, oxidative stress [12], and injury [13].

The purpose of the study was to determine whether taurine could protect male Wistar rats from the adverse effects of subacute dichlorvos toxicity. It is important to investigate the attenuation of the toxicity of DP with TA (a putative antioxidant) because oxidative stress has been identified as one of the important molecular mechanisms of DP toxicity. In addition, the protective effects of TA against DP toxicity should be explored because the residues of DP have been detected in food stuffs in several countries including Nigeria [14,15].

## 2. Methods

## 2.1. Experimental animals

The male Wistar rats used for this study were obtained from the Department of Biochemistry, Ahmadu Bello University, Zaria, Kaduna State, Nigeria. They were housed in cages in the Toxicology Laboratory of the Department of Veterinary Pharmacology and Toxicology, University of Abuja, Federal Capital Territory, Nigeria. The animals were housed under standard environmental conditions (25 °C, 12 h light/dark cycle) and had access to standard laboratory ration and tap water ad libitum. The Wistar rats were acclimatized for two weeks before the commencement of the study. The research was approved by the University of Abuja research ethics committee and the animals were handled in accordance with the guidelines of the National Institute of

\* Corresponding author.

E-mail address: [akande.ganiyat@uniabuja.edu.ng](mailto:akande.ganiyat@uniabuja.edu.ng) (M.G. Akande).

Health Guide for Care and Use of Laboratory animals [16].

## 2.2. Chemicals

Commercial grade dichlorvos (DP, 100% solution containing 100 g per liter of 2,3-dichlorovinyl dimethylphosphate) marketed as Sniper® was purchased from an agrochemical company in Abuja, Federal Capital Territory, Nigeria. DP was reconstituted in soya oil (SO, Grand Cereals and Oil Mills Limited, Jos, Nigeria) to a 5% stock solution. An analytical grade of taurine (TA, Chemical Abstract Service number: 107-35-7, purity  $\geq 99\%$ ) was obtained from Sigma Aldrich® (Steinheim, Germany). Prior to daily administration, 100 mg of TA was reconstituted in distilled water (DW) to obtain 100 mg/ml suspension.

## 2.3. Subacute toxicological study

The Wistar rats were weighed and divided randomly into five groups, with 10 rats in each group. Distilled water was given to the DW group, while the SO group was administered with soya oil (1 ml/kg). The TA group was treated with taurine only (50 mg/kg [17,18]), while the DP group received dichlorvos only (10 mg/kg, 1/7th LD<sub>50</sub>). The combination treatment group received TA first (50 mg/kg), and then DP (10 mg/kg) 30 min after the administration of TA. The treatments were administered once daily by oral gavage for 4 weeks. The rats were observed for clinical signs of toxicity during the study. At the end of the study, the rats were euthanized by decapitation [19] and 3 ml of blood samples were collected into centrifuge tubes. The blood samples were incubated at room temperature for 30 min and then centrifuged at  $1000 \times g$  for 5 min to obtain sera samples.

## 2.4. Determination of MDA concentration

The concentration of malondialdehyde (MDA) was evaluated in the serum. The method described by Draper and Hadley [20] was used. The principle of the method was based on the spectrophotometric measurement of the colour developed during the reaction of thiobarbituric acid with MDA. The solutions were cooled under tap water and the absorbance was measured with an ultraviolet (UV) spectrophotometer (T80+ UV/Visible Spectrophotometer®, PG Instruments Ltd., Leicestershire, United Kingdom) at 532 nm. The concentration of MDA in the samples was calculated by using the absorbance coefficient, MDA-TBA complex  $1.56 \times 10^5 \text{ cm}^{-1} \text{ M}^{-1}$ .

## 2.5. Assays of the activities of serum antioxidant enzymes

Superoxide dismutase (SOD) was assayed with the Northwest Life Science Specialties (NWLSTSM) SOD activity assay kit based on the method described by Martin et al. [21]. Catalase activity was measured by the NWLSTSM CAT activity assay kit according to the method described by Beers and Sizer [22]. The assay kits were purchased from Northwest Life Science Specialties, LLC, Vancouver, Washington, DC, United States of America.

## 2.6. Statistical analysis

The data obtained were expressed as mean  $\pm$  standard error of the mean. The biochemical parameters were analysed with one-way analysis of variance followed by Tukey's multiple comparison post hoc test. The statistical package used was Graphpad Prism version 4.00 for Windows (Graphpad software, San Diego, California, USA). Values of  $P < 0.05$  were considered statistically significant.




Fig. 1. Effects of treatments on serum malondialdehyde concentration. Data are expressed as mean  $\pm$  standard error of the mean,  $n = 10$  animals/group. DW: Distilled Water; SO: Soya Oil (1 ml/kg); TA: Taurine (50 mg/kg); DP: Dichlorvos (10 mg/kg); TA + DP: Taurine (50 mg/kg) + Dichlorvos (10 mg/kg).  $F_{4,45} = 1.506$ ;  $P = 0.2164$ .

## 3. Results

### 3.1. Effects of treatments on serum malondialdehyde concentration

The effects of the treatments on serum malondialdehyde (MDA) concentration are shown in Fig. 1. The highest serum MDA concentration was recorded in the DP group, while the lowest serum MDA concentration was observed in the TA + DP group.  $F_{4,45} = 1.506$ ;  $P = 0.2164$ .

### 3.2. Effects of treatments on serum superoxide dismutase activity

There was a significant increase ( $P < 0.05$ ) in the serum superoxide dismutase activity in the TA group compared to the SO group. This is illustrated in Fig. 2, where  $a = P < 0.05$ ; TA group versus SO group.  $F_{4,45} = 3.283$ ;  $P = 0.01$ .

### 3.3. Effects of treatments on serum catalase activity

The serum catalase (CAT) activity was significantly increased ( $P < 0.001$ ) in the TA group compared to those in the DW and SO groups respectively. This is depicted in Fig. 3, where  $a = p < 0.001$ , TA group versus DW and SO groups respectively. In addition, the serum CAT activity in the TA group was improved ( $P < 0.01$ ) relative to that in the DP group;  $b = p < 0.01$  TA group versus DP group. The serum CAT activity in the TA + DP group was elevated ( $P < 0.001$ ) compared to those in the DW and SO groups respectively;  $c = p < 0.001$  TA + DP group versus DW and SO groups respectively (Fig. 3). The




Fig. 2. Effects of treatments on serum superoxide dismutase activity. Data are expressed as mean  $\pm$  standard error of the mean,  $n = 10$  animals/group. DW: Distilled Water; SO: Soya Oil (1 ml/kg); TA: Taurine (50 mg/kg); DP: Dichlorvos (10 mg/kg); TA + DP: Taurine (50 mg/kg) + Dichlorvos (10 mg/kg).  $a = P < .05$  TA group versus SO group.  $F_{4,45} = 3.283$ ;  $P = 0.0191$ .

Download English Version:

<https://daneshyari.com/en/article/5558692>

Download Persian Version:

<https://daneshyari.com/article/5558692>

[Daneshyari.com](https://daneshyari.com)