



## Short communication

Antioxidant capacity of *trans*-resveratrol dietary supplements alone or combined with the mycotoxin beauvericin

Beatriz Mallebrera<sup>a,\*</sup>, Annalisa Maietti<sup>b</sup>, Paola Tedeschi<sup>b</sup>, Guillermina Font<sup>a</sup>,  
 Maria-Jose Ruiz<sup>a</sup>, Vincenzo Brandolini<sup>b</sup>

<sup>a</sup> Laboratory of Toxicology, Faculty of Pharmacy, University of Valencia, Av. Vicent Andres Estelles S/n, 46100 Burjassot, Valencia, Spain

<sup>b</sup> Department of Chemical and Pharmaceutical Sciences, University of Ferrara, Via Fossato di Mortara 17, 44121 Ferrara, Italy

## ARTICLE INFO

## Article history:

Received 20 January 2017

Received in revised form

24 March 2017

Accepted 21 April 2017

Available online 25 April 2017

## Keywords:

Dietary supplements

Beauvericin

Resveratrol

Antioxidant capacity

Photochemiluminescence

Capillary electrophoresis

## ABSTRACT

*Trans*-resveratrol (*trans*-RSV) is a polyphenol with multiples biological properties, such as anti-inflammatory, antioxidant, anti-aging, anti-diabetic, and antiplatelet. It occurs naturally in grapes and derivate, peanuts and berries. Beauvericin (BEA) is a mycotoxin present in cereals that produces cytotoxicity, intracellular reactive oxygen species and lipid peroxidation. The general objective of this research was to evaluate whether *trans*-RSV could be used as a good polyphenol against damages produced by BEA. Because *trans*-RSV can be ingested through dietary supplements, to reach this goal, the following specific objectives were proposed: to determine a) the *trans*-RSV content in different polyphenol dietary supplements by capillary electrophoresis, b) the antioxidant capacity of the *trans*-RSV in polyphenol supplements, and c) the influence of BEA in the antioxidant capacity of *trans*-RSV when they are in combination by photochemiluminescence assay. The results obtained in this study showed that all polyphenol dietary supplements present higher RSV content than the content of the label. The polyphenol supplements present antioxidant capacity. And the combination of *trans*-RSV and BEA did not affect the antioxidant capacity of *trans*-RSV. Thus, RSV could contribute to decrease oxidant effects produced by BEA.

© 2017 Elsevier Ltd. All rights reserved.

## 1. Introduction

Some oxidants are formed in response to physiological processes. A disturbance between pro-oxidants and antioxidants defense system in favor of the oxidants is defined as oxidative stress, which can contribute to the development of chronic disease and ageing process (Davies, 2000; Halliwell, 2006; Lobo et al., 2010; Rahal et al., 2014).

Antioxidant compounds ingested through diet can scavenge free radicals and protect the organisms from oxidative stress. More than 8000 compounds have been identified with antioxidant properties. Polyphenolic compounds are a great class of antioxidants. They include phenolic acids, flavonoids, stilbenes and lignans (Pandey and Rizvi, 2009). Resveratrol (3, 5, 4'-trihydroxystilbene; RSV) is a stilbene abundant in grapes and grape products such as wines and grape juice (Fernández-Mar et al., 2012). RSV exists in two

diastereomeric forms: *trans* and *cis* (Chen et al., 2007). *Trans*-RSV has biological properties such as antioxidant, anti-inflammatory, anti-aging and antiplatelet activities among others, which prevent several human diseases (Fernández-Mar et al., 2012; Li et al., 2012). This potential benefits resulted in increased consumption of *trans*-RSV supplements by several consumers. Many efforts have been made to provide a highly sensitive and selective analytical method for the determination and characterization of polyphenols in dietary supplements (Ignat et al., 2011). The polyphenol content has been determined in different food matrices by spectrophotometry (Camont et al., 2009), high-performance liquid chromatography (HPLC; Mark et al., 2005), gases chromatography (CG; Goldberg et al., 1995) and capillary electrophoresis (CE; Brandolini et al., 2002; Arribas et al., 2014; Gatea et al., 2015).

Moreover, antioxidant capacity can be determined by Trolox Equivalent Antioxidant Capacity (TEAC), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) method, oxygen radical absorbance capacity (ORAC), 1,1-diphenyl-2-picrylhydrazyl (DPPH), total radical-trapping antioxidant parameter (TRAP), ferric reducing antioxidant power (FRAP),

\* Corresponding author.

E-mail address: [beatriz.mallebrera@uv.es](mailto:beatriz.mallebrera@uv.es) (B. Mallebrera).

**Table 1**  
*Trans*-RSV content and components in the label different to *trans*-RSV in commercial dietary supplements samples.

Sample	<i>Trans</i> -RSV content in the label (mg/g)	Other compounds in the label
1	3.4	<i>Monascus purpureus</i> , octacosanol, vitamin B3, alpha lipoic acid, omega-3 fatty acid, chrome, pantothenic acid, vitamin B12 and folic acid.
2	1.23	Vitamin C (74.1 mg/g), vitamin E (12.3 mg/g), zinc, copper, selenium, omega-3 fatty acid, lutein and zeaxanthin
3	8.62	Soy isoflavones, vitamin K, vitamin D, quercetin (129.3 mg/g), calcium, magnesium and isovitexin.
4	Not declared	<i>Vitis vinifera</i> L. extract (290 mg/g), <i>Punica granatum</i> L. extract, selenium, vitamin C (26 mg/g), zinc and vitamin B2

photochemiluminescence (PCL) and thiobarbituric acid (TBA), among others (Prior et al., 2005; Alam et al., 2013).

Beauvericin (BEA) is a mycotoxin synthesized by many species of *Fusarium* fungi. BEA is a contaminant of cereals and product composed by cereals (Mahnine et al., 2011; Juan et al., 2012). It has been demonstrated that BEA is cytotoxic, decreases mitochondrial membrane potential, produces lipid peroxidation, DNA damage and cell death (Ruiz et al., 2011; Prosperini et al., 2013; Mallebrera et al., 2016), which could be related with oxidative stress produced by BEA in several cell lines (Ferrer et al., 2009; Prosperini et al., 2013; Mallebrera et al., 2015). Because of resveratrol has multiple biological properties and it can be ingested through dietary supplements, this could be used to mitigate the oxidative damage caused by beauvericin.

The general objective of this research was to evaluate whether *trans*-RSV could be used as a good polyphenol against damages produced by BEA. Because *trans*-RSV can be ingested through dietary supplements, to reach this goal, the following specific objectives were proposed: to determine a) *trans*-RSV content in dietary supplements by CE, b) the antioxidant capacity of *trans*-RSV in dietary supplements and c) the influence of BEA in the antioxidant capacity of *trans*-RSV when they are in combination with BEA by PCL.

## 2. Material and methods

### 2.1. Reagents

All reagents were purchased from Sigma-Aldrich (Milan, Italy). The luminol PCL assay was carried out using the Photochem<sup>®</sup> instrument with the ACL kit (Analytikjena, Jena, Germany).

### 2.2. Samples

Commercial samples of *trans*-RSV dietary supplements (n = 4) were collected during 2015 from different pharmacies in Italy. Table 1 shows the samples analyzed and content of *trans*-RSV in each of them according to the nutritional label.

### 2.3. Sample preparation

Briefly, 0.30 g of each sample, were extracted with 5 mL of methanol (MeOH) and mixed using a vortex every 5 min for 15 min. Then, it was centrifuged at 5000 rpm during 5 min and finally the supernatant were collected in a flask of 20 mL. This procedure was performed four times. Then, the extracts were completed to 20 mL with MeOH. Three independent extractions were performed for each sample.

### 2.4. Capillary electrophoresis

CE analyses were performed according to Brandolini et al. (2002) using a CE Beckman MDQ equipped with a diode array detector (Beckman, Fullerton, CA). The separation was obtained by

75 mm i.d. and 57 cm total length fused silica capillary column maintained in a cartridge with a detector window of 100  $\mu\text{m}$   $\times$  800  $\mu\text{m}$ . The capillary was conditioned before use by flushing 0.1 M NaOH for 1 min, then with water and, finally with buffer (20 mM Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub> and 50 mM PEG 400, 10% MeOH) for 3 min. The sample was injected into the capillary by pressure injection for 5 s. Separation was obtained at 25 kV and 25 °C for 15 min at 315 nm. After each separation the capillary was rinsed sequentially with NaOH 0.1M for 2 min and buffer analysis for 3 min. All analyses were performed in three independent assays. Data are analyzed using the Karat 32 software (Beckman Coulter, Fullerton, CA).

### 2.5. Antioxidant activity

#### 2.5.1. Antioxidant activity of *trans*-RSV dietary supplements

The antioxidant capacities of *trans*-RSV dietary supplements were determined using a PCL technique, namely, the luminol PCL assay. The determination was carried out using the Photochem<sup>®</sup> instrument with the ACL kit (Analytikjena, Jena, Germany), and following the procedure described by Popov and Lewin (1999). Two or three mL reagent 1 (solvent and dilution reagent), 200  $\mu\text{L}$  reagent 2 (buffer solution), 25  $\mu\text{L}$  reagent 3 (photosensitizer) and 10  $\mu\text{L}$  of standard or solution were mixed and measured. Trolox was used as standard to obtain a calibration curve (0.5–2 nM). The light emission curve was measured at  $\lambda_{\text{max}} = 350$  nm during 180 s, using the inhibition of superoxide anion radicals as the parameter to evaluate antioxidant effect. The antioxidant capacity was determined by using the area under the curve. The results were expressed as  $\mu\text{mol}$  Trolox equivalents (TEs) per g *trans*-RSV. Antioxidant capacity of supplements was determined replacing standard by diluted samples. Determinations were performed with 6 replicates of each sample.

#### 2.5.2. Antioxidant activity of *trans*-RSV when combined with BEA

Considering that *trans*-RSV possesses antioxidant properties and BEA increases ROS production, the *trans*-RSV antioxidant capacity against oxidant activity of BEA is an objective of interest. Thus, the antioxidant activity of *trans*-RSV, BEA and four combinations of BEA + *trans*-RSV with 1:2.5; 1:5; 2:1 and 1:1 ratio were determined using the PCL technique describe previously.

## 3. Results and discussion

During the last years, many *trans*-RSV dietary supplements have

**Table 2**  
*Trans*-RSV content (mg/g) in the dietary supplements analysed by CE. Data are expressed as mean  $\pm$  SD (n = 6).

Sample	Content (mg/g $\pm$ SD)
1	4.44 $\pm$ 0.52
2	2.41 $\pm$ 0.15
3	10.75 $\pm$ 0.48
4	24.79 $\pm$ 0.89

Download English Version:

<https://daneshyari.com/en/article/5560209>

Download Persian Version:

<https://daneshyari.com/article/5560209>

[Daneshyari.com](https://daneshyari.com)