



PREVENTION & REHABILITATION: ORIGINAL RESEARCH STUDY

The effects of strengthening exercises for wrist flexors and extensors on muscle strength and counter-stroke performance in amateur table tennis players



Raoyrin Chanavirut, Ph.D ^{a, b, *}, Nontawit Udompanich, BSc ^a, Phraophimon Udom, BSc ^a,
 Ponlapat Yonglitthipagon, PhD ^{a, b}, Wanida Donpunha, PhD ^a,
 Saowanee Nakmareong, PhD ^a, Juinichiro Yamauchi, PhD ^{c, d, e}

^a School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University, 123, Mittraphap Highway, Muang District, Khon Kaen, 40002, Thailand

^b Physical Therapy in Improving Functional Ability Group, Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, 40002, Thailand

^c Research Center in Back, Neck, Other Joint Pain and Human Performance (BNOJPH), Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, 40002, Thailand

^d Graduate School of Human Health Sciences, Tokyo Metropolitan University, Tokyo, Japan

^e Future Institute for Sport Sciences, Tokyo, Japan

ARTICLE INFO

Article history:

Received 8 September 2016

Received in revised form

31 January 2017

Accepted 27 February 2017

Keywords:

Strengthening exercise

Table tennis

Hand grip

Alternate counter test

ABSTRACT

This study aims to evaluate the effects of strengthening exercises on wrist flexor and extensor strength, hand grip strength, and counter-stroke performance. Thirty amateur table tennis players were recruited and randomly allocated into two groups: the control and the training group (n = 15/group). Pre- and post-data were collected. The training group performed home exercises for six weeks using a bucket filled with water, while the controls were asked to keep their lifestyle as usual. The general characteristics were no significant differences. The training group showed significantly higher levels of wrist flexor and extensor strength than the control group (p < 0.05). However, the difference in hand grip strength was not statistically significant. Both groups showed significant improvement in counter-stroke performance (p < 0.05), however, there was no difference between the groups. In conclusion, specific strengthening exercises increase wrist flexor and extensor strength, but they have no effect on either hand grip strength or counter-stroke performance.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

Muscle strength plays a crucial role in determining speed and power in athletic performance. Although an individual's performance is primarily determined by the specific skills, greater strength leads to a higher potential for athletic performance, especially for untrained individuals. Table tennis is a sport that requires rapid movements, and stroke speed is one of the keys to

success ([The International Table Tennis Federation Handbook, 2014–2015](#)). Exercises that strengthen wrist muscles can improve the stroke performance of table tennis players by increasing speed and power. A previous study shows hand grip strength is greater in the dominant hand versus the non-dominant hand in young top-level table tennis players ([Carrasco et al., 2010](#)). However, another study shows contradicting results in top-level female athletes ([Ivanovic and Dopsaj, 2012](#)). These opposite results may suggest the functional complexity of wrist and forearm muscles. A number of studies have been conducted on table tennis players' performance, including stroke performance. However, there are no reports about the effect of wrist-strengthening exercises on hand grip strength and counter-stroke performance. Strengthening exercises for table tennis players should be selected based on the specific movements involved, such as strokes. Therefore, this study aimed to evaluate the effect of strengthening exercises on wrist flexor and extensor

* Corresponding author. School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University, 123, Mittraphap Highway, Muang District, Khon Kaen, 40002, Thailand.

E-mail addresses: raocha@kku.ac.th (R. Chanavirut), nuinui_enzo@hotmail.com (N. Udompanich), phraophimon1992@gmail.com (P. Udom), ponlapat@kku.ac.th (P. Yonglitthipagon), waniadon@kku.ac.th (W. Donpunha), saowna@kku.ac.th (S. Nakmareong), yamauchi@tmu.ac.jp (J. Yamauchi).

strength, hand grip strength, and counter-stroke performance of amateur table tennis players.

2. Methods

2.1. Subjects

Thirty amateur table tennis players between the ages of 19 and 28 were recruited through an information poster at the university table tennis gymnasium. The subjects were randomly allocated into two groups: the control group ($n = 15$, CON) and the training group ($n = 15$, TR). A self-administered questionnaire was used to collect personal and medical history. All subjects were healthy and had no exclusionary criteria (hypertension, syncope, history of hand or forearm fracture, or previous surgery in the hand or forearm < 6 months). The study was approved by the Khon Kaen University Ethical Committee (HE582136), and written informed consent was obtained from each subject.

2.2. Experimental procedures

The subjects were randomly assigned using computer generated random numbers. Then a block randomization was employed to random the subjects into the TR and the CON groups. Wrist flexor and extensor strength, hand grip strength, and counter-stroke performance was measured for each subject before and after the six-week program using the alternate counter test. Subjects in the TR group were given a booklet in which the details inside explained wrist flexor and extensor strengthening exercise protocol at home and included the diary for recording the exercise session they performed. The subjects who had exercised adherence less than 90% were withdrawn from the study. Subjects in the CON group were asked to keep their lifestyle as usual during the six-week experiment. The protocol of this study shows in Fig. 1.

2.3. Measurements

2.3.1. Wrist flexor strength (WFS) and wrist extensor strength (WES)

WFS and WES were evaluated with a push-pull dynamometer (Baseline evaluation instruments® 250#). After 2–3 min of independent warm-up, the subjects were instructed to sit on a chair with forearm resting on a table and wrist and hand over the edge. The forearm position was pronated for wrist extensor and supinated for wrist flexor evaluation. Then, subjects were asked to extend or flex against the push pads as hard as possible when given the appropriate command. The procedure was performed two times; each contraction was held for 6 s and rest between trials was 2 min. The subjects were encouraged to breathe normally to avoid any increase in blood pressure. Then, the higher of the two force measurements was recorded.

2.3.2. Hand grip strength (HGS)

HGS was evaluated with a hand grip dynamometer (T.K.K. 5001 grip-a®). The subjects stood in an upright position with both arms resting in a natural posture alongside the body. The test hand held the dynamometer with the probe approximately 10 cm away from the body. The procedure was performed two times; rest between trials was 2 min. Then, the higher force was recorded.

2.3.3. Alternate counter test (ACT)

ACT was used to evaluate the counter-stroke performance of the table tennis players (Purashwani et al., 2010). After 5–10 min of warm-up and practice, the subjects were instructed to perform the alternate counter test. They were asked to make the number of rallies of alternate counter (one forehand and one backhand) with




Fig. 1. The flow diagram of this study.

the controller for a period of 30 s. The number of balls returned to the controller was counted by an observer. The procedure was performed two times; rest between trials was 5 min. The higher of the two numbers was recorded.

2.4. Exercise training intervention

The TR group was shown how to perform wrist flexor and extensor exercises at home with a bucket of water (Söderberg et al., 2012). The bucket weight for each subject was 60% of the 1-RM (Repetition Maximum: the maximum weight with which exactly one repetition of an exercise can be safely executed) of wrist flexor strength individually. In addition, the 1-RM was re-evaluated at the second and fourth week respectively. A scale with load capacity of up to 50 kg with graduation to 100 g precisely (SOEHNLE®, Germany) was used to weigh a bucket of water. After that, the water level was marked with a permanent marker to ensure the same water level for home exercise. Before exercise, subjects were instructed to warm-up for wrist flexor and extensor for 2 min. Then, they were instructed to perform the exercise in the same position used to evaluate wrist flexor and extensor strength. All subjects performed 8 times/set, 3 sets/day and 3 days/week. The resting period between the set was 2 min (Franklin and Balady, 2006). In addition, they were instructed to keep their diary to monitor their exercise adherence and received weekly phone calls to encourage the exercise.

2.5. Data analysis

Data analysis was performed using STATA-10.1. The Shapiro-Wilk test was used to evaluate the data normality. Then all parameters were analyzed with parametric statistics, using paired *t*-test for comparison within each group and ANCOVA for comparison between the groups, expressed as mean \pm SD. A value of $p < 0.05$ was considered to be statistically significant.

3. Results

There were no significant differences in general characteristics

Download English Version:

<https://daneshyari.com/en/article/5563956>

Download Persian Version:

<https://daneshyari.com/article/5563956>

[Daneshyari.com](https://daneshyari.com)