



Contents lists available at ScienceDirect

Journal of Exercise Science & Fitness

journal homepage: www.elsevier.com/locate/jesf

The reliability and validity of a video-based method for assessing hamstring strength in football players



Justin W.Y. Lee*, Cheng Li, Patrick S.H. Yung, Kai-Ming Chan

Department of Orthopaedics and Traumatology, Prince of Wales Hospital, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

ARTICLE INFO

Article history:

Received 22 February 2017

Received in revised form

20 April 2017

Accepted 21 April 2017

Keywords:

Strength

Performance

Musculoskeletal

Injury & prevention

ABSTRACT

Background/Objective: Evaluating hamstring strength by isokinetic dynamometry is limited by various practical issues such as time and cost. A video-based Nordic hamstring exercise is introduced as an alternative option. The aims of this study are to evaluate 1.) the between-session reliability and 2.) concurrent validity of the testing method compared to a standardized isokinetic dynamometry.

Methods: Thirty male elite footballers were recruited for the study. From the Nordic hamstring exercise, the video-analysis-determined Nordic break-point angles where the participant could no longer withstand the force of the fall (eccentric mode) and the number of seconds that the player could hold at 30° forward flexion angle (isometric mode) were measured. Intra-class correlation coefficients for between-session reliability, Pearson *r* correlations between the current method and isokinetic dynamometry were calculated.

Results: The reliability of the eccentric mode was moderate ($ICC_{(2,1)} = 0.82$) while that of isometric mode was poor ($ICC_{(2,1)} = 0.57$). The Nordic break-point angle of the eccentric mode significantly correlated with the concentric and eccentric hamstring peak torque ($r = 0.48$ and 0.58 , $p < 0.001$), while the isometric was not ($r = 0.02 - 0.07$, $p > 0.05$).

Conclusion: The eccentric mode of the video-based hamstring strength test was a moderately reliable and valid method to measure the eccentric hamstring strength in elite football players.

© 2017 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Hamstring strain injury (HSI) is the major musculoskeletal problem in team sports, accounting for 6% - 16% of overall injuries in professional football, rugby, Australian football, and baseball.^{1–6} In elite football, the HSIs accounting for more than one-third of all time-loss muscle injuries sustained and its training related injury rates have increased substantially over the past decade.⁷ Previous hamstring history generated a 2-fold higher risk of sustaining recurrent injuries. These recurrence rates ranged from 12% to 41% during the first year of returning to sports.^{2,8,9}

Understanding multifactorial risk factors and mechanisms of this injury are crucial for developing effective injury prevention measures.^{10,11} The majority of the HSI in elite football players occurs during high speed running.^{2,12,13} A large prospective cohort study on professional football players has shown lower hamstring

eccentric strength and lower quadriceps concentric strength as weak risk factors. Two studies on Australian football players have shown that low levels of hamstring strength in the preseason increased the risk for future HSI.^{14,15} However, the between-limb strength imbalance seems not a prominent risk factor.¹⁵ A meta-analysis study suggested that hamstring peak torque and hamstring-to-quadriceps (H/Q) ratio require further research to warrant their involvement in HSI because of inconclusive results, or small sample sizes.¹⁶

To date, isokinetic dynamometry had been widely adopted by previous prospective cohort studies investigating hamstring strength and strength ratios as risk factor of future HSI.^{17–19} However, this method is limited by various factors, including time, cost, and lack of portability of device.²⁰ Players may also experience a high level of delayed onset of muscle soreness after the test. Due to the above drawbacks, the clinical applicability of this test method in elite and amateur football population can be limited. Few practical field-based tests have been proposed to be the alternative measurement tools to determine hamstring injury risk. Measures

* Corresponding author.

E-mail address: justinlee@ort.cuhk.edu.hk (J.W.Y. Lee).

include maximal hamstring eccentric strength,²¹ hamstring strength endurance,¹⁴ and isometric hamstring strength.²² The usage of these tests may present some limitation, such as some techniques rely heavily on the skills and strength of the operator,^{20,23} while some strength measures' concurrent validity is yet to be determined.^{14,15,22} A previous study adopted the Nordic hamstring exercise as a field-based assessment of eccentric hamstring strength.²⁴ Their result has shown that the Nordic break-point angle that measured from a video-based motion analysis was related to eccentric hamstring peak torque.²⁴ This method allowed athletes to complete 3 trials of the test within 5 minutes. It seems to overcome many of the practical limitations of isokinetic testing. However, before using this test routinely, it is essential to investigate the reliability and concurrent validity among the elite football players.

This study aimed to evaluate 1) the between-session reliability and 2) concurrent validity of a video-based hamstring strength testing method compared to a traditional isokinetic hamstring strength testing method among semi-professional male football players.

2. Methods

2.1. Participants

A cross-sectional study design was used to determine the relationship between the performance of the video-based hamstring strength test and the isokinetic hamstrings muscle strength test among elite football players. Power analysis demonstrated that a sample of 29 was sufficient to detect an R value of 0.5.²⁴ Thirty male players (mean age = 20.4 ± 1.3 y; body weight = 64.3 ± 5.5 kg; height = 175.3 ± 5.2 cm) from 2nd division football league participated in this test. They were semi-professionals who had received training 5 times and participated in official matches once a week respectively. The exclusion criteria were a) Those had sustained a hamstring injury less than 6 months before testing. b) Those with prolonged hamstring problems, such as discomfort and inhibition during an athletic activity that hindered their participations in any tests in this study. Informed consent was signed and collected from each participant who were 18 years old or above. Informed assent was signed and gathered for athletes who were under the age of 18 years. All procedures described in this study were approved by the University clinical research ethics committee (Reference number: CRE - 2011.610 - T) and conducted in accordance with the Declaration of Helsinki.²⁵

2.2. Procedure and measurements

The video-based hamstring strength test and isokinetic strength tests have been done on two separate occasions 7 - 10 days apart. Based on Nordic hamstring exercise, the video-based hamstring strength test consisted of 2 modes: eccentric and isometric mode. Our test protocol was modified based on previous studies.^{9,24} Video demonstration and familiarization trials were provided. The two modes were tested in a randomized order. Each athlete started with a kneeling position on a yoga mat, and a tester stabilized the athlete's legs by pressing on his plantar-flexed ankles. In the eccentric mode, the athlete was instructed to lean forward in a slow, smooth, and controlled manner, with full extension of the back and hips, until he reached a maximum point where he could no longer withstand the force of the fall. Then, he was instructed to use arms and hands to buffer the landing (Fig. 1c). The Nordic break-point angle was defined as the angle between the line joining hip and knee markers and initial vertical position of each participant (Fig. 1b). The angle was determined by the first appearance of the

angular velocity that is greater than 10 °/s. The choice of 10 °/s as a cut-off point was based on our pilot study. In the pilot study, by visual inspection, the time that the participants could not withstand the force of fall during the Nordic hamstring exercise was estimated. By referring to the graph of angular velocity against time, it is observed that the angular velocity of 10 °/s was consistent to determine the Nordic break-point angle in the pilot study. In the isometric mode, the athlete was stabilized by pressing by tester 1 and held in front by tester 2 at 30°, which is the angle between the trunk and the vertical line that perpendicular to the floor. The result of isometric mode was the time (s) that athletes could maintain before the first appearance of the angular velocity of 10 °/s. Three trials of each mode were performed, and the best result was used in statistical analysis.

The athlete was videoed from the sagittal plane with a high-speed camera (Casio EX-F1, Japan) sampling at 300 frames per second. The camera set 3 m away and at 0.5 m from the floor. Three reflective circular markers were attached to the right greater trochanter, right lateral femoral condyle, and right lateral malleolus to calculate knee joint kinematics. Minimal clothing was recommended to avoid movement of markers. Video clips were digitized and transformed into a two-dimensional space using the motion analysis computer software (Ariel Performance Analysis System, Ariel Dynamics, Inc., CA, USA). All data were digitally filtered using a 4th order low-pass filter with a cutoff frequency of 5 Hz. The current motion analysis protocol was adopted from a previous study.²⁶

Gravity-corrected hamstring peak torque of the isokinetic hamstring strength test was used as the criterion for validity. Strenuous exercise was avoided 48 hours before the test. The athlete was instructed to have a 5-minute warm up by pedaling on a stationary bike and a 5-minute dynamic stretching on lower limb muscles. The measurement was taken by using a Biodex III dynamometer (Biodex Medical Systems, USA). This study adopted the testing protocol from Croisier study, which included the concentric performance of hamstring groups at 60 °/s (5 repetitions) and the eccentric performance of the hamstrings groups at 30 °/s (5 repetitions).¹⁷ The eccentric exertion of the hamstrings groups at 120 °/s was excluded from the protocol to minimize measurement error.¹⁷ Before each assessment, the participant performed four submaximal trials to familiarize himself with the protocol. All sets of testing were separated by a 1-minute rest. The tester provided verbal encouragement. The highest peak torque in Newton meters (N·m) for each measure was used. The both dominant and non-dominant legs were tested. Since there was no significant difference between the peak torques in both legs at all angular velocities ($p > 0.05$), the average peak torques were used for statistical analyses.

2.3. Data analysis

Descriptive data (means and SDs) for each variable, i.e. isokinetic hamstring peak torques and video-based strength test results were calculated. Normality of data was confirmed by the Shapiro-Wilks test. Intra-class correlation coefficient was used to examine the inter-test reliability (ICC_(2,1)) in both modes of the video-based hamstring strength test. The standard error of measurement (SEM) and minimal detectable difference (MDD) were calculated using the formulas standard deviation [(SD) × √(1 - ICC)] and (1.96 × SEM × √2) respectively.²⁷ According to previous guidelines, an ICC of 0.90 or greater was regarded as high, between 0.80 and 0.89 as moderate, and 0.79 or less as poor.²⁸ Criterion validity was calculated using the bivariate Pearson product-moment correlation coefficient (r). Correlation of the measurements from the video-based hamstring strength test and isokinetic dynamometry have been computed. The magnitudes of these correlations were described as trivial (0.0 - 0.1), small (0.1 - 0.3),

Download English Version:

<https://daneshyari.com/en/article/5579871>

Download Persian Version:

<https://daneshyari.com/article/5579871>

[Daneshyari.com](https://daneshyari.com)