Accepted Manuscript

Elevated modified shock index in early sepsis is associated with myocardial dysfunction and mortality



Namita Jayaprakash, Ognjen Gajic, Ryan D. Frank, Nathan Smischney

PII:	S0883-9441(17)30568-3
DOI:	doi: 10.1016/j.jcrc.2017.08.019
Reference:	YJCRC 52644

To appear in:

Revised date:###REVISEDDATE###Accepted date:###ACCEPTEDDATE###

Please cite this article as: Namita Jayaprakash, Ognjen Gajic, Ryan D. Frank, Nathan Smischney, Elevated modified shock index in early sepsis is associated with myocardial dysfunction and mortality, (2017), doi: 10.1016/j.jcrc.2017.08.019

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Page 1

Elevated modified shock index in early sepsis is associated with

myocardial dysfunction and mortality

Authors: Namita Jayaprakash MB BcH BAO, MRCEM^(a)*; Ognjen Gajic MD, MS^(a); Ryan D. Frank MS^(b); Nathan Smischney MD^(c)

^(a) Division of Pulmonary and Critical Care, Department of Internal Medicine, Mayo Clinic, 200

First Street SW, Rochester, MN 55905

^(b) Department Biomedical statistics and informatics, Mayo Clinic, 200 First Street SW,

Rochester, MN 55905

^(c) Department of Anesthesia/Critical Care Division, HEMAIR (HEModynamic and AIRway Management Group), Mayo Clinic, 200 First Street SW, Rochester, MN 55905

*Corresponding author

Division of Pulmonary and Critical Care Medicine

Mayo Clinic

200 1st Street SW

Rochester, MN 55905

Mobile: +1-586-533-9264

Download English Version:

https://daneshyari.com/en/article/5583177

Download Persian Version:

https://daneshyari.com/article/5583177

Daneshyari.com